

Cardinal Arithmetics

Grzegorz Bancerek
 Warsaw University
 Białystok

Summary. In the article addition, multiplication and power operation of cardinals are introduced. Presented are some properties of equipotence of Cartesian products, basic cardinal arithmetics laws (transformativity, associativity, distributivity), and some facts about finite sets.

MML Identifier: CARD_2.

The articles [12], [11], [7], [8], [3], [4], [5], [10], [2], [6], [9], and [1] provide the terminology and notation for this paper. For simplicity we follow a convention: A, B denote ordinal numbers, K, M, N denote cardinal numbers, x, x_1, x_2, y, y_1, y_2 are arbitrary, $X, Y, Z, X_1, X_2, Y_1, Y_2$ denote sets, and f denotes a function. Let us consider x . The functor $[x]$ yielding a set, is defined by:

$$[x] = x.$$

Next we state several propositions:

- (1) $[x] = x$.
- (2) $\overline{\overline{X}} \leq \overline{\overline{Y}}$ if and only if there exists f such that $X = f \circ Y$ or $X \subseteq f \circ Y$.
- (3) $\overline{\overline{f \circ X}} \leq \overline{\overline{X}}$.
- (4) If $\overline{\overline{X}} < \overline{\overline{Y}}$, then $Y \setminus X \neq \emptyset$.
- (5) If $x \in X$ and $X \approx Y$, then $Y \neq \emptyset$ and there exists x such that $x \in Y$.
- (6) $2^X \approx 2^{\overline{\overline{X}}}$ and $\overline{\overline{2^X}} = \overline{\overline{2^{\overline{\overline{X}}}}}$.
- (7) If $Z \in Y^X$, then $Z \approx X$ and $\overline{\overline{Z}} = \overline{\overline{X}}$.

We now define three new functors. Let us consider M, N . The functor $M + N$ yielding a cardinal number, is defined as follows:

$$M + N = \overline{\overline{\text{ord}(M) + \text{ord}(N)}}.$$

The functor $M \cdot N$ yielding a cardinal number, is defined by:

$$M \cdot N = \overline{\overline{\{M, N\}}}.$$

The functor M^N yielding a cardinal number, is defined by:

$$M^N = \overline{\overline{M^N}}.$$

Next we state a number of propositions:

- (8) $M + N = \overline{\text{ord}(M) + \text{ord}(N)}$.
- (9) $M \cdot N = \overline{[M, N]}$.
- (10) $M^N = \overline{\overline{M^N}}$.
- (11) $[X, Y] \approx [Y, X]$ and $\overline{[X, Y]} = \overline{[Y, X]}$.
- (12) $[[X, Y], Z] \approx [X, [Y, Z]]$ and $\overline{[[X, Y], Z]} = \overline{[X, [Y, Z]]}$.
- (13) $X \approx [X, \{x\}]$ and $\overline{X} = \overline{[X, \{x\}]}$.
- (14) (i) $[X, Y] \approx [\overline{X}, Y]$,
 (ii) $[X, Y] \approx [X, \overline{Y}]$,
 (iii) $[X, Y] \approx [\overline{X}, \overline{Y}]$,
 (iv) $\overline{[X, Y]} = \overline{[\overline{X}, Y]}$,
 (v) $\overline{[X, Y]} = \overline{[X, \overline{Y}]}$,
 (vi) $\overline{[X, Y]} = \overline{[\overline{X}, \overline{Y}]}$.
- (15) If $X_1 \approx Y_1$ and $X_2 \approx Y_2$, then $[X_1, X_2] \approx [Y_1, Y_2]$ and $\overline{[X_1, X_2]} = \overline{[Y_1, Y_2]}$.
- (16) If $x_1 \neq x_2$, then $A + B \approx [A, \{x_1\}] \cup [B, \{x_2\}]$ and $\overline{A + B} = \overline{[A, \{x_1\}] \cup [B, \{x_2\}]}$.
- (17) If $x_1 \neq x_2$, then $K + M \approx [K, \{x_1\}] \cup [M, \{x_2\}]$ and $K + M = \overline{[K, \{x_1\}] \cup [M, \{x_2\}]}$.
- (18) $A \cdot B \approx [A, B]$ and $\overline{A \cdot B} = \overline{[A, B]}$.

We now define three new functors. The cardinal number $\overline{0}$ is defined by:

$$\overline{0} = \overline{0}.$$

The cardinal number $\overline{1}$ is defined as follows:

$$\overline{1} = \overline{1}.$$

The cardinal number $\overline{2}$ is defined as follows:

$$\overline{2} = \text{succ } \overline{1}.$$

The following propositions are true:

- (19) $\overline{0} = \overline{0}$ and $\overline{1} = \overline{1}$ and $\overline{2} = \text{succ } \overline{1}$.
- (20) $\overline{0} = \mathbf{0}$ and $\overline{0} = \emptyset$ and $\overline{1} = \mathbf{1}$.
- (21) $\overline{0} = \overline{0}$ and $\overline{1} = \overline{1}$ and $\overline{2} = \overline{2}$.
- (22) $\overline{2} = \{\mathbf{0}, \mathbf{1}\}$ and $\overline{2} = \text{succ } \mathbf{1}$.
- (23) Suppose $X_1 \approx Y_1$ and $X_2 \approx Y_2$ and $x_1 \neq x_2$ and $y_1 \neq y_2$. Then $[X_1, \{x_1\}] \cup [X_2, \{x_2\}] \approx [Y_1, \{y_1\}] \cup [Y_2, \{y_2\}]$ and $\overline{[X_1, \{x_1\}] \cup [X_2, \{x_2\}]} = \overline{[Y_1, \{y_1\}] \cup [Y_2, \{y_2\}]}$.
- (24) $\overline{A + B} = \overline{A} + \overline{B}$.

- (25) $\overline{A \cdot B} = \overline{A} \cdot \overline{B}$.
- (26) $\{X, \{0\}\} \cup \{Y, \{1\}\} \approx \{Y, \{0\}\} \cup \{X, \{1\}\}$ and $\overline{\{X, \{0\}\} \cup \{Y, \{1\}\}} = \overline{\{Y, \{0\}\} \cup \{X, \{1\}\}}$.
- (27) $\{X_1, X_2\} \cup \{Y_1, Y_2\} \approx \{X_2, X_1\} \cup \{Y_2, Y_1\}$ and $\overline{\{X_1, X_2\} \cup \{Y_1, Y_2\}} = \overline{\{X_2, X_1\} \cup \{Y_2, Y_1\}}$.
- (28) If $x \neq y$, then $\overline{X} + \overline{Y} = \overline{\{X, \{x\}\} \cup \{Y, \{y\}\}}$.
- (29) $M + \overline{0} = M$ and $\overline{0} + M = M$.
- (30) $M + N = N + M$.
- (31) $(K + M) + N = K + (M + N)$.
- (32) $K \cdot \overline{0} = \overline{0}$ and $\overline{0} \cdot K = \overline{0}$.
- (33) $K \cdot \overline{1} = K$ and $\overline{1} \cdot K = K$.
- (34) $K \cdot M = M \cdot K$.
- (35) $(K \cdot M) \cdot N = K \cdot (M \cdot N)$.
- (36) $\overline{2} \cdot K = K + K$ and $K \cdot \overline{2} = K + K$.
- (37) $K \cdot (M + N) = K \cdot M + K \cdot N$ and $(M + N) \cdot K = M \cdot K + N \cdot K$.
- (38) $K^{\overline{0}} = \overline{1}$.
- (39) If $K \neq \overline{0}$, then $\overline{0}^K = \overline{0}$.
- (40) $K^{\overline{1}} = K$ and $\overline{1}^K = \overline{1}$.
- (41) $K^{M+N} = (K^M) \cdot (K^N)$.
- (42) $(K \cdot M)^N = (K^N) \cdot (M^N)$.
- (43) $K^{M \cdot N} = (K^M)^N$.
- (44) $\overline{2}^{\overline{X}} = \overline{2^X}$.
- (45) $K^{\overline{2}} = K \cdot K$.
- (46) $(K + M)^{\overline{2}} = (K \cdot K + (\overline{2} \cdot K) \cdot M) + M \cdot M$.
- (47) $\overline{X \cup Y} \leq \overline{X} + \overline{Y}$.
- (48) If $X \cap Y = \emptyset$, then $\overline{X \cup Y} = \overline{X} + \overline{Y}$.

In the sequel m, n will denote natural numbers. Next we state a number of propositions:

- (49) $\text{ord}(n + m) = \text{ord}(n) + \text{ord}(m)$.
- (50) $\text{ord}(n \cdot m) = \text{ord}(n) \cdot \text{ord}(m)$.
- (51) $\overline{n + m} = \overline{n} + \overline{m}$.
- (52) $\overline{n \cdot m} = \overline{n} \cdot \overline{m}$.
- (53) If X is finite and Y is finite and $X \cap Y = \emptyset$, then $\text{card}(X \cup Y) = \text{card } X + \text{card } Y$.
- (54) If X is finite and $x \notin X$, then $\text{card}(X \cup \{x\}) = \text{card } X + 1$.
- (55) If X is finite and Y is finite, then $\text{card } X = \text{card } Y$ if and only if $X \approx Y$.
- (56) If X is finite and Y is finite, then $\overline{X} = \overline{Y}$ if and only if $\text{card } X = \text{card } Y$.

- (57) If X is finite and Y is finite, then $\overline{\overline{X}} \leq \overline{\overline{Y}}$ if and only if $\text{card } X \leq \text{card } Y$.
- (58) If X is finite and Y is finite, then $\overline{\overline{X}} < \overline{\overline{Y}}$ if and only if $\text{card } X < \text{card } Y$.
- (59) If X is finite, then $X = \emptyset$ if and only if $\text{card } X = 0$.
- (60) If X is finite, then $\text{card } X = 1$ if and only if there exists x such that $X = \{x\}$.
- (61) If X is finite, then $X \approx \text{ord}(\text{card } X)$ and $X \approx \overline{\overline{\text{card } X}}$ and $X \approx \text{Seg}(\text{card } X)$.
- (62) If X is finite and Y is finite, then $\text{card}(X \cup Y) \leq \text{card } X + \text{card } Y$.
- (63) If $Y \subseteq X$ and X is finite, then $\text{card}(X \setminus Y) = \text{card } X - \text{card } Y$.
- (64) If X is finite and Y is finite, then $\text{card}(X \cup Y) = (\text{card } X + \text{card } Y) - \text{card}(X \cap Y)$.
- (65) If X is finite and Y is finite, then $\text{card}[X, Y] = \text{card } X \cdot \text{card } Y$.
- (66) If $X \subseteq Y$ and Y is finite, then $\text{card } X \leq \text{card } Y$.
- (67) If $X \subseteq Y$ and $X \neq Y$ and Y is finite, then $\text{card } X < \text{card } Y$ and $\overline{\overline{X}} < \overline{\overline{Y}}$.
- (68) If $\overline{\overline{X}} \leq \overline{\overline{Y}}$ or $\overline{\overline{X}} < \overline{\overline{Y}}$ but Y is finite, then X is finite.

In the sequel $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8$ are arbitrary. One can prove the following propositions:

- (69) $\text{card}\{x_1, x_2\} \leq 2$.
- (70) $\text{card}\{x_1, x_2, x_3\} \leq 3$.
- (71) $\text{card}\{x_1, x_2, x_3, x_4\} \leq 4$.
- (72) $\text{card}\{x_1, x_2, x_3, x_4, x_5\} \leq 5$.
- (73) $\text{card}\{x_1, x_2, x_3, x_4, x_5, x_6\} \leq 6$.
- (74) $\text{card}\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} \leq 7$.
- (75) $\text{card}\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} \leq 8$.
- (76) If $x_1 \neq x_2$, then $\text{card}\{x_1, x_2\} = 2$.
- (77) If $x_1 \neq x_2$ and $x_1 \neq x_3$ and $x_2 \neq x_3$, then $\text{card}\{x_1, x_2, x_3\} = 3$.
- (78) If $x_1 \neq x_2$ and $x_1 \neq x_3$ and $x_1 \neq x_4$ and $x_2 \neq x_3$ and $x_2 \neq x_3$ and $x_2 \neq x_4$ and $x_3 \neq x_4$, then $\text{card}\{x_1, x_2, x_3, x_4\} = 4$.

References

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.

- [4] Grzegorz Bancerek. Sequences of ordinal numbers. *Formalized Mathematics*, 1(2):281–290, 1990.
- [5] Grzegorz Bancerek. Zermelo theorem and axiom of choice. *Formalized Mathematics*, 1(2):265–267, 1990.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [10] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [11] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(1):25–34, 1990.
- [12] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.

Received March 6, 1990
