

# Real Exponents and Logarithms <sup>1</sup>

Konrad Raczkowski  
Warsaw University  
Białystok

Andrzej Nędzusiak  
Warsaw University  
Białystok

**Summary.** Definitions and properties of the following concepts: root, real exponent and logarithm. Also the number  $e$  is defined.

MML Identifier: POWER.

The papers [11], [2], [9], [1], [7], [5], [6], [13], [12], [4], [3], [8], and [10] provide the notation and terminology for this paper. For simplicity we follow the rules:  $a, b, c, d$  denote real numbers,  $m, n, m_1, m_2$  denote natural numbers,  $k, l$  denote integers, and  $p$  denotes a rational number. One can prove the following propositions:

- (1) If there exists  $m$  such that  $n = 2 \cdot m$ , then  $(-a)_{\mathbb{N}}^n = a_{\mathbb{N}}^n$ .
- (2) If there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $(-a)_{\mathbb{N}}^n = -a_{\mathbb{N}}^n$ .
- (3) If  $a \geq 0$  or there exists  $m$  such that  $n = 2 \cdot m$ , then  $a_{\mathbb{N}}^n \geq 0$ .

Let us consider  $n, a$ . The functor  $\sqrt[n]{a}$  yields a real number and is defined by:

- (Def.1) (i)  $\sqrt[n]{a} = \text{root}_n(a)$  if  $a \geq 0$  and  $n \geq 1$ ,  
(ii)  $\sqrt[n]{a} = -\text{root}_n(-a)$  if  $a < 0$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ .

One can prove the following propositions:

- (4) For all  $a, n$  holds if  $a \geq 0$  and  $n \geq 1$ , then  $\sqrt[n]{a} = \text{root}_n(a)$  but if  $a < 0$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a} = -\text{root}_n(-a)$ .
- (5) If  $n \geq 1$  and  $a \geq 0$  or there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a_{\mathbb{N}}^n} = a$  and  $\sqrt[n]{a_{\mathbb{N}}^n} = a$ .
- (6) If  $n \geq 1$ , then  $\sqrt[n]{0} = 0$ .
- (7) If  $n \geq 1$ , then  $\sqrt[n]{1} = 1$ .
- (8) If  $a \geq 0$  and  $n \geq 1$ , then  $\sqrt[n]{a} \geq 0$ .
- (9) If there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{-1} = -1$ .
- (10)  $\sqrt[1]{a} = a$ .

---

<sup>1</sup>Supported by RPBP-III.24.C8

- (11) If there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a} = -\sqrt[n]{-a}$ .
- (12) If  $n \geq 1$  and  $a \geq 0$  and  $b \geq 0$  or there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a \cdot b} = \sqrt[n]{a} \cdot \sqrt[n]{b}$ .
- (13) If  $a > 0$  and  $n \geq 1$  or  $a \neq 0$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{\frac{1}{a}} = \frac{1}{\sqrt[n]{a}}$ .
- (14) If  $a \geq 0$  and  $b > 0$  and  $n \geq 1$  or  $b \neq 0$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$ .
- (15) If  $a \geq 0$  and  $n \geq 1$  and  $m \geq 1$  or there exist  $m_1, m_2$  such that  $n = 2 \cdot m_1 + 1$  and  $m = 2 \cdot m_2 + 1$ , then  $\sqrt[n]{\sqrt[m]{a}} = \sqrt[n \cdot m]{a}$ .
- (16) If  $a \geq 0$  and  $n \geq 1$  and  $m \geq 1$  or there exist  $m_1, m_2$  such that  $n = 2 \cdot m_1 + 1$  and  $m = 2 \cdot m_2 + 1$ , then  $\sqrt[n]{a} \cdot \sqrt[m]{a} = \sqrt[n \cdot m]{a^{n+m}}$ .
- (17) If  $a \leq b$  but  $0 \leq a$  and  $n \geq 1$  or there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a} \leq \sqrt[n]{b}$ .
- (18) If  $a < b$  but  $a \geq 0$  and  $n \geq 1$  or there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a} < \sqrt[n]{b}$ .
- (19) If  $a \geq 1$  and  $n \geq 1$ , then  $\sqrt[n]{a} \geq 1$  and  $a \geq \sqrt[n]{a}$ .
- (20) If  $a \leq -1$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $\sqrt[n]{a} \leq -1$  and  $a \leq \sqrt[n]{a}$ .
- (21) If  $a \geq 0$  and  $a < 1$  and  $n \geq 1$ , then  $a \leq \sqrt[n]{a}$  and  $\sqrt[n]{a} < 1$ .
- (22) If  $a > -1$  and  $a \leq 0$  and there exists  $m$  such that  $n = 2 \cdot m + 1$ , then  $a \geq \sqrt[n]{a}$  and  $\sqrt[n]{a} > -1$ .
- (23) If  $a > 0$  and  $n \geq 1$ , then  $\sqrt[n]{a} - 1 \leq \frac{a-1}{n}$ .
- (24) For every sequence of real numbers  $s$  and for every  $a$  such that  $a > 0$  and for every  $n$  such that  $n \geq 1$  holds  $s(n) = \sqrt[n]{a}$  holds  $s$  is convergent and  $\lim s = 1$ .

Let us consider  $a, b$ . The functor  $a^b$  yielding a real number is defined as follows:

- (Def.2) (i)  $a^b = a^b_{\mathbb{R}}$  if  $a > 0$ ,  
(ii)  $a^b = 0$  if  $a = 0$  and  $b > 0$ ,  
(iii) there exists  $k$  such that  $k = b$  and  $a^b = a^k_{\mathbb{Z}}$  if  $a < 0$  and  $b$  is an integer.

One can prove the following propositions:

- (25) Given  $a, b$ . Then if  $a > 0$ , then  $a^b = a^b_{\mathbb{R}}$  but if  $a = 0$  and  $b > 0$ , then  $a^b = 0$  but if  $a < 0$  and  $b$  is an integer, then there exists  $k$  such that  $k = b$  and  $a^b = a^k_{\mathbb{Z}}$ .
- (26) If  $a > 0$ , then  $a^b = a^b_{\mathbb{R}}$ .
- (27) If  $b > 0$ , then  $0^b = 0$ .
- (28) If  $a < 0$ , then  $a^k = a^k_{\mathbb{Z}}$ .
- (29) If  $a \neq 0$ , then  $a^0 = 1$ .
- (30)  $a^1 = a$ .

- (31)  $1^a = 1$ .
- (32) If  $a > 0$ , then  $a^{b+c} = a^b \cdot a^c$ .
- (33) If  $a > 0$ , then  $a^{-c} = \frac{1}{a^c}$ .
- (34) If  $a > 0$ , then  $a^{b-c} = \frac{a^b}{a^c}$ .
- (35) If  $a > 0$  and  $b > 0$ , then  $(a \cdot b)^c = a^c \cdot b^c$ .
- (36) If  $a > 0$  and  $b > 0$ , then  $\frac{a^c}{b} = \frac{a^c}{b^c}$ .
- (37) If  $a > 0$ , then  $\frac{1}{a^b} = a^{-b}$ .
- (38) If  $a > 0$ , then  $(a^b)^c = a^{b \cdot c}$ .
- (39) If  $a > 0$ , then  $a^b > 0$ .
- (40) If  $a > 1$  and  $b > 0$ , then  $a^b > 1$ .
- (41) If  $a > 1$  and  $b < 0$ , then  $a^b < 1$ .
- (42) If  $a > 0$  and  $a < b$  and  $c > 0$ , then  $a^c < b^c$ .
- (43) If  $a > 0$  and  $a < b$  and  $c < 0$ , then  $a^c > b^c$ .
- (44) If  $a < b$  and  $c > 1$ , then  $c^a < c^b$ .
- (45) If  $a < b$  and  $c > 0$  and  $c < 1$ , then  $c^a > c^b$ .
- (46) If  $a \neq 0$ , then  $a^n = a_{\mathbb{N}}^n$ .
- (47) If  $n \geq 1$ , then  $a^n = a_{\mathbb{N}}^n$ .
- (48) If  $a \neq 0$ , then  $a^n = a^n$ .
- (49) If  $n \geq 1$ , then  $a^n = a^n$ .
- (50) If  $a \neq 0$ , then  $a^k = a_{\mathbb{Z}}^k$ .
- (51) If  $a > 0$ , then  $a^p = a_{\mathbb{Q}}^p$ .
- (52) If  $a \geq 0$  and  $n \geq 1$ , then  $a^{\frac{1}{n}} = \sqrt[n]{a}$ .
- (53)  $a^2 = a^2$ .
- (54) If  $a \neq 0$  and there exists  $l$  such that  $k = 2 \cdot l$ , then  $(-a)^k = a^k$ .
- (55) If  $a \neq 0$  and there exists  $l$  such that  $k = 2 \cdot l + 1$ , then  $(-a)^k = -a^k$ .

Next we state two propositions:

- (56) If  $-1 < a$ , then  $(1 + a)^n \geq 1 + n \cdot a$ .
- (57) If  $a > 0$  and  $a \neq 1$  and  $c \neq d$ , then  $a^c \neq a^d$ .

Let us consider  $a, b$ . Let us assume that  $a > 0$  and  $a \neq 1$  and  $b > 0$ . The functor  $\log_a b$  yields a real number and is defined by:

(Def.3)  $a^{\log_a b} = b$ .

The following propositions are true:

- (58) For all  $a, b, c$  such that  $a > 0$  and  $a \neq 1$  and  $b > 0$  holds  $c = \log_a b$  if and only if  $a^c = b$ .
- (59) If  $a > 0$  and  $a \neq 1$ , then  $\log_a 1 = 0$ .
- (60) If  $a > 0$  and  $a \neq 1$ , then  $\log_a a = 1$ .
- (61) If  $a > 0$  and  $a \neq 1$  and  $b > 0$  and  $c > 0$ , then  $\log_a b + \log_a c = \log_a (b \cdot c)$ .
- (62) If  $a > 0$  and  $a \neq 1$  and  $b > 0$  and  $c > 0$ , then  $\log_a b - \log_a c = \log_a \frac{b}{c}$ .

- (63) If  $a > 0$  and  $a \neq 1$  and  $b > 0$ , then  $\log_a(b^c) = c \cdot \log_a b$ .
- (64) If  $a > 0$  and  $a \neq 1$  and  $b > 0$  and  $b \neq 1$  and  $c > 0$ , then  $\log_a c = \log_a b \cdot \log_b c$ .
- (65) If  $a > 1$  and  $b > 0$  and  $c > b$ , then  $\log_a c > \log_a b$ .
- (66) If  $a > 0$  and  $a < 1$  and  $b > 0$  and  $c > b$ , then  $\log_a c < \log_a b$ .
- (67) For every sequence of real numbers  $s$  such that for every  $n$  holds  $s(n) = (1 + \frac{1}{n+1})^{n+1}$  holds  $s$  is convergent.

The real number  $e$  is defined as follows:

- (Def.4) for every sequence of real numbers  $s$  such that for every  $n$  holds  $s(n) = (1 + \frac{1}{n+1})^{n+1}$  holds  $e = \lim s$ .

## References

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [3] Andrzej Kondracki. Basic properties of rational numbers. *Formalized Mathematics*, 1(5):841–845, 1990.
- [4] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [5] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [6] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [7] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [8] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [9] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [10] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(1):125–130, 1991.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [12] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.
- [13] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.

*Received October 1, 1990*

---