

Real Function One-Side Differentiability

Ewa Burakowska
Warsaw University
Białystok

Beata Madras
Warsaw University
Białystok

Summary. We define real function one-side differentiability and one-side continuity. Main properties of one-side differentiability function are proved. Connections between one-side differential and differential real function at the point are demonstrated.

MML Identifier: `FDIFF_3`.

The terminology and notation used in this paper have been introduced in the following papers: [17], [2], [4], [1], [11], [5], [7], [14], [18], [3], [8], [9], [10], [16], [15], [12], [13], and [6]. For simplicity we follow the rules: h, h_1, h_2 are real sequences convergent to 0, c is a constant real sequence, f, f_1, f_2 are partial functions from \mathbb{R} to \mathbb{R} , x_0, r, r_1, g, g_1, g_2 are real numbers, n is a natural number, and a is a sequence of real numbers. The following propositions are true:

- (1) If there exists r such that $r > 0$ and $[x_0 - r, x_0] \subseteq \text{dom } f$, then there exist h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) < 0$.
- (2) If there exists r such that $r > 0$ and $[x_0, x_0 + r] \subseteq \text{dom } f$, then there exist h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) > 0$.
- (3) Suppose For all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) < 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent and $\{x_0\} \subseteq \text{dom } f$. Given h_1, h_2, c . Suppose $\text{rng } c = \{x_0\}$ and $\text{rng}(h_1 + c) \subseteq \text{dom } f$ and for every n holds $h_1(n) < 0$ and $\text{rng}(h_2 + c) \subseteq \text{dom } f$ and for every n holds $h_2(n) < 0$. Then $\lim(h_1^{-1}(f \cdot (h_1 + c) - f \cdot c)) = \lim(h_2^{-1}(f \cdot (h_2 + c) - f \cdot c))$.
- (4) Suppose For all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) > 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent and $\{x_0\} \subseteq \text{dom } f$. Given h_1, h_2, c . Suppose $\text{rng } c = \{x_0\}$ and $\text{rng}(h_1 + c) \subseteq$

$\text{dom } f$ and $\text{rng}(h_2 + c) \subseteq \text{dom } f$ and for every n holds $h_1(n) > 0$ and for every n holds $h_2(n) > 0$. Then $\lim(h_1^{-1}(f \cdot (h_1 + c) - f \cdot c)) = \lim(h_2^{-1}(f \cdot (h_2 + c) - f \cdot c))$.

We now define four new predicates. Let us consider f, x_0 . We say that f is left continuous in x_0 if and only if:

(Def.1) $x_0 \in \text{dom } f$ and for every a such that $\text{rng } a \subseteq]-\infty, x_0[\cap \text{dom } f$ and a is convergent and $\lim a = x_0$ holds $f \cdot a$ is convergent and $f(x_0) = \lim(f \cdot a)$.

We say that f is right continuous in x_0 if and only if:

(Def.2) $x_0 \in \text{dom } f$ and for every a such that $\text{rng } a \subseteq]x_0, +\infty[\cap \text{dom } f$ and a is convergent and $\lim a = x_0$ holds $f \cdot a$ is convergent and $f(x_0) = \lim(f \cdot a)$.

We say that f is right differentiable in x_0 if and only if the conditions (Def.3) is satisfied.

(Def.3) (i) There exists r such that $r > 0$ and $[x_0, x_0 + r] \subseteq \text{dom } f$,
(ii) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) > 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent.

We say that f is left differentiable in x_0 if and only if the conditions (Def.4) is satisfied.

(Def.4) (i) There exists r such that $r > 0$ and $[x_0 - r, x_0] \subseteq \text{dom } f$,
(ii) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) < 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent.

One can prove the following propositions:

- (5) If f is left differentiable in x_0 , then f is left continuous in x_0 .
- (6) Suppose f is left continuous in x_0 and $f(x_0) \neq g_2$ and there exists r such that $r > 0$ and $[x_0 - r, x_0] \subseteq \text{dom } f$. Then there exists r_1 such that $r_1 > 0$ and $[x_0 - r_1, x_0] \subseteq \text{dom } f$ and for every g such that $g \in [x_0 - r_1, x_0]$ holds $f(g) \neq g_2$.
- (7) If f is right differentiable in x_0 , then f is right continuous in x_0 .
- (8) Suppose f is right continuous in x_0 and $f(x_0) \neq g_2$ and there exists r such that $r > 0$ and $[x_0, x_0 + r] \subseteq \text{dom } f$. Then there exists r_1 such that $r_1 > 0$ and $[x_0, x_0 + r_1] \subseteq \text{dom } f$ and for every g such that $g \in [x_0, x_0 + r_1]$ holds $f(g) \neq g_2$.

Let us consider x_0, f . Let us assume that f is left differentiable in x_0 . The functor $f'_-(x_0)$ yielding a real number is defined by:

(Def.5) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) < 0$ holds $f'_-(x_0) = \lim(h^{-1}(f \cdot (h + c) - f \cdot c))$.

Let us consider x_0, f . Let us assume that f is right differentiable in x_0 . The functor $f'_+(x_0)$ yields a real number and is defined by:

(Def.6) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) > 0$ holds $f'_+(x_0) = \lim(h^{-1}(f \cdot (h + c) - f \cdot c))$.

The following propositions are true:

- (9) f is left differentiable in x_0 and $f'_-(x_0) = g$ if and only if the following conditions are satisfied:
- (i) there exists r such that $0 < r$ and $[x_0 - r, x_0] \subseteq \text{dom } f$,
 - (ii) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) < 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent and $\lim(h^{-1}(f \cdot (h + c) - f \cdot c)) = g$.
- (10) If f_1 is left differentiable in x_0 and f_2 is left differentiable in x_0 , then $f_1 + f_2$ is left differentiable in x_0 and $(f_1 + f_2)'_-(x_0) = f_{1-}'(x_0) + f_{2-}'(x_0)$.
- (11) If f_1 is left differentiable in x_0 and f_2 is left differentiable in x_0 , then $f_1 - f_2$ is left differentiable in x_0 and $(f_1 - f_2)'_-(x_0) = f_{1-}'(x_0) - f_{2-}'(x_0)$.
- (12) If f_1 is left differentiable in x_0 and f_2 is left differentiable in x_0 , then $f_1 f_2$ is left differentiable in x_0 and $(f_1 f_2)'_-(x_0) = f_{1-}'(x_0) \cdot f_2(x_0) + f_{2-}'(x_0) \cdot f_1(x_0)$.
- (13) If f_1 is left differentiable in x_0 and f_2 is left differentiable in x_0 and $f_2(x_0) \neq 0$, then $\frac{f_1}{f_2}$ is left differentiable in x_0 and
- $$\left(\frac{f_1}{f_2}\right)'_-(x_0) = \frac{f_{1-}'(x_0) \cdot f_2(x_0) - f_{2-}'(x_0) \cdot f_1(x_0)}{f_2(x_0)^2}.$$
- (14) If f is left differentiable in x_0 and $f(x_0) \neq 0$, then $\frac{1}{f}$ is left differentiable in x_0 and $\left(\frac{1}{f}\right)'_-(x_0) = -\frac{f'_-(x_0)}{f(x_0)^2}$.
- (15) f is right differentiable in x_0 and $f'_+(x_0) = g_1$ if and only if the following conditions are satisfied:
- (i) there exists r such that $r > 0$ and $[x_0, x_0 + r] \subseteq \text{dom } f$,
 - (ii) for all h, c such that $\text{rng } c = \{x_0\}$ and $\text{rng}(h + c) \subseteq \text{dom } f$ and for every n holds $h(n) > 0$ holds $h^{-1}(f \cdot (h + c) - f \cdot c)$ is convergent and $\lim(h^{-1}(f \cdot (h + c) - f \cdot c)) = g_1$.
- (16) If f_1 is right differentiable in x_0 and f_2 is right differentiable in x_0 , then $f_1 + f_2$ is right differentiable in x_0 and $(f_1 + f_2)'_+(x_0) = f_{1+}'(x_0) + f_{2+}'(x_0)$.
- (17) If f_1 is right differentiable in x_0 and f_2 is right differentiable in x_0 , then $f_1 - f_2$ is right differentiable in x_0 and $(f_1 - f_2)'_+(x_0) = f_{1+}'(x_0) - f_{2+}'(x_0)$.
- (18) If f_1 is right differentiable in x_0 and f_2 is right differentiable in x_0 , then $f_1 f_2$ is right differentiable in x_0 and $(f_1 f_2)'_+(x_0) = f_{1+}'(x_0) \cdot f_2(x_0) + f_{2+}'(x_0) \cdot f_1(x_0)$.
- (19) If f_1 is right differentiable in x_0 and f_2 is right differentiable in x_0 and $f_2(x_0) \neq 0$, then $\frac{f_1}{f_2}$ is right differentiable in x_0 and $\left(\frac{f_1}{f_2}\right)'_+(x_0) = \frac{f_{1+}'(x_0) \cdot f_2(x_0) - f_{2+}'(x_0) \cdot f_1(x_0)}{f_2(x_0)^2}$.
- (20) If f is right differentiable in x_0 and $f(x_0) \neq 0$, then $\frac{1}{f}$ is right differentiable in x_0 and $\left(\frac{1}{f}\right)'_+(x_0) = -\frac{f'_+(x_0)}{f(x_0)^2}$.
- (21) If f is right differentiable in x_0 and f is left differentiable in x_0 and $f'_+(x_0) = f'_-(x_0)$, then f is differentiable in x_0 and $f'(x_0) = f'_+(x_0)$ and $f'(x_0) = f'_-(x_0)$.

- (22) If f is differentiable in x_0 , then f is right differentiable in x_0 and f is left differentiable in x_0 and $f'(x_0) = f'_+(x_0)$ and $f'(x_0) = f'_-(x_0)$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [3] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [4] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [5] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [6] Jarosław Kotowicz. The limit of a real function at infinity. *Formalized Mathematics*, 2(1):17–28, 1991.
- [7] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [8] Jarosław Kotowicz. Partial functions from a domain to a domain. *Formalized Mathematics*, 1(4):697–702, 1990.
- [9] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(4):703–709, 1990.
- [10] Jarosław Kotowicz. Properties of real functions. *Formalized Mathematics*, 1(4):781–786, 1990.
- [11] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [12] Jarosław Kotowicz and Konrad Raczkowski. Real function differentiability - Part II. *Formalized Mathematics*, 2(3):407–411, 1991.
- [13] Andrzej Nędzusiak. σ -fields and probability. *Formalized Mathematics*, 1(2):401–407, 1990.
- [14] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [15] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(4):797–801, 1990.
- [16] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [17] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [18] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.

Received December 12, 1991
