

Sequences in \mathcal{E}_T^N

Agnieszka Sakowicz
 Warsaw University
 Białystok

Jarosław Gryko
 Warsaw University
 Białystok

Adam Grabowski
 Warsaw University
 Białystok

MML Identifier: TOPRNS_1.

The papers [12], [3], [4], [11], [8], [10], [1], [2], [5], [6], [9], and [7] provide the notation and terminology for this paper.

For simplicity we adopt the following rules: f denotes a function, N, n, m denote natural numbers, q, r, r_1, r_2 denote real numbers, x is arbitrary, and w, w_1, w_2, g denote points of \mathcal{E}_T^N .

Let us consider N . A sequence in \mathcal{E}_T^N is a function from \mathbb{N} into the carrier of \mathcal{E}_T^N .

In the sequel s_1, s_2, s_3, s_4, s'_1 are sequences in \mathcal{E}_T^N .

Next we state two propositions:

- (1) f is a sequence in \mathcal{E}_T^N if and only if $\text{dom } f = \mathbb{N}$ and for every x such that $x \in \mathbb{N}$ holds $f(x)$ is a point of \mathcal{E}_T^N .
- (2) f is a sequence in \mathcal{E}_T^N iff $\text{dom } f = \mathbb{N}$ and for every n holds $f(n)$ is a point of \mathcal{E}_T^N .

Let us consider N, s_1, n . Then $s_1(n)$ is a point of \mathcal{E}_T^N .

Let us consider N . A sequence in \mathcal{E}_T^N is non-zero if:

(Def.1) $\text{rng it} \subseteq (\text{the carrier of } \mathcal{E}_T^N) \setminus \{0_{\mathcal{E}_T^N}\}$.

We now state several propositions:

- (3) s_1 is non-zero iff for every x such that $x \in \mathbb{N}$ holds $s_1(x) \neq 0_{\mathcal{E}_T^N}$.
- (4) s_1 is non-zero iff for every n holds $s_1(n) \neq 0_{\mathcal{E}_T^N}$.
- (5) For all N, s_1, s_2 such that for every x such that $x \in \mathbb{N}$ holds $s_1(x) = s_2(x)$ holds $s_1 = s_2$.
- (6) For all N, s_1, s_2 such that for every n holds $s_1(n) = s_2(n)$ holds $s_1 = s_2$.
- (7) For every point w of \mathcal{E}_T^N there exists s_1 such that $\text{rng } s_1 = \{w\}$.

The scheme *ExTopRealNSeq* deals with a natural number \mathcal{A} and a unary functor \mathcal{F} yielding a point of $\mathcal{E}_T^{\mathcal{A}}$, and states that:

There exists a sequence s_1 in \mathcal{E}_T^A such that for every n holds $s_1(n) = \mathcal{F}(n)$

for all values of the parameters.

Let us consider N, s_2, s_3 . The functor $s_2 + s_3$ yielding a sequence in \mathcal{E}_T^N is defined by:

(Def.2) For every n holds $(s_2 + s_3)(n) = s_2(n) + s_3(n)$.

Let us consider r, N, s_1 . The functor $r \cdot s_1$ yields a sequence in \mathcal{E}_T^N and is defined by:

(Def.3) For every n holds $(r \cdot s_1)(n) = r \cdot s_1(n)$.

Let us consider N, s_1 . The functor $-s_1$ yields a sequence in \mathcal{E}_T^N and is defined as follows:

(Def.4) For every n holds $(-s_1)(n) = -s_1(n)$.

Let us consider N, s_2, s_3 . The functor $s_2 - s_3$ yields a sequence in \mathcal{E}_T^N and is defined by:

(Def.5) $s_2 - s_3 = s_2 + -s_3$.

Let us consider N and let x be a point of \mathcal{E}_T^N . The functor $|x|$ yields a real number and is defined by:

(Def.6) There exists a finite sequence y of elements of \mathbb{R} such that $x = y$ and $|x| = |y|$.

Let us consider N, s_1 . The functor $|s_1|$ yielding a sequence of real numbers is defined by:

(Def.7) For every n holds $|s_1|(n) = |s_1(n)|$.

We now state a number of propositions:

$$(8) \quad |r| \cdot |w| = |r \cdot w|.$$

$$(9) \quad |r \cdot s_1| = |r| |s_1|.$$

$$(10) \quad s_2 + s_3 = s_3 + s_2.$$

$$(11) \quad (s_2 + s_3) + s_4 = s_2 + (s_3 + s_4).$$

$$(12) \quad -s_1 = (-1) \cdot s_1.$$

$$(13) \quad r \cdot (s_2 + s_3) = r \cdot s_2 + r \cdot s_3.$$

$$(14) \quad (r \cdot q) \cdot s_1 = r \cdot (q \cdot s_1).$$

$$(15) \quad r \cdot (s_2 - s_3) = r \cdot s_2 - r \cdot s_3.$$

$$(16) \quad s_2 - (s_3 + s_4) = s_2 - s_3 - s_4.$$

$$(17) \quad 1 \cdot s_1 = s_1.$$

$$(18) \quad --s_1 = s_1.$$

$$(19) \quad s_2 - -s_3 = s_2 + s_3.$$

$$(20) \quad s_2 - (s_3 - s_4) = (s_2 - s_3) + s_4.$$

$$(21) \quad s_2 + (s_3 - s_4) = (s_2 + s_3) - s_4.$$

(22) If $r \neq 0$ and s_1 is non-zero, then $r \cdot s_1$ is non-zero.

(23) If s_1 is non-zero, then $-s_1$ is non-zero.

$$(24) \quad |0_{\mathcal{E}_T^N}| = 0.$$

- (25) If $|w| = 0$, then $w = 0_{\mathcal{E}_T^N}$.
- (26) $|w| \geq 0$.
- (27) $|-w| = |w|$.
- (28) $|w_1 - w_2| = |w_2 - w_1|$.
- (29) $|w_1 - w_2| = 0$ iff $w_1 = w_2$.
- (30) $|w_1 + w_2| \leq |w_1| + |w_2|$.
- (31) $|w_1 - w_2| \leq |w_1| + |w_2|$.
- (32) $|w_1| - |w_2| \leq |w_1 + w_2|$.
- (33) $|w_1| - |w_2| \leq |w_1 - w_2|$.
- (34) If $w_1 \neq w_2$, then $|w_1 - w_2| > 0$.
- (35) $|w_1 - w_2| \leq |w_1 - w| + |w - w_2|$.
- (36) If $0 \leq |w_1|$ and $0 \leq r_1$ and $|w_1| < |w_2|$ and $r_1 < r_2$, then $|w_1| \cdot r_1 < |w_2| \cdot r_2$.
- (38)¹ $-|w| < r$ and $r < |w|$ iff $|r| < |w|$.

Let us consider N . A sequence in \mathcal{E}_T^N is bounded if:

- (Def.8) There exists r such that for every n holds $|\text{it}(n)| < r$.

The following proposition is true

- (39) For every n there exists r such that $0 < r$ and for every m such that $m \leq n$ holds $|s_1(m)| < r$.

Let us consider N . A sequence in \mathcal{E}_T^N is convergent if:

- (Def.9) There exists g such that for every r such that $0 < r$ there exists n such that for every m such that $n \leq m$ holds $|\text{it}(m) - g| < r$.

Let us consider N , s_1 . Let us assume that s_1 is convergent. The functor $\lim s_1$ yields a point of \mathcal{E}_T^N and is defined by:

- (Def.10) For every r such that $0 < r$ there exists n such that for every m such that $n \leq m$ holds $|s_1(m) - \lim s_1| < r$.

The following propositions are true:

- (40) Suppose s_1 is convergent. Then $\lim s_1 = g$ if and only if for every r such that $0 < r$ there exists n such that for every m such that $n \leq m$ holds $|s_1(m) - g| < r$.
- (41) If s_1 is convergent and s'_1 is convergent, then $s_1 + s'_1$ is convergent.
- (42) If s_1 is convergent and s'_1 is convergent, then $\lim(s_1 + s'_1) = \lim s_1 + \lim s'_1$.
- (43) If s_1 is convergent, then $r \cdot s_1$ is convergent.
- (44) If s_1 is convergent, then $\lim(r \cdot s_1) = r \cdot \lim s_1$.
- (45) If s_1 is convergent, then $-s_1$ is convergent.
- (46) If s_1 is convergent, then $\lim(-s_1) = -\lim s_1$.
- (47) If s_1 is convergent and s'_1 is convergent, then $s_1 - s'_1$ is convergent.

¹The proposition (37) has been removed.

- (48) If s_1 is convergent and s'_1 is convergent, then $\lim(s_1 - s'_1) = \lim s_1 - \lim s'_1$.
- (50)² If s_1 is convergent, then s_1 is bounded.
- (51) If s_1 is convergent, then if $\lim s_1 \neq 0_{\mathcal{E}_T^N}$, then there exists n such that for every m such that $n \leq m$ holds $\frac{|\lim s_1|}{2} < |s_1(m)|$.

REFERENCES

- [1] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [2] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [6] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [7] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [8] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [9] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [10] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [12] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.

Received May 10, 1994

²The proposition (49) has been removed.