

On Kolmogorov Topological Spaces ¹

Zbigniew Karno
Warsaw University
Białystok

Summary. Let X be a topological space. X is said to be T_0 -space (or *Kolmogorov space*) provided for every pair of distinct points $x, y \in X$ there exists an open subset of X containing exactly one of these points; equivalently, for every pair of distinct points $x, y \in X$ there exists a closed subset of X containing exactly one of these points (see [1], [6], [2]).

The purpose is to list some of the standard facts on Kolmogorov spaces, using Mizar formalism. As a sample we formulate the following characteristics of such spaces: *X is a Kolmogorov space iff for every pair of distinct points $x, y \in X$ the closures $\overline{\{x\}}$ and $\overline{\{y\}}$ are distinct.*

There is also reviewed analogous facts on Kolmogorov subspaces of topological spaces. In the presented approach T_0 -subsets are introduced and some of their properties developed.

MML Identifier: TSP_1.

The articles [10], [11], [9], [7], [8], [5], [4], and [3] provide the terminology and notation for this paper.

1. SUBSPACES

Let Y be a non empty topological structure. We see that the subspace of Y is a non empty topological structure and it can be characterized by the following (equivalent) condition:

- (Def.1) (i) The carrier of it \subseteq the carrier of Y , and
(ii) for every subset G_0 of it holds G_0 is open iff there exists a subset G of Y such that G is open and $G_0 = G \cap$ (the carrier of it).

Next we state two propositions:

¹Presented at Mizar Conference: Mathematics in Mizar (Białystok, September 12–14, 1994).

- (1) Let Y be a non empty topological structure, and let Y_0 be a subspace of Y , and let G_0 be a subset of Y_0 . Then G_0 is open if and only if there exists a subset G of Y such that G is open and $G_0 = G \cap$ (the carrier of Y_0).
- (2) Let Y be a non empty topological structure, and let Y_0 be a subspace of Y , and let G be a subset of Y . Suppose G is open. Then there exists a subset G_0 of Y_0 such that G_0 is open and $G_0 = G \cap$ (the carrier of Y_0).

Let Y be a non empty topological structure. We see that the subspace of Y is a non empty topological structure and it can be characterized by the following (equivalent) condition:

- (Def.2) (i) The carrier of it \subseteq the carrier of Y , and
(ii) for every subset F_0 of it holds F_0 is closed iff there exists a subset F of Y such that F is closed and $F_0 = F \cap$ (the carrier of it).

We now state two propositions:

- (3) Let Y be a non empty topological structure, and let Y_0 be a subspace of Y , and let F_0 be a subset of Y_0 . Then F_0 is closed if and only if there exists a subset F of Y such that F is closed and $F_0 = F \cap$ (the carrier of Y_0).
- (4) Let Y be a non empty topological structure, and let Y_0 be a subspace of Y , and let F be a subset of Y . Suppose F is closed. Then there exists a subset F_0 of Y_0 such that F_0 is closed and $F_0 = F \cap$ (the carrier of Y_0).

2. KOLMOGOROV SPACES

A topological structure is T_0 if it satisfies the condition (Def.3).

- (Def.3) Let x, y be points of it. Suppose $x \neq y$. Then
(i) there exists a subset V of it such that V is open and $x \in V$ and $y \notin V$,
or
(ii) there exists a subset W of it such that W is open and $x \notin W$ and $y \in W$.

Let us observe that a non empty topological structure is T_0 if it satisfies the condition (Def.4).

- (Def.4) Let x, y be points of it. Suppose $x \neq y$. Then
(i) there exists a subset E of it such that E is closed and $x \in E$ and $y \notin E$,
or
(ii) there exists a subset F of it such that F is closed and $x \notin F$ and $y \in F$.

Let us mention that every non empty topological structure which is trivial is also T_0 and every non empty topological structure which is non T_0 is also non trivial.

One can verify that there exists a topological space which is strict T_0 and non empty and there exists a topological space which is strict non T_0 and non empty.

One can check the following observations:

- * every topological space which is discrete is also T_0 ,
- * every topological space which is non T_0 is also non discrete,
- * every topological space which is anti-discrete and non trivial is also non T_0 ,
- * every topological space which is anti-discrete and T_0 is also trivial, and
- * every topological space which is T_0 and non trivial is also non anti-discrete.

Let us observe that a topological space is T_0 if:

(Def.5) For all points x, y of it such that $x \neq y$ holds $\overline{\{x\}} \neq \overline{\{y\}}$.

Let us observe that a topological space is T_0 if:

(Def.6) For all points x, y of it such that $x \neq y$ holds $x \notin \overline{\{y\}}$ or $y \notin \overline{\{x\}}$.

Let us observe that a topological space is T_0 if:

(Def.7) For all points x, y of it such that $x \neq y$ and $x \in \overline{\{y\}}$ holds $\overline{\{y\}} \not\subseteq \overline{\{x\}}$.

One can verify the following observations:

- * every topological space which is almost discrete and T_0 is also discrete,
- * every topological space which is almost discrete and non discrete is also non T_0 , and
- * every topological space which is non discrete and T_0 is also non almost discrete.

A Kolmogorov space is a T_0 topological space. A non-Kolmogorov space is a non T_0 topological space.

Let us observe that there exists a Kolmogorov space which is non trivial and strict and there exists a non-Kolmogorov space which is non trivial and strict.

3. T_0 -SUBSETS

Let Y be a topological structure. A subset of Y is T_0 if it satisfies the condition (Def.8).

(Def.8) Let x, y be points of Y . Suppose $x \in it$ and $y \in it$ and $x \neq y$. Then there exists a subset V of Y such that V is open and $x \in V$ and $y \notin V$ or there exists a subset W of Y such that W is open and $x \notin W$ and $y \in W$.

Let Y be a non empty topological structure. Let us observe that a subset of Y is T_0 if it satisfies the condition (Def.9).

(Def.9) Let x, y be points of Y . Suppose $x \in it$ and $y \in it$ and $x \neq y$. Then

- (i) there exists a subset E of Y such that E is closed and $x \in E$ and $y \notin E$, or
- (ii) there exists a subset F of Y such that F is closed and $x \notin F$ and $y \in F$.

Next we state two propositions:

- (5) Let Y_0, Y_1 be topological structures, and let D_0 be a subset of Y_0 , and let D_1 be a subset of Y_1 . Suppose the topological structure of $Y_0 =$ the topological structure of Y_1 and $D_0 = D_1$. If D_0 is T_0 , then D_1 is T_0 .
- (6) Let Y be a non empty topological structure and let A be a subset of Y . Suppose $A =$ the carrier of Y . Then A is T_0 if and only if Y is T_0 .

In the sequel Y will denote a non empty topological structure.

The following propositions are true:

- (7) For all subsets A, B of Y such that $B \subseteq A$ holds if A is T_0 , then B is T_0 .
- (8) For all subsets A, B of Y such that A is T_0 or B is T_0 holds $A \cap B$ is T_0 .
- (9) Let A, B be subsets of Y . Suppose A is open or B is open. If A is T_0 and B is T_0 , then $A \cup B$ is T_0 .
- (10) Let A, B be subsets of Y . Suppose A is closed or B is closed. If A is T_0 and B is T_0 , then $A \cup B$ is T_0 .
- (11) For every subset A of Y such that A is discrete holds A is T_0 .
- (12) For every non empty subset A of Y such that A is anti-discrete and A is not trivial holds A is not T_0 .

Let X be a topological space. Let us observe that a subset of X is T_0 if:

- (Def.10) For all points x, y of X such that $x \in$ it and $y \in$ it and $x \neq y$ holds $\overline{\{x\}} \neq \overline{\{y\}}$.

Let X be a topological space. Let us observe that a subset of X is T_0 if:

- (Def.11) For all points x, y of X such that $x \in$ it and $y \in$ it and $x \neq y$ holds $x \notin \overline{\{y\}}$ or $y \notin \overline{\{x\}}$.

Let X be a topological space. Let us observe that a subset of X is T_0 if:

- (Def.12) For all points x, y of X such that $x \in$ it and $y \in$ it and $x \neq y$ holds if $x \in \overline{\{y\}}$, then $\overline{\{y\}} \not\subseteq \overline{\{x\}}$.

In the sequel X will denote a topological space.

The following two propositions are true:

- (13) Every empty subset of X is T_0 .
- (14) For every point x of X holds $\{x\}$ is T_0 .

4. KOLMOGOROV SUBSPACES

Let Y be a non empty topological structure. Observe that there exists a subspace of Y which is strict and T_0 .

Let Y be a non empty topological structure. Let us observe that a subspace of Y is T_0 if it satisfies the condition (Def.13).

- (Def.13) Let x, y be points of Y . Suppose x is a point of it and y is a point of it and $x \neq y$. Then there exists a subset V of Y such that V is open and

$x \in V$ and $y \notin V$ or there exists a subset W of Y such that W is open and $x \notin W$ and $y \in W$.

Let Y be a non empty topological structure. Let us observe that a subspace of Y is T_0 if it satisfies the condition (Def.14).

(Def.14) Let x, y be points of Y . Suppose x is a point of it and y is a point of it and $x \neq y$. Then

- (i) there exists a subset E of Y such that E is closed and $x \in E$ and $y \notin E$, or
- (ii) there exists a subset F of Y such that F is closed and $x \notin F$ and $y \in F$.

In the sequel Y denotes a non empty topological structure.

The following propositions are true:

- (15) Let Y_0 be a subspace of Y and let A be a subset of Y . Suppose $A =$ the carrier of Y_0 . Then A is T_0 if and only if Y_0 is T_0 .
- (16) Let Y_0 be a subspace of Y and let Y_1 be a T_0 subspace of Y . If Y_0 is a subspace of Y_1 , then Y_0 is T_0 .

Let X be a topological space. One can check that there exists a subspace of X which is strict and T_0 .

In the sequel X is a topological space.

The following propositions are true:

- (17) For every T_0 subspace X_1 of X and for every subspace X_2 of X such that X_1 meets X_2 holds $X_1 \cap X_2$ is T_0 .
- (18) For all T_0 subspaces X_1, X_2 of X such that X_1 is open or X_2 is open holds $X_1 \cup X_2$ is T_0 .
- (19) For all T_0 subspaces X_1, X_2 of X such that X_1 is closed or X_2 is closed holds $X_1 \cup X_2$ is T_0 .

Let X be a topological space. A Kolmogorov subspace of X is a T_0 subspace of X .

Next we state the proposition

- (20) Let X be a topological space and let A_0 be a non empty subset of X . Suppose A_0 is T_0 . Then there exists a strict Kolmogorov subspace X_0 of X such that $A_0 =$ the carrier of X_0 .

Let X be a non trivial topological space. One can verify that there exists a Kolmogorov subspace of X which is proper and strict.

Let X be a Kolmogorov space. Observe that every subspace of X is T_0 .

Let X be a non-Kolmogorov space. One can check that every subspace of X which is non proper is also non T_0 and every subspace of X which is T_0 is also proper.

Let X be a non-Kolmogorov space. Note that there exists a subspace of X which is strict and non T_0 .

Let X be a non-Kolmogorov space. A non-Kolmogorov subspace of X is a non T_0 subspace of X .

We now state the proposition

- (21) Let X be a non-Kolmogorov space and let A_0 be a subset of X . Suppose A_0 is not T_0 . Then there exists a strict non-Kolmogorov subspace X_0 of X such that $A_0 =$ the carrier of X_0 .

REFERENCES

- [1] P. Alexandroff and H. H. Hopf. *Topologie I*. Springer-Verlag, Berlin, 1935.
- [2] Ryszard Engelking. *General Topology*. Volume 60 of *Monografie Matematyczne*, PWN - Polish Scientific Publishers, Warsaw, 1977.
- [3] Zbigniew Karno. Maximal anti-discrete subspaces of topological spaces. *Formalized Mathematics*, 5(1):109–118, 1996.
- [4] Zbigniew Karno. Maximal discrete subspaces of almost discrete topological spaces. *Formalized Mathematics*, 4(1):125–135, 1993.
- [5] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(5):665–674, 1991.
- [6] Kazimierz Kuratowski. *Topology*. Volume I, PWN - Polish Scientific Publishers, Academic Press, Warsaw, New York and London, 1966.
- [7] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [8] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Formalized Mathematics*, 2(4):535–545, 1991.
- [9] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(1):25–34, 1990.
- [10] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [11] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received July 26, 1994
