

Inverse Limits of Many Sorted Algebras

Adam Grabowski
Warsaw University
Białystok

Summary. This article introduces the construction of an inverse limit of many sorted algebras. A few preliminary notions such as an ordered family of many sorted algebras and a binding of family are formulated. Definitions of a set of many sorted signatures and a set of signature morphisms are also given.

MML Identifier: MSALIMIT.

The terminology and notation used here are introduced in the following articles: [21], [25], [12], [22], [26], [9], [28], [10], [5], [23], [8], [18], [27], [11], [3], [7], [24], [2], [1], [20], [15], [19], [6], [14], [17], [16], [4], and [13].

1. INVERSE LIMITS OF MANY SORTED ALGEBRAS

We adopt the following rules: P denotes a non empty poset, i, j, k denote elements of P , and S denotes a non void non empty many sorted signature.

Let I be a non empty set, let us consider S , let A_1 be an algebra family of I over S , let i be an element of I , and let o be an operation symbol of S . One can verify that $(\text{OPER}(A_1))(i)(o)$ is function-like and relation-like.

Let I be a non empty set, let us consider S , let A_1 be an algebra family of I over S , and let s be a sort symbol of S . Note that $(\text{SORTS}(A_1))(s)$ is functional.

Let us consider P, S . An algebra family of the carrier of P over S is called a family of algebras over S ordered by P if it satisfies the condition (Def. 1).

(Def. 1) There exists a many sorted function F of the internal relation of P such that for all i, j, k if $i \geq j$ and $j \geq k$, then there exists a many sorted function f_1 from $\text{it}(i)$ into $\text{it}(j)$ and there exists a many sorted function f_2 from $\text{it}(j)$ into $\text{it}(k)$ such that $f_1 = F(j, i)$ and $f_2 = F(k, j)$ and $F(k, i) = f_2 \circ f_1$ and f_1 is a homomorphism of $\text{it}(i)$ into $\text{it}(j)$.

In the sequel O_1 is a family of algebras over S ordered by P .

Let us consider P, S, O_1 . A many sorted function of the internal relation of P is called a binding of O_1 if it satisfies the condition (Def. 2).

(Def. 2) Given i, j, k . Suppose $i \geq j$ and $j \geq k$. Then there exists a many sorted function f_1 from $O_1(i)$ into $O_1(j)$ and there exists a many sorted function f_2 from $O_1(j)$ into $O_1(k)$ such that $f_1 = \text{it}(j, i)$ and $f_2 = \text{it}(k, j)$ and $\text{it}(k, i) = f_2 \circ f_1$ and f_1 is a homomorphism of $O_1(i)$ into $O_1(j)$.

Let us consider P, S, O_1 , let B be a binding of O_1 , and let us consider i, j . Let us assume that $i \geq j$. The functor $\text{bind}(B, i, j)$ yielding a many sorted function from $O_1(i)$ into $O_1(j)$ is defined by:

(Def. 3) $\text{bind}(B, i, j) = B(j, i)$.

In the sequel B will be a binding of O_1 .

Next we state the proposition

(1) If $i \geq j$ and $j \geq k$, then $\text{bind}(B, j, k) \circ \text{bind}(B, i, j) = \text{bind}(B, i, k)$.

Let us consider P, S, O_1 and let I_1 be a binding of O_1 . We say that I_1 is normalized if and only if:

(Def. 4) For every i holds $I_1(i, i) = \text{id}_{(\text{the sorts of } O_1(i))}$.

We now state the proposition

(2) Given P, S, O_1, B, i, j . Suppose $i \geq j$. Let f be a many sorted function from $O_1(i)$ into $O_1(j)$. If $f = \text{bind}(B, i, j)$, then f is a homomorphism of $O_1(i)$ into $O_1(j)$.

Let us consider P, S, O_1, B . The functor $\text{Normalized}(B)$ yields a binding of O_1 and is defined as follows:

(Def. 5) For all i, j such that $i \geq j$ holds $(\text{Normalized}(B))(j, i) = (j = i \rightarrow \text{id}_{(\text{the sorts of } O_1(i))}, \text{bind}(B, i, j) \circ \text{id}_{(\text{the sorts of } O_1(i))})$.

Next we state the proposition

(3) For all i, j such that $i \geq j$ and $i \neq j$ holds $B(j, i) = (\text{Normalized}(B))(j, i)$.

Let us consider P, S, O_1, B . One can verify that $\text{Normalized}(B)$ is normalized.

Let us consider P, S, O_1 . Note that there exists a binding of O_1 which is normalized.

The following proposition is true

(4) For every normalized binding N_1 of O_1 and for all i, j such that $i \geq j$ holds $(\text{Normalized}(N_1))(j, i) = N_1(j, i)$.

Let us consider P, S, O_1 and let B be a binding of O_1 . The functor $\varprojlim B$ yields a strict subalgebra of $\prod O_1$ and is defined by the condition (Def. 6).

(Def. 6) Let s be a sort symbol of S and let f be an element of $(\text{SORTS}(O_1))(s)$. Then $f \in (\text{the sorts of } \varprojlim B)(s)$ if and only if for all i, j such that $i \geq j$ holds $(\text{bind}(B, i, j))(s)(f(i)) = f(j)$.

Next we state the proposition

- (5) Let D_1 be a discrete non empty poset, and given S , and let O_1 be a family of algebras over S ordered by D_1 , and let B be a normalized binding of O_1 . Then $\varprojlim B = \prod O_1$.

2. SETS AND MORPHISMS OF MANY SORTED SIGNATURES

In the sequel x will be a set and A will be a non empty set.

Let X be a set. We say that X is MSS-membered if and only if:

- (Def. 7) If $x \in X$, then x is a strict non empty non void many sorted signature.

One can verify that there exists a set which is non empty and MSS-membered.

The strict many sorted signature TrivialMSSign is defined by:

- (Def. 8) TrivialMSSign is empty and void.

Let us note that TrivialMSSign is empty and void.

One can check that there exists a many sorted signature which is strict, empty, and void.

The following proposition is true

- (6) Let S be a void many sorted signature. Then $\text{id}_{(\text{the carrier of } S)}$ and $\text{id}_{(\text{the operation symbols of } S)}$ form morphism between S and S .

Let us consider A . The functor $\text{MSS-set}(A)$ is defined by the condition

- (Def. 9).

- (Def. 9) $x \in \text{MSS-set}(A)$ if and only if there exists a strict non empty non void many sorted signature S such that $x = S$ and the carrier of $S \subseteq A$ and the operation symbols of $S \subseteq A$.

Let us consider A . One can check that $\text{MSS-set}(A)$ is non empty and MSS-membered.

Let A be a non empty MSS-membered set. We see that the element of A is a strict non empty non void many sorted signature.

The following proposition is true

- (7) Let x be an element of $\text{MSS-set}(A)$. Then $\text{id}_{(\text{the carrier of } x)}$ and $\text{id}_{(\text{the operation symbols of } x)}$ form morphism between x and x .

Let S_1, S_2 be many sorted signatures. The functor $\text{MSS-morph}(S_1, S_2)$ is defined by:

- (Def. 10) $x \in \text{MSS-morph}(S_1, S_2)$ iff there exist functions f, g such that $x = \langle f, g \rangle$ and f and g form morphism between S_1 and S_2 .

REFERENCES

- [1] Grzegorz Bancerek. Cartesian product of functions. *Formalized Mathematics*, 2(4):547–552, 1991.
- [2] Grzegorz Bancerek. Curried and uncurried functions. *Formalized Mathematics*, 1(3):537–541, 1990.
- [3] Grzegorz Bancerek. König’s theorem. *Formalized Mathematics*, 1(3):589–593, 1990.

- [4] Grzegorz Bancerek. Minimal signature for partial algebra. *Formalized Mathematics*, 5(3):405–414, 1996.
- [5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [6] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Formalized Mathematics*, 5(1):47–54, 1996.
- [7] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [8] Czesław Byliński. A classical first order language. *Formalized Mathematics*, 1(4):669–676, 1990.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [11] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [12] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [13] Adam Grabowski. On the category of posets. *Formalized Mathematics*, 5(4):501–505, 1996.
- [14] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Formalized Mathematics*, 5(1):61–65, 1996.
- [15] Beata Madras. Product of family of universal algebras. *Formalized Mathematics*, 4(1):103–108, 1993.
- [16] Beata Madras. Products of many sorted algebras. *Formalized Mathematics*, 5(1):55–60, 1996.
- [17] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Formalized Mathematics*, 5(2):167–172, 1996.
- [18] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(3):495–500, 1990.
- [19] Andrzej Trybulec. Many sorted algebras. *Formalized Mathematics*, 5(1):37–42, 1996.
- [20] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [22] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(1):97–105, 1990.
- [23] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(2):313–319, 1990.
- [24] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [25] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.
- [26] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [27] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [28] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(1):85–89, 1990.

Received June 11, 1996
