

# Binary Arithmetics. Binary Sequences

Robert Milewski  
 University of Białystok

MML Identifier: BINARI\_3.

The notation and terminology used here are introduced in the following papers: [10], [9], [7], [3], [2], [4], [12], [6], [5], [14], [1], [8], [15], [11], and [13].

## 1. BINARY ARITHMETICS

The following propositions are true:

- (1) For every non empty natural number  $n$  and for every tuple  $F$  of  $n$  and *Boolean* holds  $\text{Absval}(F) <$  the  $n$ -th power of 2.
- (2) For every non empty natural number  $n$  and for all tuples  $F_1, F_2$  of  $n$  and *Boolean* such that  $\text{Absval}(F_1) = \text{Absval}(F_2)$  holds  $F_1 = F_2$ .
- (3) For all finite sequences  $t_1, t_2$  such that  $\text{Rev}(t_1) = \text{Rev}(t_2)$  holds  $t_1 = t_2$ .
- (4) For every natural number  $n$  holds  $\underbrace{\langle 0, \dots, 0 \rangle}_{n+1} = \underbrace{\langle 0, \dots, 0 \rangle}_n \hat{\ } \langle 0 \rangle$ .
- (5) For every natural number  $n$  holds  $\underbrace{\langle 0, \dots, 0 \rangle}_n \in \text{Boolean}^*$ .
- (6) For every natural number  $n$  and for every tuple  $y$  of  $n$  and *Boolean* such that  $y = \underbrace{\langle 0, \dots, 0 \rangle}_n$  holds  $\neg y = n \mapsto 1$ .
- (7) For every non empty natural number  $n$  and for every tuple  $F$  of  $n$  and *Boolean* such that  $F = \underbrace{\langle 0, \dots, 0 \rangle}_n$  holds  $\text{Absval}(F) = 0$ .
- (8) Let  $n$  be a non empty natural number and  $F$  be a tuple of  $n$  and *Boolean*. If  $F = \underbrace{\langle 0, \dots, 0 \rangle}_n$ , then  $\text{Absval}(\neg F) = (\text{the } n\text{-th power of } 2) - 1$ .

- (9) For every natural number  $n$  holds  $\text{Rev}(\underbrace{\langle 0, \dots, 0 \rangle}_n) = \underbrace{\langle 0, \dots, 0 \rangle}_n$ .
- (10) For every natural number  $n$  and for every tuple  $y$  of  $n$  and *Boolean* such that  $y = \underbrace{\langle 0, \dots, 0 \rangle}_n$  holds  $\text{Rev}(\neg y) = \neg y$ .
- (11)  $\text{Bin1}(1) = \langle \text{true} \rangle$ .
- (12) For every non empty natural number  $n$  holds  $\text{Absval}(\text{Bin1}(n)) = 1$ .
- (13) For all elements  $x, y$  of *Boolean* holds  $x \vee y = \text{true}$  iff  $x = \text{true}$  or  $y = \text{true}$  and  $x \vee y = \text{false}$  iff  $x = \text{false}$  and  $y = \text{false}$ .
- (14) For all elements  $x, y$  of *Boolean* holds  $\text{add\_ovfl}(\langle x \rangle, \langle y \rangle) = \text{true}$  iff  $x = \text{true}$  and  $y = \text{true}$ .
- (15)  $\neg \langle \text{false} \rangle = \langle \text{true} \rangle$ .
- (16)  $\neg \langle \text{true} \rangle = \langle \text{false} \rangle$ .
- (17)  $\langle \text{false} \rangle + \langle \text{false} \rangle = \langle \text{false} \rangle$ .
- (18)  $\langle \text{false} \rangle + \langle \text{true} \rangle = \langle \text{true} \rangle$  and  $\langle \text{true} \rangle + \langle \text{false} \rangle = \langle \text{true} \rangle$ .
- (19)  $\langle \text{true} \rangle + \langle \text{true} \rangle = \langle \text{false} \rangle$ .
- (20) Let  $n$  be a non empty natural number and  $x, y$  be tuples of  $n$  and *Boolean*. Suppose  $\pi_n x = \text{true}$  and  $\pi_n \text{carry}(x, \text{Bin1}(n)) = \text{true}$ . Let  $k$  be a non empty natural number. If  $k \neq 1$  and  $k \leq n$ , then  $\pi_k x = \text{true}$  and  $\pi_k \text{carry}(x, \text{Bin1}(n)) = \text{true}$ .
- (21) For every non empty natural number  $n$  and for every tuple  $x$  of  $n$  and *Boolean* such that  $\pi_n x = \text{true}$  and  $\pi_n \text{carry}(x, \text{Bin1}(n)) = \text{true}$  holds  $\text{carry}(x, \text{Bin1}(n)) = \neg \text{Bin1}(n)$ .
- (22) Let  $n$  be a non empty natural number and  $x, y$  be tuples of  $n$  and *Boolean*. If  $y = \underbrace{\langle 0, \dots, 0 \rangle}_n$  and  $\pi_n x = \text{true}$  and  $\pi_n \text{carry}(x, \text{Bin1}(n)) = \text{true}$ , then  $x = \neg y$ .
- (23) For every non empty natural number  $n$  and for every tuple  $y$  of  $n$  and *Boolean* such that  $y = \underbrace{\langle 0, \dots, 0 \rangle}_n$  holds  $\text{carry}(\neg y, \text{Bin1}(n)) = \neg \text{Bin1}(n)$ .
- (24) Let  $n$  be a non empty natural number and  $x, y$  be tuples of  $n$  and *Boolean*. If  $y = \underbrace{\langle 0, \dots, 0 \rangle}_n$ , then  $\text{add\_ovfl}(x, \text{Bin1}(n)) = \text{true}$  iff  $x = \neg y$ .
- (25) For every non empty natural number  $n$  and for every tuple  $z$  of  $n$  and *Boolean* such that  $z = \underbrace{\langle 0, \dots, 0 \rangle}_n$  holds  $\neg z + \text{Bin1}(n) = z$ .

## 2. BINARY SEQUENCES

Let  $n, k$  be natural numbers. The functor  $n$ -BinarySequence( $k$ ) yielding a tuple of  $n$  and *Boolean* is defined by:

- (Def. 1) For every natural number  $i$  such that  $i \in \text{Seg } n$  holds  
 $\pi_i(n\text{-BinarySequence}(k)) = ((k \div (\text{the } (i - 1)\text{-th power of } 2)) \bmod 2 = 0 \rightarrow \text{false}, \text{true}).$

One can prove the following propositions:

- (26) For every natural number  $n$  holds  $n\text{-BinarySequence}(0) = \underbrace{\langle 0, \dots, 0 \rangle}_n$ .
- (27) For all natural numbers  $n, k$  such that  $k < \text{the } n\text{-th power of } 2$  holds  
 $((n + 1)\text{-BinarySequence}(k))(n + 1) = \text{false}.$
- (28) Let  $n$  be a non empty natural number and  $k$  be a natural number. If  $k < \text{the } n\text{-th power of } 2$ , then  $(n + 1)\text{-BinarySequence}(k) = (n\text{-BinarySequence}(k)) \wedge \langle \text{false} \rangle$ .
- (29) For every non empty natural number  $n$  holds  $(n + 1)\text{-BinarySequence}(\text{the } n\text{-th power of } 2) = \underbrace{\langle 0, \dots, 0 \rangle}_n \wedge \langle \text{true} \rangle$ .
- (30) Let  $n$  be a non empty natural number and  $k$  be a natural number. Suppose the  $n$ -th power of  $2 \leq k$  and  $k < \text{the } (n + 1)\text{-th power of } 2$ . Then  $((n + 1)\text{-BinarySequence}(k))(n + 1) = \text{true}.$
- (31) Let  $n$  be a non empty natural number and  $k$  be a natural number. Suppose the  $n$ -th power of  $2 \leq k$  and  $k < \text{the } (n + 1)\text{-th power of } 2$ . Then  $(n + 1)\text{-BinarySequence}(k) = (n\text{-BinarySequence}(k - (\text{the } n\text{-th power of } 2))) \wedge \langle \text{true} \rangle$ .
- (32) Let  $n$  be a non empty natural number and  $k$  be a natural number. Suppose  $k < \text{the } n\text{-th power of } 2$ . Let  $x$  be a tuple of  $n$  and *Boolean*. If  $x = \underbrace{\langle 0, \dots, 0 \rangle}_n$ , then  $n\text{-BinarySequence}(k) = \neg x$  iff  $k = (\text{the } n\text{-th power of } 2) - 1$ .
- (33) Let  $n$  be a non empty natural number and  $k$  be a natural number. If  $k + 1 < \text{the } n\text{-th power of } 2$ , then  $\text{add\_ovfl}(n\text{-BinarySequence}(k), \text{Bin1}(n)) = \text{false}.$
- (34) Let  $n$  be a non empty natural number and  $k$  be a natural number. If  $k + 1 < \text{the } n\text{-th power of } 2$ , then  $n\text{-BinarySequence}(k + 1) = (n\text{-BinarySequence}(k)) + \text{Bin1}(n)$ .
- (35) For all natural numbers  $n, i$  holds  $(n + 1)\text{-BinarySequence}(i) = \langle i \bmod 2 \rangle \wedge (n\text{-BinarySequence}(i \div 2))$ .
- (36) For every non empty natural number  $n$  and for every natural number  $k$

such that  $k < 2^n$  holds  $\text{Absval}(n\text{-BinarySequence}(k)) = k$ .

- (37) For every non empty natural number  $n$  and for every tuple  $x$  of  $n$  and *Boolean* holds  $n\text{-BinarySequence}(\text{Absval}(x)) = x$ .

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [4] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [6] Czesław Byliński. Some properties of restrictions of finite sequences. *Formalized Mathematics*, 5(2):241–245, 1996.
- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [9] Yasuho Mizuhara and Takaya Nishiyama. Binary arithmetics, addition and subtraction of integers. *Formalized Mathematics*, 5(1):27–29, 1996.
- [10] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [11] Konrad Raczkowski and Andrzej Nędzusiak. Serieses. *Formalized Mathematics*, 2(4):449–452, 1991.
- [12] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [13] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [14] Edmund Woronowicz. Many–argument relations. *Formalized Mathematics*, 1(4):733–737, 1990.
- [15] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

Received February 24, 1998

---