

# First-countable, Sequential, and Frechet Spaces

Bartłomiej Skorulski  
University of Białystok

**Summary.** This article contains a definition of three classes of topological spaces: first-countable, Frechet, and sequential. Next there are some facts about them, that every first-countable space is Frechet and every Frechet space is sequential. Next section contains a formalized construction of topological space which is Frechet but not first-countable. This article is based on [9, pp. 73–81].

MML Identifier: FRECHET.

The notation and terminology used here are introduced in the following papers: [19], [2], [15], [4], [5], [6], [11], [1], [13], [3], [12], [14], [10], [20], [21], [18], [16], [8], [7], and [17].

## 1. PRELIMINARIES

One can prove the following proposition

- (1) For every non empty 1-sorted structure  $T$  and for every sequence  $S$  of  $T$  holds  $\text{rng } S$  is a subset of  $T$ .

Let  $T$  be a non empty 1-sorted structure and let  $S$  be a sequence of  $T$ . Then  $\text{rng } S$  is a subset of  $T$ .

The following propositions are true:

- (2) Let  $T_1$  be a non empty 1-sorted structure,  $T_2$  be a 1-sorted structure, and  $S$  be a sequence of  $T_1$ . If  $\text{rng } S \subseteq \text{carrier of } T_2$ , then  $S$  is a sequence of  $T_2$ .
- (3) For every non empty topological space  $T$  and for every point  $x$  of  $T$  and for every basis  $B$  of  $x$  holds  $B \neq \emptyset$ .

Let  $T$  be a non empty topological space and let  $x$  be a point of  $T$ . Note that every basis of  $x$  is non empty.

We now state a number of propositions:

- (4) For every topological space  $T$  and for all subsets  $A, B$  of  $T$  such that  $A$  is open and  $B$  is closed holds  $A \setminus B$  is open.
- (5) Let  $T$  be a topological structure. Suppose that
  - (i)  $\emptyset_T$  is closed,
  - (ii)  $\Omega_T$  is closed,
  - (iii) for all subsets  $A, B$  of  $T$  such that  $A$  is closed and  $B$  is closed holds  $A \cup B$  is closed, and
  - (iv) for every family  $F$  of subsets of  $T$  such that  $F$  is closed holds  $\bigcap F$  is closed.

Then  $T$  is a topological space.

- (6) Let  $T$  be a topological space,  $S$  be a non empty topological structure, and  $f$  be a map from  $T$  into  $S$ . Suppose that for every subset  $A$  of  $S$  holds  $A$  is closed iff  $f^{-1}(A)$  is closed. Then  $S$  is a topological space.
- (7) Let  $x$  be a point of the metric space of real numbers and  $x', r$  be real numbers. If  $x' = x$  and  $r > 0$ , then  $\text{Ball}(x, r) = ]x' - r, x' + r[$ .
- (8) Let  $A$  be a subset of  $\mathbb{R}^1$ . Then  $A$  is open if and only if for every real number  $x$  such that  $x \in A$  there exists a real number  $r$  such that  $r > 0$  and  $]x - r, x + r[ \subseteq A$ .
- (9) For every sequence  $S$  of  $\mathbb{R}^1$  such that for every natural number  $n$  holds  $S(n) \in ]n - \frac{1}{4}, n + \frac{1}{4}[$  holds  $\text{rng } S$  is closed.
- (10) For every subset  $B$  of  $\mathbb{R}^1$  such that  $B = \mathbb{N}$  holds  $B$  is closed.
- (11) Let  $M$  be a metric space,  $x$  be a point of  $M_{\text{top}}$ , and  $x'$  be a point of  $M$ . Suppose  $x = x'$ . Then there exists a basis  $B$  of  $x$  such that
  - (i)  $B = \{\text{Ball}(x', \frac{1}{n}); n \text{ ranges over natural numbers: } n \neq 0\}$ ,
  - (ii)  $B$  is countable, and
  - (iii) there exists a function  $f$  from  $\mathbb{N}$  into  $B$  such that for every set  $n$  such that  $n \in \mathbb{N}$  there exists a natural number  $n'$  such that  $n = n'$  and  $f(n) = \text{Ball}(x', \frac{1}{n'+1})$ .
- (12) For all functions  $f, g$  holds  $\text{rng}(f+g) = f^\circ(\text{dom } f \setminus \text{dom } g) \cup \text{rng } g$ .
- (13) For all sets  $A, B$  such that  $B \subseteq A$  holds  $(\text{id}_A)^\circ B = B$ .
- (14) For all sets  $B, x$  holds  $\text{dom}(B \mapsto x) = B$ .
- (15) For all sets  $A, B, x$  holds  $\text{dom}(\text{id}_A + \cdot (B \mapsto x)) = A \cup B$ .
- (16) For all sets  $A, B, x$  such that  $B \neq \emptyset$  holds  $\text{rng}(\text{id}_A + \cdot (B \mapsto x)) = (A \setminus B) \cup \{x\}$ .
- (17) For all sets  $A, B, C, x$  such that  $C \subseteq A$  holds  $(\text{id}_A + \cdot (B \mapsto x))^{-1}(C \setminus \{x\}) = C \setminus B \setminus \{x\}$ .
- (18) For all sets  $A, B, x$  such that  $x \notin A$  holds  $(\text{id}_A + \cdot (B \mapsto x))^{-1}(\{x\}) = B$ .

- (19) For all sets  $A, B, C, x$  such that  $C \subseteq A$  and  $x \notin A$  holds  $(\text{id}_A + \cdot (B \mapsto x))^{-1}(C \cup \{x\}) = C \cup B$ .
- (20) For all sets  $A, B, C, x$  such that  $C \subseteq A$  and  $x \notin A$  holds  $(\text{id}_A + \cdot (B \mapsto x))^{-1}(C \setminus \{x\}) = C \setminus B$ .

## 2. FIRST-COUNTABLE, SEQUENTIAL, AND FRECHET SPACES

Let  $T$  be a non empty topological structure. We say that  $T$  is first-countable if and only if:

- (Def. 1) For every point  $x$  of  $T$  holds there exists a basis of  $x$  which is countable.

The following two propositions are true:

- (21) For every metric space  $M$  holds  $M_{\text{top}}$  is first-countable.
- (22)  $\mathbb{R}^1$  is first-countable.

Let us note that  $\mathbb{R}^1$  is first-countable.

Let  $T$  be a topological structure, let  $S$  be a sequence of  $T$ , and let  $x$  be a point of  $T$ . We say that  $S$  is convergent to  $x$  if and only if the condition (Def. 2) is satisfied.

- (Def. 2) Let  $U_1$  be a subset of  $T$ . Suppose  $U_1$  is open and  $x \in U_1$ . Then there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $S(m) \in U_1$ .

The following proposition is true

- (23) Let  $T$  be a non empty topological structure,  $x$  be a point of  $T$ , and  $S$  be a sequence of  $T$ . If  $S = \mathbb{N} \mapsto x$ , then  $S$  is convergent to  $x$ .

Let  $T$  be a topological structure and let  $S$  be a sequence of  $T$ . We say that  $S$  is convergent if and only if:

- (Def. 3) There exists a point  $x$  of  $T$  such that  $S$  is convergent to  $x$ .

Let  $T$  be a non empty topological structure and let  $S$  be a sequence of  $T$ .

The functor  $\text{Lim } S$  yields a subset of  $T$  and is defined as follows:

- (Def. 4) For every point  $x$  of  $T$  holds  $x \in \text{Lim } S$  iff  $S$  is convergent to  $x$ .

Let  $T$  be a non empty topological structure. We say that  $T$  is Frechet if and only if the condition (Def. 5) is satisfied.

- (Def. 5) Let  $A$  be a subset of  $T$  and  $x$  be a point of  $T$ . If  $x \in \overline{A}$ , then there exists a sequence  $S$  of  $T$  such that  $\text{rng } S \subseteq A$  and  $x \in \text{Lim } S$ .

Let  $T$  be a non empty topological structure. We say that  $T$  is sequential if and only if the condition (Def. 6) is satisfied.

- (Def. 6) Let  $A$  be a subset of  $T$ . Then  $A$  is closed if and only if for every sequence  $S$  of  $T$  such that  $S$  is convergent and  $\text{rng } S \subseteq A$  holds  $\text{Lim } S \subseteq A$ .

The following proposition is true

- (24) For every non empty topological space  $T$  such that  $T$  is first-countable holds  $T$  is Frechet.

Let us observe that every non empty topological space which is first-countable is also Frechet.

We now state four propositions:

- (25)  $\mathbb{R}^1$  is Frechet.
- (26) Let  $T$  be a non empty topological space and  $A$  be a subset of  $T$ . Suppose  $A$  is closed. Let  $S$  be a sequence of  $T$ . If  $S$  is convergent and  $\text{rng } S \subseteq A$ , then  $\text{Lim } S \subseteq A$ .
- (27) Let  $T$  be a non empty topological space. Suppose that for every subset  $A$  of  $T$  such that for every sequence  $S$  of  $T$  such that  $S$  is convergent and  $\text{rng } S \subseteq A$  holds  $\text{Lim } S \subseteq A$  holds  $A$  is closed. Then  $T$  is sequential.
- (28) For every non empty topological space  $T$  such that  $T$  is Frechet holds  $T$  is sequential.

Let us mention that every non empty topological space which is Frechet is also sequential.

Next we state the proposition

- (29)  $\mathbb{R}^1$  is sequential.

### 3. COUNTEREXAMPLE OF FRECHET BUT NOT FIRST-COUNTABLE SPACE

The strict non empty topological space  $\mathbb{R}^1_{/\mathbb{N}}$  is defined by the conditions (Def. 7).

- (Def. 7)(i) The carrier of  $\mathbb{R}^1_{/\mathbb{N}} = (\mathbb{R} \setminus \mathbb{N}) \cup \{\mathbb{R}\}$ , and
- (ii) there exists a map  $f$  from  $\mathbb{R}^1$  into  $\mathbb{R}^1_{/\mathbb{N}}$  such that  $f = \text{id}_{\mathbb{R}} + \cdot (\mathbb{N} \mapsto \mathbb{R})$  and for every subset  $A$  of  $\mathbb{R}^1_{/\mathbb{N}}$  holds  $A$  is closed iff  $f^{-1}(A)$  is closed.

We now state several propositions:

- (30)  $\mathbb{R}$  is a point of  $\mathbb{R}^1_{/\mathbb{N}}$ .
- (31) Let  $A$  be a subset of  $\mathbb{R}^1_{/\mathbb{N}}$ . Then  $A$  is open and  $\mathbb{R} \in A$  if and only if there exists a subset  $O$  of  $\mathbb{R}^1$  such that  $O$  is open and  $\mathbb{N} \subseteq O$  and  $A = (O \setminus \mathbb{N}) \cup \{\mathbb{R}\}$ .
- (32) For every set  $A$  holds  $A$  is a subset of  $\mathbb{R}^1_{/\mathbb{N}}$  and  $\mathbb{R} \notin A$  iff  $A$  is a subset of  $\mathbb{R}^1$  and  $\mathbb{N} \cap A = \emptyset$ .
- (33) Let  $A$  be a subset of  $\mathbb{R}^1$  and  $B$  be a subset of  $\mathbb{R}^1_{/\mathbb{N}}$ . If  $A = B$ , then  $\mathbb{N} \cap A = \emptyset$  and  $A$  is open iff  $\mathbb{R} \notin B$  and  $B$  is open.
- (34) For every subset  $A$  of  $\mathbb{R}^1_{/\mathbb{N}}$  such that  $A = \{\mathbb{R}\}$  holds  $A$  is closed.

- (35)  $\mathbb{R}^1/\mathbb{N}$  is not first-countable.
- (36)  $\mathbb{R}^1/\mathbb{N}$  is Frechet.
- (37) It is not true that for every non empty topological space  $T$  such that  $T$  is Frechet holds  $T$  is first-countable.

#### 4. AUXILIARY THEOREMS

Next we state three propositions:

- (38)  $\frac{1}{4} > 0$  and  $\frac{1}{4} < \frac{1}{2}$ .
- (39) For every real number  $r$  there exists a natural number  $n$  such that  $r < n$ .
- (40) For every real number  $r$  such that  $r > 0$  there exists a natural number  $n$  such that  $\frac{1}{n} < r$  and  $n \neq 0$ .

#### REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Countable sets and Hessenberg's theorem. *Formalized Mathematics*, 2(1):65–69, 1991.
- [3] Leszek Borys. Paracompact and metrizable spaces. *Formalized Mathematics*, 2(4):481–485, 1991.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [7] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257–261, 1990.
- [8] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces - fundamental concepts. *Formalized Mathematics*, 2(4):605–608, 1991.
- [9] Ryszard Engelking. *General Topology*, volume 60 of *Monografie Matematyczne*. PWN - Polish Scientific Publishers, Warsaw, 1977.
- [10] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [11] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Formalized Mathematics*, 1(3):607–610, 1990.
- [12] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [13] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [14] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [15] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [17] Andrzej Trybulec. Baire spaces, Sober spaces. *Formalized Mathematics*, 6(2):289–294, 1997.
- [18] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

- [19] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.
- [20] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [21] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

*Received May 13, 1998*

---