

# On the Dividing Function of the Simple Closed Curve into Segments

Yatsuka Nakamura  
Shinshu University  
Nagano

**Summary.** At the beginning, the concept of the segment of the simple closed curve in 2-dimensional Euclidean space is defined. Some properties of segments are shown in the succeeding theorems. At the end, the existence of the function which can divide the simple closed curve into segments is shown. We can make the diameter of segments as small as we want.

MML Identifier: JORDAN7.

The terminology and notation used in this paper are introduced in the following papers: [17], [5], [7], [2], [15], [3], [11], [12], [13], [1], [14], [4], [18], [16], [10], [8], [9], and [6].

## 1. DEFINITION OF THE SEGMENT AND ITS PROPERTY

In this paper  $p, p_1, q$  are points of  $\mathcal{E}_T^2$ .

The following three propositions are true:

- (1) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$ . Suppose  $P$  is a simple closed curve. Then  $W\text{-min } P \in \text{LowerArc } P$  and  $E\text{-max } P \in \text{LowerArc } P$  and  $W\text{-min } P \in \text{UpperArc } P$  and  $E\text{-max } P \in \text{UpperArc } P$ .
- (2) For every compact non empty subset  $P$  of  $\mathcal{E}_T^2$  and for every  $q$  such that  $P$  is a simple closed curve and  $LE(q, W\text{-min } P, P)$  holds  $q = W\text{-min } P$ .
- (3) For every compact non empty subset  $P$  of  $\mathcal{E}_T^2$  and for every  $q$  such that  $P$  is a simple closed curve and  $q \in P$  holds  $LE(W\text{-min } P, q, P)$ .

Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and let  $q_1, q_2$  be points of  $\mathcal{E}_T^2$ . The functor  $\text{Segment}(q_1, q_2, P)$  yields a subset of  $\mathcal{E}_T^2$  and is defined by:

$$(\text{Def. 1}) \quad \text{Segment}(q_1, q_2, P) = \begin{cases} \{p : \text{LE}(q_1, p, P) \wedge \text{LE}(p, q_2, P)\}, \\ \text{if } q_2 \neq \text{W-min } P, \\ \{p_1 : \text{LE}(q_1, p_1, P) \vee q_1 \in P \wedge p_1 = \text{W-min } P\}, \\ \text{otherwise.} \end{cases}$$

One can prove the following propositions:

- (4) For every compact non empty subset  $P$  of  $\mathcal{E}_T^2$  such that  $P$  is a simple closed curve holds  $\text{Segment}(\text{W-min } P, \text{E-max } P, P) = \text{UpperArc } P$  and  $\text{Segment}(\text{E-max } P, \text{W-min } P, P) = \text{LowerArc } P$ .
- (5) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2$  be points of  $\mathcal{E}_T^2$ . If  $P$  is a simple closed curve and  $\text{LE}(q_1, q_2, P)$ , then  $q_1 \in P$  and  $q_2 \in P$ .
- (6) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2$  be points of  $\mathcal{E}_T^2$ . If  $P$  is a simple closed curve and  $\text{LE}(q_1, q_2, P)$ , then  $q_1 \in \text{Segment}(q_1, q_2, P)$  and  $q_2 \in \text{Segment}(q_1, q_2, P)$ .
- (7) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q$  be a point of  $\mathcal{E}_T^2$ . If  $P$  is a simple closed curve and  $q \in P$  and  $q \neq \text{W-min } P$ , then  $\text{Segment}(q, q, P) = \{q\}$ .
- (8) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2$  be points of  $\mathcal{E}_T^2$ . If  $P$  is a simple closed curve and  $q_1 \neq \text{W-min } P$  and  $q_2 \neq \text{W-min } P$ , then  $\text{W-min } P \notin \text{Segment}(q_1, q_2, P)$ .
- (9) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2, q_3$  be points of  $\mathcal{E}_T^2$ . Suppose  $P$  is a simple closed curve and  $\text{LE}(q_1, q_2, P)$  and  $\text{LE}(q_2, q_3, P)$  and  $q_1 = q_2$  and  $q_1 = \text{W-min } P$  and  $q_1 \neq q_3$  and  $q_2 = q_3$  and  $q_2 = \text{W-min } P$ . Then  $\text{Segment}(q_1, q_2, P) \cap \text{Segment}(q_2, q_3, P) = \{q_2\}$ .
- (10) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2$  be points of  $\mathcal{E}_T^2$ . Suppose  $P$  is a simple closed curve and  $\text{LE}(q_1, q_2, P)$  and  $q_1 \neq q_2$  and  $q_1 \neq \text{W-min } P$ . Then  $\text{Segment}(q_2, \text{W-min } P, P) \cap \text{Segment}(\text{W-min } P, q_1, P) = \{\text{W-min } P\}$ .
- (11) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $q_1, q_2, q_3, q_4$  be points of  $\mathcal{E}_T^2$ . Suppose  $P$  is a simple closed curve and  $\text{LE}(q_1, q_2, P)$  and  $\text{LE}(q_2, q_3, P)$  and  $\text{LE}(q_3, q_4, P)$  and  $q_1 \neq q_2$  and  $q_2 \neq q_3$ . Then  $\text{Segment}(q_1, q_2, P) \cap \text{Segment}(q_3, q_4, P) = \emptyset$ .

## 2. A FUNCTION TO DIVIDE THE SIMPLE CLOSED CURVE

In the sequel  $n$  is a natural number.

We now state three propositions:

- (12) Let  $P$  be a non empty subset of the carrier of  $\mathcal{E}_T^n$  and  $f$  be a map from  $\mathbb{I}$  into  $(\mathcal{E}_T^n) \upharpoonright P$ . Suppose  $P \neq \emptyset$  and  $f$  is a homeomorphism. Then there exists a map  $g$  from  $\mathbb{I}$  into  $\mathcal{E}_T^n$  such that  $f = g$  and  $g$  is continuous and one-to-one.
- (13) For every finite sequence  $f$  of elements of  $\mathbb{R}$  such that  $f$  is increasing holds  $f$  is one-to-one.
- (14) Let  $P$  be a compact non empty subset of  $\mathcal{E}_T^2$  and  $e$  be a real number. Suppose  $P$  is a simple closed curve and  $e > 0$ . Then there exists a finite sequence  $h$  of elements of the carrier of  $\mathcal{E}_T^2$  such that
- (i)  $h(1) = \text{W-min } P$ ,
  - (ii)  $h$  is one-to-one,
  - (iii)  $8 \leq \text{len } h$ ,
  - (iv)  $\text{rng } h \subseteq P$ ,
  - (v) for every natural number  $i$  such that  $1 \leq i$  and  $i < \text{len } h$  holds  $\text{LE}(\pi_i h, \pi_{i+1} h, P)$ ,
  - (vi) for every natural number  $i$  and for every subset  $W$  of the carrier of  $\mathcal{E}^2$  such that  $1 \leq i$  and  $i < \text{len } h$  and  $W = \text{Segment}(\pi_i h, \pi_{i+1} h, P)$  holds  $\emptyset W < e$ ,
  - (vii) for every subset  $W$  of the carrier of  $\mathcal{E}^2$  such that  $W = \text{Segment}(\pi_{\text{len } h} h, \pi_1 h, P)$  holds  $\emptyset W < e$ ,
  - (viii) for every natural number  $i$  such that  $1 \leq i$  and  $i + 1 < \text{len } h$  holds  $\text{Segment}(\pi_i h, \pi_{i+1} h, P) \cap \text{Segment}(\pi_{i+1} h, \pi_{i+2} h, P) = \{\pi_{i+1} h\}$ ,
  - (ix)  $\text{Segment}(\pi_{\text{len } h} h, \pi_1 h, P) \cap \text{Segment}(\pi_1 h, \pi_2 h, P) = \{\pi_1 h\}$ , and
  - (x) for all natural numbers  $i, j$  such that  $1 \leq i$  and  $i < \text{len } h$  and  $1 \leq j$  and  $j < \text{len } h$  and  $i \neq j$  and  $i$  and  $j$  are not adjacent holds  $\text{Segment}(\pi_i h, \pi_{i+1} h, P) \cap \text{Segment}(\pi_j h, \pi_{j+1} h, P) = \emptyset$ .

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in  $\mathcal{E}^2$ . *Formalized Mathematics*, 6(3):427–440, 1997.
- [5] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(2):383–386, 1990.
- [6] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257–261, 1990.

- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces - fundamental concepts. *Formalized Mathematics*, 2(4):605–608, 1991.
- [9] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Simple closed curves. *Formalized Mathematics*, 2(5):663–664, 1991.
- [10] Alicia de la Cruz. Totally bounded metric spaces. *Formalized Mathematics*, 2(4):559–562, 1991.
- [11] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-board - part I. *Formalized Mathematics*, 3(1):107–115, 1992.
- [12] Yatsuka Nakamura and Andrzej Trybulec. Adjacency concept for pairs of natural numbers. *Formalized Mathematics*, 6(1):1–3, 1997.
- [13] Yatsuka Nakamura and Andrzej Trybulec. A decomposition of a simple closed curves and the order of their points. *Formalized Mathematics*, 6(4):563–572, 1997.
- [14] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [15] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [16] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [17] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.
- [18] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

*Received June 16, 1998*

---