

Natural Numbers

Robert Milewski
University of Białystok

MML Identifier: NAT_2.

The papers [6], [4], [2], [7], [1], [3], [5], and [8] provide the terminology and notation for this paper.

1. PRELIMINARIES

In this article we present several logical schemes. The scheme *NonUniqRecExD* deals with a non empty set \mathcal{A} , an element \mathcal{B} of \mathcal{A} , and a ternary predicate \mathcal{P} , and states that:

There exists a function f from \mathbb{N} into \mathcal{A} such that $f(0) = \mathcal{B}$ and for every element n of \mathbb{N} holds $\mathcal{P}[n, f(n), f(n + 1)]$

provided the following condition is satisfied:

- For every natural number n and for every element x of \mathcal{A} there exists an element y of \mathcal{A} such that $\mathcal{P}[n, x, y]$.

The scheme *NonUniqFinRecExD* deals with a non empty set \mathcal{A} , an element \mathcal{B} of \mathcal{A} , a natural number \mathcal{C} , and a ternary predicate \mathcal{P} , and states that:

There exists a finite sequence p of elements of \mathcal{A} such that $\text{len } p = \mathcal{C}$ but $p(1) = \mathcal{B}$ or $\mathcal{C} = 0$ but for every natural number n such that $1 \leq n$ and $n \leq \mathcal{C} - 1$ holds $\mathcal{P}[n, p(n), p(n + 1)]$

provided the parameters meet the following requirement:

- Let n be a natural number. Suppose $1 \leq n$ and $n \leq \mathcal{C} - 1$. Let x be an element of \mathcal{A} . Then there exists an element y of \mathcal{A} such that $\mathcal{P}[n, x, y]$.

The scheme *NonUniqPiFinRecExD* deals with a non empty set \mathcal{A} , an element \mathcal{B} of \mathcal{A} , a natural number \mathcal{C} , and a ternary predicate \mathcal{P} , and states that:

There exists a finite sequence p of elements of \mathcal{A} such that $\text{len } p = \mathcal{C}$ but $\pi_1 p = \mathcal{B}$ or $\mathcal{C} = 0$ but for every natural number n such that $1 \leq n$ and $n \leq \mathcal{C} - 1$ holds $\mathcal{P}[n, \pi_n p, \pi_{n+1} p]$

provided the following condition is met:

- Let n be a natural number. Suppose $1 \leq n$ and $n \leq \mathcal{C} - 1$. Let x be an element of \mathcal{A} . Then there exists an element y of \mathcal{A} such that $\mathcal{P}[n, x, y]$.

The following two propositions are true:

- (1) For every real number x holds $x < \lfloor x \rfloor + 1$.
- (2) For all real numbers x, y such that $x \geq 0$ and $y > 0$ holds $\frac{x}{\lfloor \frac{x}{y} \rfloor + 1} < y$.

2. DIVISION AND REST OF DIVISION

The following propositions are true:

- (3) For every natural number n holds n is empty iff $n = 0$.
- (4) For every natural number n holds $0 \div n = 0$.
- (5) For every non empty natural number n holds $n \div n = 1$.
- (6) For every natural number n holds $n \div 1 = n$.
- (7) For all natural numbers i, j, k, l such that $i \leq j$ and $k \leq j$ holds if $i = (j -' k) + l$, then $k = (j -' i) + l$.
- (8) For all natural numbers i, n such that $i \in \text{Seg } n$ holds $(n -' i) + 1 \in \text{Seg } n$.
- (9) For all natural numbers i, j such that $j < i$ holds $(i -' (j + 1)) + 1 = i -' j$.
- (10) For all natural numbers i, j such that $i \geq j$ holds $j -' i = 0$.
- (11) For all non empty natural numbers i, j holds $i -' j < i$.
- (12) Let n, k be natural numbers. Suppose $k \leq n$. Then the n -th power of 2 = (the k -th power of 2) · (the $(n -' k)$ -th power of 2).
- (13) For all natural numbers n, k such that $k \leq n$ holds the k -th power of 2 | the n -th power of 2.
- (14) For all natural numbers n, k such that $k > 0$ and $n \div k = 0$ holds $n < k$.
- (15) For all natural numbers n, k such that $k > 0$ and $k \leq n$ holds $n \div k \geq 1$.
- (16) For all natural numbers n, k such that $k \neq 0$ holds $(n + k) \div k = (n \div k) + 1$.
- (17) For all natural numbers n, k, i such that $k | n$ and $1 \leq n$ and $1 \leq i$ and $i \leq k$ holds $(n -' i) \div k = (n \div k) - 1$.
- (18) Let n, k be natural numbers. Suppose $k \leq n$. Then (the n -th power of 2) \div (the k -th power of 2) = the $(n -' k)$ -th power of 2.
- (19) For every natural number n such that $n > 0$ holds (the n -th power of 2) mod 2 = 0.

- (20) For every natural number n such that $n > 0$ holds $n \bmod 2 = 0$ iff $(n - ' 1) \bmod 2 = 1$.
- (21) For every non empty natural number n such that $n \neq 1$ holds $n > 1$.
- (22) For all natural numbers n, k such that $n \leq k$ and $k < n + n$ holds $k \div n = 1$.
- (23) For every natural number n holds n is even iff $n \bmod 2 = 0$.
- (24) For every natural number n holds n is odd iff $n \bmod 2 = 1$.
- (25) For all natural numbers n, k, t such that $1 \leq t$ and $k \leq n$ and $2 \cdot t \mid k$ holds $n \div t$ is even iff $(n - ' k) \div t$ is even.
- (26) For all natural numbers n, m, k such that $n \leq m$ holds $n \div k \leq m \div k$.
- (27) For all natural numbers n, k such that $k \leq 2 \cdot n$ holds $(k + 1) \div 2 \leq n$.
- (28) For every even natural number n holds $n \div 2 = (n + 1) \div 2$.
- (29) For all natural numbers n, k, i holds $n \div k \div i = n \div k \cdot i$.

Let n be a natural number. We say that n is non trivial if and only if:

(Def. 1) $n \neq 0$ and $n \neq 1$.

One can verify that every natural number which is non trivial is also non empty.

One can check that there exists a natural number which is non trivial.

The following two propositions are true:

- (30) For every natural number k holds k is non trivial iff k is non empty and $k \neq 1$.
- (31) For every non trivial natural number k holds $k \geq 2$.

The scheme *Ind from 2* concerns a unary predicate \mathcal{P} , and states that:

For every non trivial natural number k holds $\mathcal{P}[k]$

provided the following conditions are met:

- $\mathcal{P}[2]$, and
- For every non trivial natural number k such that $\mathcal{P}[k]$ holds $\mathcal{P}[k + 1]$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [4] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [5] Konrad Raczkowski and Andrzej Nędzusiak. Serieses. *Formalized Mathematics*, 2(4):449–452, 1991.
- [6] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [7] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.

- [8] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received February 23, 1998
