

Completely-Irreducible Elements¹

Robert Milewski
University of Białystok

Summary. The article is a translation of [5, 92–93].

MML Identifier: WAYBEL16.

The terminology and notation used here are introduced in the following articles: [16], [1], [14], [12], [15], [13], [3], [4], [9], [6], [10], [11], [2], [7], and [8].

1. PRELIMINARIES

The following propositions are true:

- (1) For every sup-semilattice L and for all elements x, y of L holds $\bigsqcup_L(\uparrow x \cap \uparrow y) = x \sqcup y$.
- (2) For every semilattice L and for all elements x, y of L holds $\bigsqcup_L(\downarrow x \cap \downarrow y) = x \sqcap y$.
- (3) Let L be a non empty relational structure and x, y be elements of L . If x is maximal in $(\text{the carrier of } L) \setminus \uparrow y$, then $\uparrow x \setminus \{x\} = \uparrow x \cap \uparrow y$.
- (4) Let L be a non empty relational structure and x, y be elements of L . If x is minimal in $(\text{the carrier of } L) \setminus \downarrow y$, then $\downarrow x \setminus \{x\} = \downarrow x \cap \downarrow y$.
- (5) Let L be a poset with l.u.b.'s, X, Y be subsets of L , and X', Y' be subsets of L^{op} . If $X = X'$ and $Y = Y'$, then $X \sqcup Y = X' \sqcap Y'$.
- (6) Let L be a poset with g.l.b.'s, X, Y be subsets of L , and X', Y' be subsets of L^{op} . If $X = X'$ and $Y = Y'$, then $X \sqcap Y = X' \sqcup Y'$.
- (7) For every non empty reflexive transitive relational structure L holds $\text{Filt}(L) = \text{Ids}(L^{\text{op}})$.

¹This work has been supported by KBN Grant 8 T11C 018 12.

- (8) For every non empty reflexive transitive relational structure L holds $\text{Ids}(L) = \text{Filt}(L^{\text{op}})$.

2. FREE GENERATION SET

Let S, T be complete non empty posets. A map from S into T is said to be a CLHomomorphism of S, T if:

- (Def. 1) It is directed-sups-preserving and infs-preserving.

Let S be a continuous complete non empty poset and let A be a subset of S . We say that A is a free generator set if and only if the condition (Def. 2) is satisfied.

- (Def. 2) Let T be a continuous complete non empty poset and f be a function from A into the carrier of T . Then there exists a CLHomomorphism h of S, T such that $h \upharpoonright A = f$ and for every CLHomomorphism h' of S, T such that $h' \upharpoonright A = f$ holds $h' = h$.

Let L be an upper-bounded non empty poset. One can check that $\text{Filt}(L)$ is non empty.

The following propositions are true:

- (9) For every set X and for every non empty subset Y of $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ holds $\bigcap Y$ is a filter of $2_{\underline{C}}^X$.
- (10) For every set X and for every non empty subset Y of $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ holds $\inf Y$ exists in $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ and $\bigcap_{(\text{Filt}(2_{\underline{C}}^X), \subseteq)} Y = \bigcap Y$.
- (11) For every set X holds 2^X is a filter of $2_{\underline{C}}^X$.
- (12) For every set X holds $\{X\}$ is a filter of $2_{\underline{C}}^X$.
- (13) For every set X holds $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ is upper-bounded.
- (14) For every set X holds $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ is lower-bounded.
- (15) For every set X holds $\top_{\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle} = 2^X$.
- (16) For every set X holds $\perp_{\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle} = \{X\}$.
- (17) For every non empty set X and for every non empty subset Y of $\langle X, \subseteq \rangle$ such that $\sup Y$ exists in $\langle X, \subseteq \rangle$ holds $\bigcup Y \subseteq \sup Y$.
- (18) For every upper-bounded semilattice L holds $\langle \text{Filt}(L), \subseteq \rangle$ is complete.

Let L be an upper-bounded semilattice. Note that $\langle \text{Filt}(L), \subseteq \rangle$ is complete.

3. COMPLETELY-IRREDUCIBLE ELEMENTS

Let L be a non empty relational structure and let p be an element of L . We say that p is completely-irreducible if and only if:

(Def. 3) $\text{Min } \uparrow p \setminus \{p\}$ exists in L .

We now state the proposition

- (19) Let L be a non empty relational structure and p be an element of L . If p is completely-irreducible, then $\prod_L(\uparrow p \setminus \{p\}) \neq p$.

Let L be a non empty relational structure. The functor $\text{Irr } L$ yielding a subset of L is defined by:

(Def. 4) For every element x of L holds $x \in \text{Irr } L$ iff x is completely-irreducible.

The following propositions are true:

- (20) Let L be a non empty poset and p be an element of L . Then p is completely-irreducible if and only if there exists an element q of L such that $p < q$ and for every element s of L such that $p < s$ holds $q \leq s$ and $\uparrow p = \{p\} \cup \uparrow q$.
- (21) For every upper-bounded non empty poset L holds $\top_L \notin \text{Irr } L$.
- (22) For every semilattice L holds $\text{Irr } L \subseteq \text{IRR}(L)$.
- (23) For every semilattice L and for every element x of L such that x is completely-irreducible holds x is irreducible.
- (24) Let L be a non empty poset and x be an element of L . Suppose x is completely-irreducible. Let X be a subset of L . If $\inf X$ exists in L and $x = \inf X$, then $x \in X$.
- (25) For every non empty poset L and for every subset X of L such that X is order-generating holds $\text{Irr } L \subseteq X$.
- (26) Let L be a complete lattice and p be an element of L . Given an element k of L such that p is maximal in $(\text{the carrier of } L) \setminus \uparrow k$. Then p is completely-irreducible.
- (27) Let L be a transitive antisymmetric relational structure with l.u.b.'s and p, q, u be elements of L . Suppose $p < q$ and for every element s of L such that $p < s$ holds $q \leq s$ and $u \not\leq p$. Then $p \sqcup u = q \sqcup u$.
- (28) Let L be a distributive lattice and p, q, u be elements of L . Suppose $p < q$ and for every element s of L such that $p < s$ holds $q \leq s$ and $u \not\leq p$. Then $u \sqcap q \not\leq p$.
- (29) Let L be a distributive complete lattice. Suppose L^{op} is meet-continuous. Let p be an element of L . Suppose p is completely-irreducible. Then $(\text{the carrier of } L) \setminus \downarrow p$ is an open filter of L .
- (30) Let L be a distributive complete lattice. Suppose L^{op} is meet-continuous. Let p be an element of L . Suppose p is completely-irreducible. Then there exists an element k of L such that $k \in \text{the carrier of } \text{CompactSublatt}(L)$ and p is maximal in $(\text{the carrier of } L) \setminus \uparrow k$.
- (31) Let L be a lower-bounded algebraic lattice and x, y be elements of L . Suppose $y \not\leq x$. Then there exists an element p of L such that p is

completely-irreducible and $x \leq p$ and $y \not\leq p$.

- (32) Let L be a lower-bounded algebraic lattice. Then $\text{Irr } L$ is order-generating and for every subset X of L such that X is order-generating holds $\text{Irr } L \subseteq X$.
- (33) For every lower-bounded algebraic lattice L and for every element s of L holds $s = \bigcap_L (\uparrow s \cap \text{Irr } L)$.
- (34) Let L be a complete non empty poset, X be a subset of L , and p be an element of L . If p is completely-irreducible and $p = \inf X$, then $p \in X$.
- (35) Let L be a complete algebraic lattice and p be an element of L . Suppose p is completely-irreducible. Then $p = \bigcap_L \{x; x \text{ ranges over elements of } L: x \in \uparrow p \wedge \bigvee_{k: \text{element of } L} (k \in \text{the carrier of } \text{CompactSublatt}(L) \wedge x \text{ is maximal in } (\text{the carrier of } L) \setminus \uparrow k)\}$.
- (36) Let L be a complete algebraic lattice and p be an element of L . Then there exists an element k of L such that $k \in \text{the carrier of } \text{CompactSublatt}(L)$ and p is maximal in $(\text{the carrier of } L) \setminus \uparrow k$ if and only if p is completely-irreducible.

REFERENCES

- [1] Grzegorz Bancerek. Complete lattices. *Formalized Mathematics*, 2(5):719–725, 1991.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. *Formalized Mathematics*, 6(1):81–91, 1997.
- [3] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Formalized Mathematics*, 6(1):93–107, 1997.
- [4] Czesław Byliński. Galois connections. *Formalized Mathematics*, 6(1):131–143, 1997.
- [5] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [6] Adam Grabowski. Auxiliary and approximating relations. *Formalized Mathematics*, 6(2):179–188, 1997.
- [7] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Formalized Mathematics*, 6(1):117–121, 1997.
- [8] Artur Korniłowicz. Definitions and properties of the join and meet of subsets. *Formalized Mathematics*, 6(1):153–158, 1997.
- [9] Artur Korniłowicz. Meet-continuous lattices. *Formalized Mathematics*, 6(1):159–167, 1997.
- [10] Beata Madras. Irreducible and prime elements. *Formalized Mathematics*, 6(2):233–239, 1997.
- [11] Robert Milewski. Algebraic lattices. *Formalized Mathematics*, 6(2):249–254, 1997.
- [12] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [14] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(2):313–319, 1990.
- [15] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [16] Zinaida Trybulec and Halina Święczkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.

Received February 9, 1998
