

Correctness of a Cyclic Redundancy Check Code Generator

Yuguang Yang
Shinshu University
Nagano

Katsumi Wasaki
Shinshu University
Nagano

Yasushi Fuwa
Shinshu University
Nagano

Yatsuka Nakamura
Shinshu University
Nagano

Summary. We prove the correctness of the division circuit and the CRC (cyclic redundancy checks) circuit by verifying the contents of the register after one shift. Circuits with 12-bit register and 16-bit register are taken as examples. All the proofs are done formally.

MML Identifier: GATE_4.

The terminology and notation used here are introduced in the article [1].

1. CORRECTNESS OF DIVISION CIRCUITS WITH 12-BIT REGISTER AND 16-BIT REGISTER

One can prove the following propositions:

- (1) Let $g_0, g_1, g_2, g_3, g_4, g_5, g_6, g_7, g_8, g_9, g_{10}, g_{11}, g_{12}, a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}, b_0, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}, b_{11}, p$ be sets such that NE g_0 and NE g_{12} and NE b_0 iff NE XOR2($p, \text{AND2}(g_0, a_{11})$) and NE b_1 iff NE XOR2($a_0, \text{AND2}(g_1, a_{11})$) and NE b_2 iff NE XOR2($a_1, \text{AND2}(g_2, a_{11})$) and NE b_3 iff NE XOR2($a_2, \text{AND2}(g_3, a_{11})$) and NE b_4 iff NE XOR2($a_3, \text{AND2}(g_4, a_{11})$) and NE b_5 iff NE XOR2($a_4, \text{AND2}(g_5, a_{11})$) and NE b_6 iff NE XOR2($a_5, \text{AND2}(g_6, a_{11})$) and NE b_7 iff NE XOR2($a_6, \text{AND2}(g_7, a_{11})$)

and NE b_8 iff NE XOR2(a_7 , AND2(g_8 , a_{11})) and NE b_9 iff NE XOR2(a_8 , AND2(g_9 , a_{11})) and NE b_{10} iff NE XOR2(a_9 , AND2(g_{10} , a_{11})) and NE b_{11} iff NE XOR2(a_{10} , AND2(g_{11} , a_{11})). Then

- (i) NE a_{11} iff NE AND2(g_{12} , a_{11}),
- (ii) NE a_{10} iff NE XOR2(b_{11} , AND2(g_{11} , a_{11})),
- (iii) NE a_9 iff NE XOR2(b_{10} , AND2(g_{10} , a_{11})),
- (iv) NE a_8 iff NE XOR2(b_9 , AND2(g_9 , a_{11})),
- (v) NE a_7 iff NE XOR2(b_8 , AND2(g_8 , a_{11})),
- (vi) NE a_6 iff NE XOR2(b_7 , AND2(g_7 , a_{11})),
- (vii) NE a_5 iff NE XOR2(b_6 , AND2(g_6 , a_{11})),
- (viii) NE a_4 iff NE XOR2(b_5 , AND2(g_5 , a_{11})),
- (ix) NE a_3 iff NE XOR2(b_4 , AND2(g_4 , a_{11})),
- (x) NE a_2 iff NE XOR2(b_3 , AND2(g_3 , a_{11})),
- (xi) NE a_1 iff NE XOR2(b_2 , AND2(g_2 , a_{11})),
- (xii) NE a_0 iff NE XOR2(b_1 , AND2(g_1 , a_{11})), and
- (xiii) NE p iff NE XOR2(b_0 , AND2(g_0 , a_{11})).

- (2) Let $g_0, g_1, g_2, g_3, g_4, g_5, g_6, g_7, g_8, g_9, g_{10}, g_{11}, g_{12}, g_{13}, g_{14}, g_{15}, g_{16}, a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}, a_{12}, a_{13}, a_{14}, a_{15}, b_0, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}, b_{11}, b_{12}, b_{13}, b_{14}, b_{15}, p$ be sets such that NE g_0 and NE g_{16} and NE b_0 iff NE XOR2(p , AND2(g_0 , a_{15})) and NE b_1 iff NE XOR2(a_0 , AND2(g_1 , a_{15})) and NE b_2 iff NE XOR2(a_1 , AND2(g_2 , a_{15})) and NE b_3 iff NE XOR2(a_2 , AND2(g_3 , a_{15})) and NE b_4 iff NE XOR2(a_3 , AND2(g_4 , a_{15})) and NE b_5 iff NE XOR2(a_4 , AND2(g_5 , a_{15})) and NE b_6 iff NE XOR2(a_5 , AND2(g_6 , a_{15})) and NE b_7 iff NE XOR2(a_6 , AND2(g_7 , a_{15})) and NE b_8 iff NE XOR2(a_7 , AND2(g_8 , a_{15})) and NE b_9 iff NE XOR2(a_8 , AND2(g_9 , a_{15})) and NE b_{10} iff NE XOR2(a_9 , AND2(g_{10} , a_{15})) and NE b_{11} iff NE XOR2(a_{10} , AND2(g_{11} , a_{15})) and NE b_{12} iff NE XOR2(a_{11} , AND2(g_{12} , a_{15})) and NE b_{13} iff NE XOR2(a_{12} , AND2(g_{13} , a_{15})) and NE b_{14} iff NE XOR2(a_{13} , AND2(g_{14} , a_{15})) and NE b_{15} iff NE XOR2(a_{14} , AND2(g_{15} , a_{15})). Then

- (i) NE a_{15} iff NE AND2(g_{16} , a_{15}),
- (ii) NE a_{14} iff NE XOR2(b_{15} , AND2(g_{15} , a_{15})),
- (iii) NE a_{13} iff NE XOR2(b_{14} , AND2(g_{14} , a_{15})),
- (iv) NE a_{12} iff NE XOR2(b_{13} , AND2(g_{13} , a_{15})),
- (v) NE a_{11} iff NE XOR2(b_{12} , AND2(g_{12} , a_{15})),
- (vi) NE a_{10} iff NE XOR2(b_{11} , AND2(g_{11} , a_{15})),
- (vii) NE a_9 iff NE XOR2(b_{10} , AND2(g_{10} , a_{15})),
- (viii) NE a_8 iff NE XOR2(b_9 , AND2(g_9 , a_{15})),
- (ix) NE a_7 iff NE XOR2(b_8 , AND2(g_8 , a_{15})),
- (x) NE a_6 iff NE XOR2(b_7 , AND2(g_7 , a_{15})),
- (xi) NE a_5 iff NE XOR2(b_6 , AND2(g_6 , a_{15})),
- (xii) NE a_4 iff NE XOR2(b_5 , AND2(g_5 , a_{15})),

- (xiii) NE a_3 iff NE XOR2(b_4 , AND2(g_4 , a_{15})),
- (xiv) NE a_2 iff NE XOR2(b_3 , AND2(g_3 , a_{15})),
- (xv) NE a_1 iff NE XOR2(b_2 , AND2(g_2 , a_{15})),
- (xvi) NE a_0 iff NE XOR2(b_1 , AND2(g_1 , a_{15})), and
- (xvii) NE p iff NE XOR2(b_0 , AND2(g_0 , a_{15})).

2. CORRECTNESS OF CRC CIRCUITS WITH GENERATOR POLYNOMIAL OF DEGREE 12 AND 16

Next we state two propositions:

- (3) Let $g_0, g_1, g_2, g_3, g_4, g_5, g_6, g_7, g_8, g_9, g_{10}, g_{11}, g_{12}, a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}, b_0, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}, b_{11}, z, p$ be sets such that NE g_0 and NE g_{12} and not NE z and NE b_0 iff NE XOR2(p, a_{11}) and NE b_1 iff NE XOR2(a_0 , AND2(g_1, b_0)) and NE b_2 iff NE XOR2(a_1 , AND2(g_2, b_0)) and NE b_3 iff NE XOR2(a_2 , AND2(g_3, b_0)) and NE b_4 iff NE XOR2(a_3 , AND2(g_4, b_0)) and NE b_5 iff NE XOR2(a_4 , AND2(g_5, b_0)) and NE b_6 iff NE XOR2(a_5 , AND2(g_6, b_0)) and NE b_7 iff NE XOR2(a_6 , AND2(g_7, b_0)) and NE b_8 iff NE XOR2(a_7 , AND2(g_8, b_0)) and NE b_9 iff NE XOR2(a_8 , AND2(g_9, b_0)) and NE b_{10} iff NE XOR2(a_9 , AND2(g_{10}, b_0)) and NE b_{11} iff NE XOR2(a_{10} , AND2(g_{11}, b_0)). Then
- (i) NE b_{11} iff NE XOR2(XOR2(a_{10} , AND2(g_{11}, a_{11})), XOR2(z , AND2(g_{11}, p))),
 - (ii) NE b_{10} iff NE XOR2(XOR2(a_9 , AND2(g_{10}, a_{11})), XOR2(z , AND2(g_{10}, p))),
 - (iii) NE b_9 iff NE XOR2(XOR2(a_8 , AND2(g_9, a_{11})), XOR2(z , AND2(g_9, p))),
 - (iv) NE b_8 iff NE XOR2(XOR2(a_7 , AND2(g_8, a_{11})), XOR2(z , AND2(g_8, p))),
 - (v) NE b_7 iff NE XOR2(XOR2(a_6 , AND2(g_7, a_{11})), XOR2(z , AND2(g_7, p))),
 - (vi) NE b_6 iff NE XOR2(XOR2(a_5 , AND2(g_6, a_{11})), XOR2(z , AND2(g_6, p))),
 - (vii) NE b_5 iff NE XOR2(XOR2(a_4 , AND2(g_5, a_{11})), XOR2(z , AND2(g_5, p))),
 - (viii) NE b_4 iff NE XOR2(XOR2(a_3 , AND2(g_4, a_{11})), XOR2(z , AND2(g_4, p))),
 - (ix) NE b_3 iff NE XOR2(XOR2(a_2 , AND2(g_3, a_{11})), XOR2(z , AND2(g_3, p))),
 - (x) NE b_2 iff NE XOR2(XOR2(a_1 , AND2(g_2, a_{11})), XOR2(z , AND2(g_2, p))),
 - (xi) NE b_1 iff NE XOR2(XOR2(a_0 , AND2(g_1, a_{11})), XOR2(z , AND2(g_1, p))),
- and
- (xii) NE b_0 iff NE XOR2(XOR2(z , AND2(g_0, a_{11})), XOR2(z , AND2(g_0, p))).
- (4) Let $g_0, g_1, g_2, g_3, g_4, g_5, g_6, g_7, g_8, g_9, g_{10}, g_{11}, g_{12}, g_{13}, g_{14}, g_{15}, g_{16}, a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}, a_{12}, a_{13}, a_{14}, a_{15}, b_0, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}, b_{11}, b_{12}, b_{13}, b_{14}, b_{15}, z, p$ be sets such that NE g_0 and NE g_{16} and not NE z and NE b_0 iff NE XOR2(p, a_{15}) and NE b_1 iff NE XOR2(a_0 , AND2(g_1, b_0)) and NE b_2 iff NE XOR2(a_1 , AND2(g_2, b_0)) and NE b_3 iff NE XOR2(a_2 , AND2(g_3, b_0))

and NE b_4 iff NE XOR2(a_3 , AND2(g_4 , b_0)) and NE b_5 iff NE XOR2(a_4 , AND2(g_5 , b_0)) and NE b_6 iff NE XOR2(a_5 , AND2(g_6 , b_0)) and NE b_7 iff NE XOR2(a_6 , AND2(g_7 , b_0)) and NE b_8 iff NE XOR2(a_7 , AND2(g_8 , b_0)) and NE b_9 iff NE XOR2(a_8 , AND2(g_9 , b_0)) and NE b_{10} iff NE XOR2(a_9 , AND2(g_{10} , b_0)) and NE b_{11} iff NE XOR2(a_{10} , AND2(g_{11} , b_0)) and NE b_{12} iff NE XOR2(a_{11} , AND2(g_{12} , b_0)) and NE b_{13} iff NE XOR2(a_{12} , AND2(g_{13} , b_0)) and NE b_{14} iff NE XOR2(a_{13} , AND2(g_{14} , b_0)) and NE b_{15} iff NE XOR2(a_{14} , AND2(g_{15} , b_0)).

Then

- (i) NE b_{15} iff NE XOR2(XOR2(a_{14} , AND2(g_{15} , a_{15})), XOR2(z , AND2(g_{15} , p))),
 - (ii) NE b_{14} iff NE XOR2(XOR2(a_{13} , AND2(g_{14} , a_{15})), XOR2(z , AND2(g_{14} , p))),
 - (iii) NE b_{13} iff NE XOR2(XOR2(a_{12} , AND2(g_{13} , a_{15})), XOR2(z , AND2(g_{13} , p))),
 - (iv) NE b_{12} iff NE XOR2(XOR2(a_{11} , AND2(g_{12} , a_{15})), XOR2(z , AND2(g_{12} , p))),
 - (v) NE b_{11} iff NE XOR2(XOR2(a_{10} , AND2(g_{11} , a_{15})), XOR2(z , AND2(g_{11} , p))),
 - (vi) NE b_{10} iff NE XOR2(XOR2(a_9 , AND2(g_{10} , a_{15})), XOR2(z , AND2(g_{10} , p))),
 - (vii) NE b_9 iff NE XOR2(XOR2(a_8 , AND2(g_9 , a_{15})), XOR2(z , AND2(g_9 , p))),
 - (viii) NE b_8 iff NE XOR2(XOR2(a_7 , AND2(g_8 , a_{15})), XOR2(z , AND2(g_8 , p))),
 - (ix) NE b_7 iff NE XOR2(XOR2(a_6 , AND2(g_7 , a_{15})), XOR2(z , AND2(g_7 , p))),
 - (x) NE b_6 iff NE XOR2(XOR2(a_5 , AND2(g_6 , a_{15})), XOR2(z , AND2(g_6 , p))),
 - (xi) NE b_5 iff NE XOR2(XOR2(a_4 , AND2(g_5 , a_{15})), XOR2(z , AND2(g_5 , p))),
 - (xii) NE b_4 iff NE XOR2(XOR2(a_3 , AND2(g_4 , a_{15})), XOR2(z , AND2(g_4 , p))),
 - (xiii) NE b_3 iff NE XOR2(XOR2(a_2 , AND2(g_3 , a_{15})), XOR2(z , AND2(g_3 , p))),
 - (xiv) NE b_2 iff NE XOR2(XOR2(a_1 , AND2(g_2 , a_{15})), XOR2(z , AND2(g_2 , p))),
 - (xv) NE b_1 iff NE XOR2(XOR2(a_0 , AND2(g_1 , a_{15})), XOR2(z , AND2(g_1 , p))),
- and
- (xvi) NE b_0 iff NE XOR2(XOR2(z , AND2(g_0 , a_{15})), XOR2(z , AND2(g_0 , p))).

REFERENCES

- [1] Yatsuka Nakamura. Logic gates and logical equivalence of adders. *Formalized Mathematics*, 8(1):35–45, 1999.

Received April 16, 1999
