

Dynkin's Lemma in Measure Theory

Franz Merkl
University of Bielefeld

Summary. This article formalizes the proof of Dynkin's lemma in measure theory. Dynkin's lemma is a useful tool in measure theory and probability theory: it helps frequently to generalize a statement about all elements of a intersection-stable set system to all elements of the sigma-field generated by that system.

MML Identifier: DYNKIN.

The terminology and notation used in this paper have been introduced in the following articles: [5], [11], [1], [4], [2], [3], [7], [6], [12], [13], [8], [10], and [9].

1. PRELIMINARIES

For simplicity, we adopt the following rules: O_1 denotes a non empty set, f denotes a sequence of subsets of O_1 , X , A , B denote subsets of O_1 , D denotes a non empty subset of 2^{O_1} , n , m denote natural numbers, F denotes a non empty set, and x , Y denote sets.

Next we state two propositions:

- (1) For every sequence f of subsets of O_1 and for every x holds $x \in \text{rng } f$ iff there exists n such that $f(n) = x$.
- (2) For every n holds $\text{PSeg } n$ is finite.

Let us consider n . One can verify that $\text{PSeg } n$ is finite.

Next we state the proposition

- (3) For all sets x , y , z such that $x \subseteq y$ holds x misses $z \setminus y$.

Let a , b , c be sets. The functor a, b followed by c is defined as follows:

(Def. 1) a, b followed by $c = (\mathbb{N} \mapsto c) + [0 \mapsto a, 1 \mapsto b]$.

Let a, b, c be sets. Observe that a, b followed by c is function-like and relation-like.

Let X be a non empty set and let a, b, c be elements of X . Then a, b followed by c is a function from \mathbb{N} into X .

Next we state the proposition

- (4) For every non empty set X and for all elements a, b, c of X holds a, b followed by c is a function from \mathbb{N} into X .

Let O_1 be a non empty set and let a, b, c be subsets of O_1 . Then a, b followed by c is a sequence of subsets of O_1 .

One can prove the following propositions:

- (5) For all sets a, b, c holds $(a, b \text{ followed by } c)(0) = a$ and $(a, b \text{ followed by } c)(1) = b$ and for every n such that $n \neq 0$ and $n \neq 1$ holds $(a, b \text{ followed by } c)(n) = c$.
- (6) For all subsets a, b of O_1 holds $\bigcup \text{rng}(a, b \text{ followed by } \emptyset) = a \cup b$.

Let O_1 be a non empty set, let f be a sequence of subsets of O_1 , and let X be a subset of O_1 . The functor $\text{seqIntersection}(X, f)$ yields a sequence of subsets of O_1 and is defined by:

- (Def. 2) For every n holds $(\text{seqIntersection}(X, f))(n) = X \cap f(n)$.

2. DISJOINT-VALUED FUNCTIONS AND INTERSECTION

Let us consider O_1 and let us consider f . Let us observe that f is disjoint valued if and only if:

- (Def. 3) If $n < m$, then $f(n)$ misses $f(m)$.

We now state the proposition

- (7) For every non empty set Y and for every x holds $x \subseteq \bigcap Y$ iff for every element y of Y holds $x \subseteq y$.

Let x be a set. We introduce x is intersection stable as a synonym of x is multiplicative.

Let O_1 be a non empty set, let f be a sequence of subsets of O_1 , and let n be an element of \mathbb{N} . The functor $\text{disjointify}(f, n)$ yielding an element of 2^{O_1} is defined by:

- (Def. 5)¹ $\text{disjointify}(f, n) = f(n) \setminus \bigcup \text{rng}(f \upharpoonright \text{PSeg } n)$.

Let O_1 be a non empty set and let g be a sequence of subsets of O_1 . The functor $\text{disjointify } g$ yielding a sequence of subsets of O_1 is defined as follows:

- (Def. 6) For every n holds $(\text{disjointify } g)(n) = \text{disjointify}(g, n)$.

The following propositions are true:

¹The definition (Def. 4) has been removed.

- (8) For every n holds $(\text{disjointify } f)(n) = f(n) \setminus \bigcup \text{rng}(f \upharpoonright \text{PSeg } n)$.
- (9) For every sequence f of subsets of O_1 holds $\text{disjointify } f$ is disjoint valued.
- (10) For every sequence f of subsets of O_1 holds $\bigcup \text{rng } \text{disjointify } f = \bigcup \text{rng } f$.
- (11) For all subsets x, y of O_1 such that x misses y holds x, y followed by $\emptyset_{(O_1)}$ is disjoint valued.
- (12) Let f be a sequence of subsets of O_1 . Suppose f is disjoint valued. Let X be a subset of O_1 . Then $\text{seqIntersection}(X, f)$ is disjoint valued.
- (13) For every sequence f of subsets of O_1 and for every subset X of O_1 holds $X \cap \text{Union } f = \text{Union } \text{seqIntersection}(X, f)$.

3. DYNKIN SYSTEMS: DEFINITION AND CLOSURE PROPERTIES

Let us consider O_1 . A subset of 2^{O_1} is called a Dynkin system of O_1 if:

- (Def. 7) For every f such that $\text{rng } f \subseteq \text{it}$ and f is disjoint valued holds $\text{Union } f \in \text{it}$ and for every X such that $X \in \text{it}$ holds $X^c \in \text{it}$ and $\emptyset \in \text{it}$.

Let us consider O_1 . One can check that every Dynkin system of O_1 is non empty.

The following propositions are true:

- (14) 2^{O_1} is a Dynkin system of O_1 .
- (15) If for every Y such that $Y \in F$ holds Y is a Dynkin system of O_1 , then $\bigcap F$ is a Dynkin system of O_1 .
- (16) If D is a Dynkin system of O_1 and intersection stable, then if $A \in D$ and $B \in D$, then $A \setminus B \in D$.
- (17) If D is a Dynkin system of O_1 and intersection stable, then if $A \in D$ and $B \in D$, then $A \cup B \in D$.
- (18) Suppose D is a Dynkin system of O_1 and intersection stable. Let x be a finite set. If $x \subseteq D$, then $\bigcup x \in D$.
- (19) Suppose D is a Dynkin system of O_1 and intersection stable. Let f be a sequence of subsets of O_1 . If $\text{rng } f \subseteq D$, then $\text{rng } \text{disjointify } f \subseteq D$.
- (20) Suppose D is a Dynkin system of O_1 and intersection stable. Let f be a sequence of subsets of O_1 . If $\text{rng } f \subseteq D$, then $\bigcup \text{rng } f \in D$.
- (21) For every Dynkin system D of O_1 and for all elements x, y of D such that x misses y holds $x \cup y \in D$.
- (22) For every Dynkin system D of O_1 and for all elements x, y of D such that $x \subseteq y$ holds $y \setminus x \in D$.

4. MAIN STEPS FOR DYNKIN'S LEMMA

One can prove the following proposition

- (23) If D is a Dynkin system of O_1 and intersection stable, then D is a σ -field of subsets of O_1 .

Let O_1 be a non empty set and let E be a subset of 2^{O_1} . The functor $\text{GenDynSys } E$ yielding a Dynkin system of O_1 is defined by:

- (Def. 8) $E \subseteq \text{GenDynSys } E$ and for every Dynkin system D of O_1 such that $E \subseteq D$ holds $\text{GenDynSys } E \subseteq D$.

Let O_1 be a non empty set, let G be a set, and let X be a subset of O_1 . The functor $\text{DynSys}(X, G)$ yields a subset of 2^{O_1} and is defined as follows:

- (Def. 9) For every subset A of O_1 holds $A \in \text{DynSys}(X, G)$ iff $A \cap X \in G$.

Let O_1 be a non empty set, let G be a Dynkin system of O_1 , and let X be an element of G . Then $\text{DynSys}(X, G)$ is a Dynkin system of O_1 .

Next we state four propositions:

- (24) Let E be a subset of 2^{O_1} and X, Y be subsets of O_1 . If $X \in E$ and $Y \in \text{GenDynSys } E$ and E is intersection stable, then $X \cap Y \in \text{GenDynSys } E$.
- (25) Let E be a subset of 2^{O_1} and X, Y be subsets of O_1 . If $X \in \text{GenDynSys } E$ and $Y \in \text{GenDynSys } E$ and E is intersection stable, then $X \cap Y \in \text{GenDynSys } E$.
- (26) For every subset E of 2^{O_1} such that E is intersection stable holds $\text{GenDynSys } E$ is intersection stable.
- (27) Let E be a subset of 2^{O_1} . Suppose E is intersection stable. Let D be a Dynkin system of O_1 . If $E \subseteq D$, then $\sigma(E) \subseteq D$.

REFERENCES

- [1] Grzegorz Bancerek. Continuous, stable, and linear maps of coherence spaces. *Formalized Mathematics*, 5(3):381–393, 1996.
- [2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [4] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [5] Michał Muzalewski and Lesław W. Szczërba. Construction of finite sequences over ring and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(1):97–104, 1991.
- [6] Andrzej Nędzusiak. Probability. *Formalized Mathematics*, 1(4):745–749, 1990.
- [7] Andrzej Nędzusiak. σ -fields and probability. *Formalized Mathematics*, 1(2):401–407, 1990.
- [8] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [9] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [10] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [11] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.
- [12] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

- [13] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received November 27, 2000
