

The Correctness of the High Speed Array Multiplier Circuits

Hiroshi Yamazaki
Shinshu University
Nagano

Katsumi Wasaki
Shinshu University
Nagano

Summary. This article introduces the verification of the correctness for the operations and the specification of the high speed array multiplier. We formalize the concepts of 2-by-2 and 3-by-3 bit Plain array multiplier, 3-by-3 Wallace tree multiplier circuit, and show that outputs of the array multiplier are equivalent to outputs of normal (sequential) multiplier.

MML Identifier: GATE_5.

The articles [3], [1], and [2] provide the terminology and notation for this paper.

1. PRELIMINARIES

Let x_0, x_1, y_0, y_1 be sets. The functor $MULT_{210}(x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

(Def. 1) $MULT_{210}(x_1, y_1, x_0, y_0) = AND2(x_0, y_0)$.

The functor $MULT_{211}(x_1, y_1, x_0, y_0)$ yields a set and is defined by:

(Def. 2) $MULT_{211}(x_1, y_1, x_0, y_0) = ADD1(AND2(x_1, y_0), AND2(x_0, y_1), \emptyset)$.

The functor $MULT_{212}(x_1, y_1, x_0, y_0)$ yielding a set is defined as follows:

(Def. 3) $MULT_{212}(x_1, y_1, x_0, y_0) = ADD2(\emptyset, AND2(x_1, y_1), AND2(x_1, y_0), AND2(x_0, y_1), \emptyset)$.

The functor $MULT_{213}(x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

(Def. 4) $MULT_{213}(x_1, y_1, x_0, y_0) = CARR2(\emptyset, AND2(x_1, y_1), AND2(x_1, y_0), AND2(x_0, y_1), \emptyset)$.

We now state the proposition

- (1) Let $x_0, x_1, y_0, y_1, z_0, z_1, z_2, z_3, q_0, q_1, c_1, q_{11}, c_{11}$ be sets such that NE q_0 iff NE $\text{AND2}(x_0, y_0)$ and NE q_1 iff NE $\text{XOR3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE c_1 iff NE $\text{MAJ3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE q_{11} iff NE $\text{XOR3}(\text{AND2}(x_1, y_1), \emptyset, c_1)$ and NE c_{11} iff NE $\text{MAJ3}(\text{AND2}(x_1, y_1), \emptyset, c_1)$ and NE z_0 iff NE q_0 and NE z_1 iff NE q_1 and NE z_2 iff NE q_{11} and NE z_3 iff NE c_{11} . Then
- (i) NE z_0 iff NE $\text{MULT}_{210}(x_1, y_1, x_0, y_0)$,
 - (ii) NE z_1 iff NE $\text{MULT}_{211}(x_1, y_1, x_0, y_0)$,
 - (iii) NE z_2 iff NE $\text{MULT}_{212}(x_1, y_1, x_0, y_0)$, and
 - (iv) NE z_3 iff NE $\text{MULT}_{213}(x_1, y_1, x_0, y_0)$.

Let x_0, x_1, x_2, y_0, y_1 be sets. The functor $\text{MULT}_{310}(x_2, x_1, y_1, x_0, y_0)$ yields a set and is defined by:

$$\text{(Def. 5)} \quad \text{MULT}_{310}(x_2, x_1, y_1, x_0, y_0) = \text{AND2}(x_0, y_0).$$

The functor $\text{MULT}_{311}(x_2, x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

$$\text{(Def. 6)} \quad \text{MULT}_{311}(x_2, x_1, y_1, x_0, y_0) = \text{ADD1}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset).$$

The functor $\text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

$$\text{(Def. 7)} \quad \text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0) = \text{ADD2}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset).$$

The functor $\text{MULT}_{313}(x_2, x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

$$\text{(Def. 8)} \quad \text{MULT}_{313}(x_2, x_1, y_1, x_0, y_0) = \text{ADD3}(\emptyset, \text{AND2}(x_2, y_1), \text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset).$$

The functor $\text{MULT}_{314}(x_2, x_1, y_1, x_0, y_0)$ yielding a set is defined by:

$$\text{(Def. 9)} \quad \text{MULT}_{314}(x_2, x_1, y_1, x_0, y_0) = \text{CARR3}(\emptyset, \text{AND2}(x_2, y_1), \text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset).$$

Let $x_0, x_1, x_2, y_0, y_1, y_2$ be sets. The functor $\text{MULT}_{321}(x_2, y_2, x_1, y_1, x_0, y_0)$ yields a set and is defined by:

$$\text{(Def. 10)} \quad \text{MULT}_{321}(x_2, y_2, x_1, y_1, x_0, y_0) = \text{ADD1}(\text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_0, y_2), \emptyset).$$

The functor $\text{MULT}_{322}(x_2, y_2, x_1, y_1, x_0, y_0)$ yields a set and is defined by:

$$\text{(Def. 11)} \quad \text{MULT}_{322}(x_2, y_2, x_1, y_1, x_0, y_0) = \text{ADD2}(\text{MULT}_{313}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_1, y_2), \text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_0, y_2), \emptyset).$$

The functor $\text{MULT}_{323}(x_2, y_2, x_1, y_1, x_0, y_0)$ yields a set and is defined as follows:

$$\text{(Def. 12)} \quad \text{MULT}_{323}(x_2, y_2, x_1, y_1, x_0, y_0) = \text{ADD3}(\text{MULT}_{314}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_2, y_2), \text{MULT}_{313}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_1, y_2), \text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_0, y_2), \emptyset).$$

The functor $\text{MULT}_{324}(x_2, y_2, x_1, y_1, x_0, y_0)$ yielding a set is defined as follows:

$$\text{(Def. 13)} \quad \text{MULT}_{324}(x_2, y_2, x_1, y_1, x_0, y_0) = \text{CARR3}(\text{MULT}_{314}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_2, y_2), \text{MULT}_{313}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_1, y_2), \text{MULT}_{312}(x_2, x_1, y_1, x_0, y_0), \text{AND2}(x_0, y_2), \emptyset).$$

$y_1, x_0, y_0), \text{AND2}(x_0, y_2), \emptyset)$.

Next we state the proposition

- (2) Let $x_0, x_1, x_2, y_0, y_1, y_2, z_0, z_1, z_2, z_3, z_4, z_5, q_0, q_1, q_2, c_1, c_2, q_{11}, q_{12}, c_{11}, c_{12}, q_{21}, q_{22}, c_{21}, c_{22}$ be sets such that NE q_0 iff NE $\text{AND2}(x_0, y_0)$ and NE q_1 iff NE $\text{XOR3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE c_1 iff NE $\text{MAJ3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE q_2 iff NE $\text{XOR3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \emptyset)$ and NE c_2 iff NE $\text{MAJ3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \emptyset)$ and NE q_{11} iff NE $\text{XOR3}(q_2, \text{AND2}(x_0, y_2), c_1)$ and NE c_{11} iff NE $\text{MAJ3}(q_2, \text{AND2}(x_0, y_2), c_1)$ and NE q_{12} iff NE $\text{XOR3}(\text{AND2}(x_2, y_1), \text{AND2}(x_1, y_2), c_2)$ and NE c_{12} iff NE $\text{MAJ3}(\text{AND2}(x_2, y_1), \text{AND2}(x_1, y_2), c_2)$ and NE q_{21} iff NE $\text{XOR3}(q_{12}, \emptyset, c_{11})$ and NE c_{21} iff NE $\text{MAJ3}(q_{12}, \emptyset, c_{11})$ and NE q_{22} iff NE $\text{XOR3}(\text{AND2}(x_2, y_2), c_{21}, c_{12})$ and NE c_{22} iff NE $\text{MAJ3}(\text{AND2}(x_2, y_2), c_{21}, c_{12})$ and NE z_0 iff NE q_0 and NE z_1 iff NE q_1 and NE z_2 iff NE q_{11} and NE z_3 iff NE q_{21} and NE z_4 iff NE q_{22} and NE z_5 iff NE c_{22} . Then
- (i) NE z_0 iff NE $\text{MULT}_{310}(x_2, x_1, y_1, x_0, y_0)$,
 - (ii) NE z_1 iff NE $\text{MULT}_{311}(x_2, x_1, y_1, x_0, y_0)$,
 - (iii) NE z_2 iff NE $\text{MULT}_{321}(x_2, y_2, x_1, y_1, x_0, y_0)$,
 - (iv) NE z_3 iff NE $\text{MULT}_{322}(x_2, y_2, x_1, y_1, x_0, y_0)$,
 - (v) NE z_4 iff NE $\text{MULT}_{323}(x_2, y_2, x_1, y_1, x_0, y_0)$, and
 - (vi) NE z_5 iff NE $\text{MULT}_{324}(x_2, y_2, x_1, y_1, x_0, y_0)$.

2. LOGICAL EQUIVALENCE OF WALLACE TREE MULTIPLIER

One can prove the following proposition

- (3) Let $x_0, x_1, x_2, y_0, y_1, y_2, z_0, z_1, z_2, z_3, z_4, z_5, q_0, q_1, q_2, q_3, c_1, c_2, c_3, q_{11}, q_{12}, q_{13}, c_{11}, c_{12}, c_{13}$ be sets such that NE q_0 iff NE $\text{AND2}(x_0, y_0)$ and NE q_1 iff NE $\text{XOR3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE c_1 iff NE $\text{MAJ3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE q_2 iff NE $\text{XOR3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_0, y_2))$ and NE c_2 iff NE $\text{MAJ3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_0, y_2))$ and NE q_3 iff NE $\text{XOR3}(\text{AND2}(x_2, y_1), \text{AND2}(x_1, y_2), \emptyset)$ and NE c_3 iff NE $\text{MAJ3}(\text{AND2}(x_2, y_1), \text{AND2}(x_1, y_2), \emptyset)$ and NE q_{11} iff NE $\text{XOR3}(q_2, c_1, \emptyset)$ and NE c_{11} iff NE $\text{MAJ3}(q_2, c_1, \emptyset)$ and NE q_{12} iff NE $\text{XOR3}(q_3, c_2, c_{11})$ and NE c_{12} iff NE $\text{MAJ3}(q_3, c_2, c_{11})$ and NE q_{13} iff NE $\text{XOR3}(\text{AND2}(x_2, y_2), c_3, c_{12})$ and NE c_{13} iff NE $\text{MAJ3}(\text{AND2}(x_2, y_2), c_3, c_{12})$ and NE z_0 iff NE q_0 and NE z_1 iff NE q_1 and NE z_2 iff NE q_{11} and NE z_3 iff NE q_{12} and NE z_4 iff NE q_{13} and NE z_5 iff NE c_{13} . Then
- (i) NE z_0 iff NE $\text{MULT}_{310}(x_2, x_1, y_1, x_0, y_0)$,
 - (ii) NE z_1 iff NE $\text{MULT}_{311}(x_2, x_1, y_1, x_0, y_0)$,

- (iii) NE z_2 iff NE $\text{MULT}_{321}(x_2, y_2, x_1, y_1, x_0, y_0)$,
- (iv) NE z_3 iff NE $\text{MULT}_{322}(x_2, y_2, x_1, y_1, x_0, y_0)$,
- (v) NE z_4 iff NE $\text{MULT}_{323}(x_2, y_2, x_1, y_1, x_0, y_0)$, and
- (vi) NE z_5 iff NE $\text{MULT}_{324}(x_2, y_2, x_1, y_1, x_0, y_0)$.

Let a_1, b_1, c be sets. We introduce $\text{CLAADD1}(a_1, b_1, c)$ as a synonym of $\text{XOR3}(a_1, b_1, c)$. We introduce $\text{CLACARR1}(a_1, b_1, c)$ as a synonym of $\text{MAJ3}(a_1, b_1, c)$.

Let a_1, b_1, a_2, b_2, c be sets. The functor $\text{CLAADD2}(a_2, b_2, a_1, b_1, c)$ yields a set and is defined by:

$$\text{(Def. 16)}^1 \quad \text{CLAADD2}(a_2, b_2, a_1, b_1, c) = \text{XOR3}(a_2, b_2, \text{MAJ3}(a_1, b_1, c)).$$

The functor $\text{CLACARR2}(a_2, b_2, a_1, b_1, c)$ yields a set and is defined by:

$$\text{(Def. 17)} \quad \text{CLACARR2}(a_2, b_2, a_1, b_1, c) = \text{OR2}(\text{AND2}(a_2, b_2), \text{AND2}(\text{OR2}(a_2, b_2), \text{MAJ3}(a_1, b_1, c))).$$

Let $a_1, b_1, a_2, b_2, a_3, b_3, c$ be sets. The functor $\text{CLAADD3}(a_3, b_3, a_2, b_2, a_1, b_1, c)$ yielding a set is defined as follows:

$$\text{(Def. 18)} \quad \text{CLAADD3}(a_3, b_3, a_2, b_2, a_1, b_1, c) = \text{XOR3}(a_3, b_3, \text{CLACARR2}(a_2, b_2, a_1, b_1, c)).$$

The functor $\text{CLACARR3}(a_3, b_3, a_2, b_2, a_1, b_1, c)$ yields a set and is defined by:

$$\text{(Def. 19)} \quad \text{CLACARR3}(a_3, b_3, a_2, b_2, a_1, b_1, c) = \text{OR3}(\text{AND2}(a_3, b_3), \text{AND2}(\text{OR2}(a_3, b_3), \text{AND2}(a_2, b_2))), \text{AND3}(\text{OR2}(a_3, b_3), \text{OR2}(a_2, b_2), \text{MAJ3}(a_1, b_1, c))).$$

Let $a_1, b_1, a_2, b_2, a_3, b_3, a_4, b_4, c$ be sets. The functor $\text{CLAADD4}(a_4, b_4, a_3, b_3, a_2, b_2, a_1, b_1, c)$ yielding a set is defined by:

$$\text{(Def. 20)} \quad \text{CLAADD4}(a_4, b_4, a_3, b_3, a_2, b_2, a_1, b_1, c) = \text{XOR3}(a_4, b_4, \text{CLACARR3}(a_3, b_3, a_2, b_2, a_1, b_1, c)).$$

The functor $\text{CLACARR4}(a_4, b_4, a_3, b_3, a_2, b_2, a_1, b_1, c)$ yielding a set is defined as follows:

$$\text{(Def. 21)} \quad \text{CLACARR4}(a_4, b_4, a_3, b_3, a_2, b_2, a_1, b_1, c) = \text{OR4}(\text{AND2}(a_4, b_4), \text{AND2}(\text{OR2}(a_4, b_4), \text{AND2}(a_3, b_3))), \text{AND3}(\text{OR2}(a_4, b_4), \text{OR2}(a_3, b_3), \text{AND2}(a_2, b_2)), \text{AND4}(\text{OR2}(a_4, b_4), \text{OR2}(a_3, b_3), \text{OR2}(a_2, b_2), \text{MAJ3}(a_1, b_1, c))).$$

One can prove the following proposition

- (4) Let $x_0, x_1, x_2, y_0, y_1, y_2, z_0, z_1, z_2, z_3, z_4, z_5, q_0, q_1, q_2, q_3, c_1, c_2, c_3$ be sets such that NE q_0 iff NE $\text{AND2}(x_0, y_0)$ and NE q_1 iff NE $\text{XOR3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE c_1 iff NE $\text{MAJ3}(\text{AND2}(x_1, y_0), \text{AND2}(x_0, y_1), \emptyset)$ and NE q_2 iff NE $\text{XOR3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_0, y_2))$ and NE c_2 iff NE $\text{MAJ3}(\text{AND2}(x_2, y_0), \text{AND2}(x_1, y_1), \text{AND2}(x_0, y_2))$ and NE q_3 iff NE $\text{XOR3}(\text{AND2}(x_2, y_1), \text{AND2}(x_1, y_2), \emptyset)$ and NE c_3 iff NE $\text{MAJ3}(\text{AND2}(x_2,$

¹The definitions (Def. 14) and (Def. 15) have been removed.

y_1), $\text{AND2}(x_1, y_2), \emptyset$) and $\text{NE } z_0$ iff $\text{NE } q_0$ and $\text{NE } z_1$ iff $\text{NE } q_1$ and $\text{NE } z_2$ iff $\text{NE } \text{CLAADD1}(q_2, c_1, \emptyset)$ and $\text{NE } z_3$ iff $\text{NE } \text{CLAADD2}(q_3, c_2, q_2, c_1, \emptyset)$ and $\text{NE } z_4$ iff $\text{NE } \text{CLAADD3}(\text{AND2}(x_2, y_2), c_3, q_3, c_2, q_2, c_1, \emptyset)$ and $\text{NE } z_5$ iff $\text{NE } \text{CLACARR3}(\text{AND2}(x_2, y_2), c_3, q_3, c_2, q_2, c_1, \emptyset)$. Then

- (i) $\text{NE } z_0$ iff $\text{NE } \text{MULT}_{310}(x_2, x_1, y_1, x_0, y_0)$,
- (ii) $\text{NE } z_1$ iff $\text{NE } \text{MULT}_{311}(x_2, x_1, y_1, x_0, y_0)$,
- (iii) $\text{NE } z_2$ iff $\text{NE } \text{MULT}_{321}(x_2, y_2, x_1, y_1, x_0, y_0)$,
- (iv) $\text{NE } z_3$ iff $\text{NE } \text{MULT}_{322}(x_2, y_2, x_1, y_1, x_0, y_0)$,
- (v) $\text{NE } z_4$ iff $\text{NE } \text{MULT}_{323}(x_2, y_2, x_1, y_1, x_0, y_0)$, and
- (vi) $\text{NE } z_5$ iff $\text{NE } \text{MULT}_{324}(x_2, y_2, x_1, y_1, x_0, y_0)$.

REFERENCES

- [1] Yatsuka Nakamura. Logic gates and logical equivalence of adders. *Formalized Mathematics*, 8(1):35–45, 1999.
- [2] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [3] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.

Received August 28, 2000
