

Some Properties of Dyadic Numbers and Intervals

Józef Białaś
Łódź University

Yatsuka Nakamura
Shinshu University
Nagano

Summary. The article is the second part of a paper proving the fundamental Urysohn Theorem concerning the existence of a real valued continuous function on a normal topological space. The paper is divided into two parts. In the first part, we introduce some definitions and theorems concerning properties of intervals; in the second we prove some of properties of dyadic numbers used in proving Urysohn Lemma.

MML Identifier: URYSOHN2.

The terminology and notation used here have been introduced in the following articles: [9], [10], [11], [3], [4], [8], [7], [6], [12], [1], [2], and [5].

The following proposition is true

- (1) For every interval A such that $A \neq \emptyset$ holds if $\inf A < \sup A$, then $\text{vol}(A) = \sup A - \inf A$ and if $\sup A = \inf A$, then $\text{vol}(A) = 0_{\mathbb{R}}$.

Let A be a subset of \mathbb{R} and let x be a real number. The functor $x \cdot A$ yielding a subset of \mathbb{R} is defined as follows:

- (Def. 1) For every real number y holds $y \in x \cdot A$ iff there exists a real number z such that $z \in A$ and $y = x \cdot z$.

Next we state a number of propositions:

- (2) For every subset A of \mathbb{R} and for every real number x such that $x \neq 0$ holds $x^{-1} \cdot (x \cdot A) = A$.
- (3) For every real number x such that $x \neq 0$ and for every subset A of \mathbb{R} such that $A = \mathbb{R}$ holds $x \cdot A = A$.
- (4) For every subset A of \mathbb{R} such that $A \neq \emptyset$ holds $0 \cdot A = \{0\}$.
- (5) For every subset A of \mathbb{R} such that $A \neq \emptyset$ holds $0 \cdot A = \{0\}$.

- (6) For every real number x holds $x \cdot \emptyset = \emptyset$.
- (7) For every real number y holds $y < 0$ or $y = 0$ or $0 < y$.
- (8) Let a, b be extended real numbers. Suppose $a \leq b$. Then $a = -\infty$ and $b = -\infty$ or $a = -\infty$ and $b \in \mathbb{R}$ or $a = -\infty$ and $b = +\infty$ or $a \in \mathbb{R}$ and $b \in \mathbb{R}$ or $a \in \mathbb{R}$ and $b = +\infty$ or $a = +\infty$ and $b = +\infty$.
- (9) For every extended real number x holds $[x, x]$ is an interval.
- (10) For every interval A holds $0 \cdot A$ is an interval.
- (11) For all real numbers q, x such that $x \neq 0$ holds $q = x \cdot \frac{q}{x}$.
- (12) For all real numbers p, q, x such that $0 < x$ and $x \cdot p < x \cdot q$ holds $p < q$.
- (13) For all real numbers p, q, x such that $x < 0$ and $x \cdot p < x \cdot q$ holds $q < p$.
- (14) For all real numbers p, q, x such that $0 < x$ and $x \cdot p \leq x \cdot q$ holds $p \leq q$.
- (15) For all real numbers p, q, x such that $x < 0$ and $x \cdot p \leq x \cdot q$ holds $q \leq p$.
- (16) Let A be an interval and x be a real number. If $x \neq 0$, then if A is open interval, then $x \cdot A$ is open interval.
- (17) Let A be an interval and x be a real number. If $x \neq 0$, then if A is closed interval, then $x \cdot A$ is closed interval.
- (18) Let A be an interval and x be a real number. Suppose $0 < x$. If A is right open interval, then $x \cdot A$ is right open interval.
- (19) Let A be an interval and x be a real number. Suppose $x < 0$. If A is right open interval, then $x \cdot A$ is left open interval.
- (20) Let A be an interval and x be a real number. Suppose $0 < x$. If A is left open interval, then $x \cdot A$ is left open interval.
- (21) Let A be an interval and x be a real number. Suppose $x < 0$. If A is left open interval, then $x \cdot A$ is right open interval.
- (22) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A]$. Then $B = [\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (23) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A]$. Then $B =]\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (24) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A[$. Then $B =]\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (25) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A[$.

Then $B = [\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.

- (26) For every interval A and for every real number x holds $x \cdot A$ is an interval.
Let A be an interval and let x be a real number. Observe that $x \cdot A$ is interval.
The following propositions are true:
- (27) Let A be an interval and x be a real number. If $0 \leq x$, then for every real number y such that $y = \text{vol}(A)$ holds $x \cdot y = \text{vol}(x \cdot A)$.
- (28) For all real numbers x, y, z such that $x < y$ and $y \leq z$ or $x \leq y$ and $y < z$ holds $x < z$.
- (29) For every natural number n holds $n < 2^n$.
- (30) For every integer n such that $0 \leq n$ holds n is a natural number.
- (31) For all natural numbers n, m such that $n < m$ holds $2^n < 2^m$.
- (32) For every real number e_1 such that $0 < e_1$ there exists a natural number n such that $1 < 2^n \cdot e_1$.
- (33) For all real numbers a, b such that $0 \leq a$ and $1 < b - a$ there exists a natural number n such that $a < n$ and $n < b$.
- (34) For every integer n such that $0 < n$ holds n is a natural number.
- (35) For every rational number n such that $0 \leq n$ holds $0 \leq \text{num } n$.
- (36) For every rational number n such that $0 < n$ holds $0 < \text{num } n$.
- (37) For all real numbers a, b, c, d such that $0 < b$ and $0 < d$ or $b < 0$ and $d < 0$ holds if $\frac{a}{b} < \frac{c}{d}$, then $a \cdot d < c \cdot b$.
- (38) For every natural number n holds $\text{dyadic}(n) \subseteq \text{DYADIC}$.
- (39) For all real numbers a, b such that $a < b$ and $0 \leq a$ and $b \leq 1$ there exists a real number c such that $c \in \text{DYADIC}$ and $a < c$ and $c < b$.
- (40) For all real numbers a, b such that $a < b$ there exists a real number c such that $c \in \text{DOM}$ and $a < c$ and $c < b$.
- (41) For every non empty subset A of $\overline{\mathbb{R}}$ and for all extended real numbers a, b such that $A \subseteq [a, b]$ holds $a \leq \inf A$ and $\sup A \leq b$.
- (42) $0 \in \text{DYADIC}$ and $1 \in \text{DYADIC}$.
- (43) For all extended real numbers a, b such that $a = 0$ and $b = 1$ holds $\text{DYADIC} \subseteq [a, b]$.
- (44) For all natural numbers n, k such that $n \leq k$ holds $\text{dyadic}(n) \subseteq \text{dyadic}(k)$.
- (45) For all real numbers a, b, c, d such that $a < c$ and $c < b$ and $a < d$ and $d < b$ holds $|d - c| < b - a$.
- (46) Let e_1 be a real number. Suppose $0 < e_1$. Let d be a real number. Suppose $0 < d$ and $d \leq 1$. Then there exist real numbers r_1, r_2 such that $r_1 \in \text{DYADIC} \cup \mathbb{R}_{>1}$ and $r_2 \in \text{DYADIC} \cup \mathbb{R}_{>1}$ and $0 < r_1$ and $r_1 < d$ and $d < r_2$ and $r_2 - r_1 < e_1$.

REFERENCES

- [1] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Formalized Mathematics*, 2(1):163–171, 1991.
- [2] Józef Białas. Series of positive real numbers. Measure theory. *Formalized Mathematics*, 2(1):173–183, 1991.
- [3] Józef Białas. Properties of the intervals of real numbers. *Formalized Mathematics*, 3(2):263–269, 1992.
- [4] Józef Białas. Some properties of the intervals. *Formalized Mathematics*, 5(1):21–26, 1996.
- [5] Józef Białas and Yatsuka Nakamura. Dyadic numbers and T_4 topological spaces. *Formalized Mathematics*, 5(3):361–366, 1996.
- [6] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [7] Andrzej Kondracki. Basic properties of rational numbers. *Formalized Mathematics*, 1(5):841–845, 1990.
- [8] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [9] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [10] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [11] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [12] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received February 16, 2001
