

On Polynomials with Coefficients in a Ring of Polynomials

Barbara Dzienis
University of Białystok

Summary. The main result of the paper is, that the ring of polynomials with o_1 variables and coefficients in the ring of polynomials with o_2 variables and coefficient in a ring L is isomorphic with the ring with $o_1 + o_2$ variables, and coefficients in L .

MML Identifier: POLYNOM6.

The papers [18], [4], [3], [6], [15], [14], [9], [1], [2], [13], [12], [10], [5], [16], [7], [17], [8], and [11] provide the notation and terminology for this paper.

1. PRELIMINARIES

In this paper o_1, o_2 are ordinal numbers.

Let L_1, L_2 be non empty double loop structures. Let us note that the predicate L_1 is ring isomorphic to L_2 is reflexive. We introduce L_1 and L_2 are isomorphic as a synonym of L_1 is ring isomorphic to L_2 .

We now state the proposition

- (1) Let B be a set. Suppose that for every set x holds $x \in B$ iff there exists an ordinal number o such that $x = o_1 + o$ and $o \in o_2$. Then $o_1 + o_2 = o_1 \cup B$.

Let o_1 be an ordinal number and let o_2 be a non empty ordinal number. Note that $o_1 + o_2$ is non empty and $o_2 + o_1$ is non empty.

One can prove the following proposition

- (2) Let n be an ordinal number and a, b be bags of n . Suppose $a < b$. Then there exists an ordinal number o such that $o \in n$ and $a(o) < b(o)$ and for every ordinal number l such that $l \in o$ holds $a(l) = b(l)$.

2. ABOUT BAGS

Let o_1, o_2 be ordinal numbers, let a be an element of $\text{Bags } o_1$, and let b be an element of $\text{Bags } o_2$. The functor $a + b$ yielding an element of $\text{Bags}(o_1 + o_2)$ is defined as follows:

- (Def. 1) For every ordinal number o holds if $o \in o_1$, then $(a + b)(o) = a(o)$ and if $o \in (o_1 + o_2) \setminus o_1$, then $(a + b)(o) = b(o - o_1)$.

One can prove the following propositions:

- (3) For every element a of $\text{Bags } o_1$ and for every element b of $\text{Bags } o_2$ such that $o_2 = \emptyset$ holds $a + b = a$.
- (4) For every element a of $\text{Bags } o_1$ and for every element b of $\text{Bags } o_2$ such that $o_1 = \emptyset$ holds $a + b = b$.
- (5) For every element b_1 of $\text{Bags } o_1$ and for every element b_2 of $\text{Bags } o_2$ holds $b_1 + b_2 = \text{EmptyBag}(o_1 + o_2)$ iff $b_1 = \text{EmptyBag } o_1$ and $b_2 = \text{EmptyBag } o_2$.
- (6) For every element c of $\text{Bags}(o_1 + o_2)$ there exists an element c_1 of $\text{Bags } o_1$ and there exists an element c_2 of $\text{Bags } o_2$ such that $c = c_1 + c_2$.
- (7) For all elements b_1, c_1 of $\text{Bags } o_1$ and for all elements b_2, c_2 of $\text{Bags } o_2$ such that $b_1 + b_2 = c_1 + c_2$ holds $b_1 = c_1$ and $b_2 = c_2$.
- (8) Let n be an ordinal number, L be an Abelian add-associative right zeroed right complementable distributive associative non empty double loop structure, and p, q, r be serieses of n, L . Then $(p + q) * r = p * r + q * r$.

3. MAIN RESULTS

Let n be an ordinal number and let L be a right zeroed Abelian add-associative right complementable unital distributive associative non trivial non empty double loop structure. Observe that $\text{Polynom-Ring}(n, L)$ is non trivial and distributive.

Let o_1, o_2 be non empty ordinal numbers, let L be a right zeroed add-associative right complementable unital distributive non trivial non empty double loop structure, and let P be a polynomial of $o_1, \text{Polynom-Ring}(o_2, L)$. The functor $\text{Compress } P$ yields a polynomial of $o_1 + o_2, L$ and is defined by the condition (Def. 2).

- (Def. 2) Let b be an element of $\text{Bags}(o_1 + o_2)$. Then there exists an element b_1 of $\text{Bags } o_1$ and there exists an element b_2 of $\text{Bags } o_2$ and there exists a polynomial Q_1 of o_2, L such that $Q_1 = P(b_1)$ and $b = b_1 + b_2$ and $(\text{Compress } P)(b) = Q_1(b_2)$.

Next we state several propositions:

- (9) For all elements b_1, c_1 of Bags o_1 and for all elements b_2, c_2 of Bags o_2 such that $b_1 \mid c_1$ and $b_2 \mid c_2$ holds $b_1 + b_2 \mid c_1 + c_2$.
- (10) Let b be a bag of $o_1 + o_2$, b_1 be an element of Bags o_1 , and b_2 be an element of Bags o_2 . Suppose $b \mid b_1 + b_2$. Then there exists an element c_1 of Bags o_1 and there exists an element c_2 of Bags o_2 such that $c_1 \mid b_1$ and $c_2 \mid b_2$ and $b = c_1 + c_2$.
- (11) For all elements a_1, b_1 of Bags o_1 and for all elements a_2, b_2 of Bags o_2 holds $a_1 + a_2 < b_1 + b_2$ iff $a_1 < b_1$ or $a_1 = b_1$ and $a_2 < b_2$.
- (12) Let b_1 be an element of Bags o_1 , b_2 be an element of Bags o_2 , and G be a finite sequence of elements of $(\text{Bags}(o_1 + o_2))^*$. Suppose that
 - (i) $\text{dom } G = \text{Seg len divisors } b_1$, and
 - (ii) for every natural number i such that $i \in \text{Seg len divisors } b_1$ there exists an element a'_1 of Bags o_1 and there exists a finite sequence F_1 of elements of $\text{Bags}(o_1 + o_2)$ such that $F_1 = G_i$ and $\pi_i \text{ divisors } b_1 = a'_1$ and $\text{len } F_1 = \text{len divisors } b_2$ and for every natural number m such that $m \in \text{dom } F_1$ there exists an element a''_1 of Bags o_2 such that $\pi_m \text{ divisors } b_2 = a''_1$ and $\pi_m F_1 = a'_1 + a''_1$.
 Then $\text{divisors}(b_1 + b_2) = \text{Flat}(G)$.
- (13) For all elements a_1, b_1, c_1 of Bags o_1 and for all elements a_2, b_2, c_2 of Bags o_2 such that $c_1 = b_1 -' a_1$ and $c_2 = b_2 -' a_2$ holds $(b_1 + b_2) -' (a_1 + a_2) = c_1 + c_2$.
- (14) Let b_1 be an element of Bags o_1 , b_2 be an element of Bags o_2 , and G be a finite sequence of elements of $(\text{Bags}(o_1 + o_2))^2$. Suppose that
 - (i) $\text{dom } G = \text{Seg len decomp } b_1$, and
 - (ii) for every natural number i such that $i \in \text{Seg len decomp } b_1$ there exist elements a'_1, b'_1 of Bags o_1 and there exists a finite sequence F_1 of elements of $(\text{Bags}(o_1 + o_2))^2$ such that $F_1 = G_i$ and $\pi_i \text{ decomp } b_1 = \langle a'_1, b'_1 \rangle$ and $\text{len } F_1 = \text{len decomp } b_2$ and for every natural number m such that $m \in \text{dom } F_1$ there exist elements a''_1, b''_1 of Bags o_2 such that $\pi_m \text{ decomp } b_2 = \langle a''_1, b''_1 \rangle$ and $\pi_m F_1 = \langle a'_1 + a''_1, b'_1 + b''_1 \rangle$.
 Then $\text{decomp}(b_1 + b_2) = \text{Flat}(G)$.
- (15) Let o_1, o_2 be non empty ordinal numbers and L be an Abelian right zeroed add-associative right complementable unital distributive associative well unital non trivial non empty double loop structure. Then $\text{Polynom-Ring}(o_1, \text{Polynom-Ring}(o_2, L))$ and $\text{Polynom-Ring}(o_1 + o_2, L)$ are isomorphic.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Grzegorz Bancerek. Sequences of ordinal numbers. *Formalized Mathematics*, 1(2):281–290, 1990.

- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Formalized Mathematics*, 4(1):91–101, 1993.
- [5] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(3):433–439, 1990.
- [6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [7] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [8] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [9] Robert Milewski. Associated matrix of linear map. *Formalized Mathematics*, 5(3):339–345, 1996.
- [10] Robert Milewski. The ring of polynomials. *Formalized Mathematics*, 9(2):339–346, 2001.
- [11] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(1):3–11, 1991.
- [12] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Formalized Mathematics*, 9(1):95–110, 2001.
- [13] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [14] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [15] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [16] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [17] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [18] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.

Received August 10, 2001
