

# On the Decompositions of Intervals and Simple Closed Curves<sup>1</sup>

Adam Grabowski  
University of Białystok

**Summary.** The aim of the paper is to show that the only subcontinua of the Jordan curve are arcs, the whole curve, and singletons of its points. Additionally, it has been shown that the only subcontinua of the unit interval  $\mathbb{I}$  are closed intervals.

MML Identifier: BORSUK\_4.

The articles [21], [23], [13], [24], [2], [1], [3], [25], [19], [6], [4], [20], [8], [10], [11], [15], [26], [18], [22], [14], [16], [9], [17], [5], [12], and [7] provide the terminology and notation for this paper.

## 1. PRELIMINARIES

Let us note that every simple closed curve is non trivial.

Let  $T$  be a non empty topological space. One can check that there exists a subset of  $T$  which is non empty, compact, and connected.

Let us observe that every element of the carrier of  $\mathbb{I}$  is real.

Next we state two propositions:

- (1) Let  $X$  be a non empty set and  $A, B$  be non empty subsets of  $X$ . If  $A \subset B$ , then there exists an element  $p$  of  $X$  such that  $p \in B$  and  $A \subseteq B \setminus \{p\}$ .
- (2) Let  $X$  be a non empty set and  $A$  be a non empty subset of  $X$ . Then  $A$  is trivial if and only if there exists an element  $x$  of  $X$  such that  $A = \{x\}$ .

---

<sup>1</sup>This work has been partially supported by the European Community TYPES grant IST-1999-29001 and CALCULEMUS grant HPRN-CT-2000-00102.

Let  $T$  be a non trivial 1-sorted structure. Observe that there exists a subset of the carrier of  $T$  which is non trivial.

The following proposition is true

- (3) For every non trivial set  $X$  and for every set  $p$  there exists an element  $q$  of  $X$  such that  $q \neq p$ .

Let  $X$  be a non trivial set. Observe that there exists a subset of  $X$  which is non trivial.

We now state a number of propositions:

- (4) Let  $T$  be a non trivial set,  $X$  be a non trivial subset of  $T$ , and  $p$  be a set. Then there exists an element  $q$  of  $T$  such that  $q \in X$  and  $q \neq p$ .
- (5) Let  $f, g$  be functions and  $a$  be a set. Suppose  $f$  is one-to-one and  $g$  is one-to-one and  $\text{dom } f \cap \text{dom } g = \{a\}$  and  $\text{rng } f \cap \text{rng } g = \{f(a)\}$ . Then  $f + \cdot g$  is one-to-one.
- (6) Let  $f, g$  be functions and  $a$  be a set. Suppose  $f$  is one-to-one and  $g$  is one-to-one and  $\text{dom } f \cap \text{dom } g = \{a\}$  and  $\text{rng } f \cap \text{rng } g = \{f(a)\}$  and  $f(a) = g(a)$ . Then  $(f + \cdot g)^{-1} = f^{-1} + \cdot g^{-1}$ .
- (7) Let  $n$  be a natural number,  $A$  be a non empty subset of the carrier of  $\mathcal{E}_T^n$ , and  $p, q$  be points of  $\mathcal{E}_T^n$ . If  $A$  is an arc from  $p$  to  $q$ , then  $A \setminus \{p\}$  is non empty.
- (8) For every natural number  $n$  and for all points  $a, b$  of  $\mathcal{E}_T^n$  holds  $\mathcal{L}(a, b)$  is  $n$ -convex.
- (9) For all real numbers  $s_1, s_3, s_4, l$  such that  $s_1 \leq s_3$  and  $s_1 < s_4$  and  $0 \leq l$  and  $l \leq 1$  holds  $s_1 \leq (1 - l) \cdot s_3 + l \cdot s_4$ .
- (10) For every set  $x$  and for all real numbers  $a, b$  such that  $a \leq b$  and  $x \in [a, b]$  holds  $x \in ]a, b[$  or  $x = a$  or  $x = b$ .
- (11) For all real numbers  $a, b, c, d$  such that  $]a, b[$  meets  $[c, d]$  holds  $b > c$ .
- (12) For all real numbers  $a, b, c, d$  such that  $b \leq c$  holds  $[a, b]$  misses  $]c, d[$ .
- (13) For all real numbers  $a, b, c, d$  such that  $b \leq c$  holds  $]a, b[$  misses  $[c, d]$ .
- (14) For all real numbers  $a, b, c, d$  such that  $a < b$  and  $[a, b] \subseteq [c, d]$  holds  $c \leq a$  and  $b \leq d$ .
- (15) For all real numbers  $a, b, c, d$  such that  $a < b$  and  $]a, b[ \subseteq [c, d]$  holds  $c \leq a$  and  $b \leq d$ .
- (16) For all real numbers  $a, b, c, d$  such that  $a < b$  and  $]a, b[ \subseteq [c, d]$  holds  $[a, b] \subseteq [c, d]$ .
- (17) Let  $A$  be a subset of the carrier of  $\mathbb{I}$  and  $a, b$  be real numbers. If  $a < b$  and  $A = ]a, b[$ , then  $[a, b] \subseteq$  the carrier of  $\mathbb{I}$ .
- (18) Let  $A$  be a subset of the carrier of  $\mathbb{I}$  and  $a, b$  be real numbers. If  $a < b$  and  $A = ]a, b]$ , then  $[a, b] \subseteq$  the carrier of  $\mathbb{I}$ .

- (19) Let  $A$  be a subset of the carrier of  $\mathbb{I}$  and  $a, b$  be real numbers. If  $a < b$  and  $A = [a, b[$ , then  $[a, b] \subseteq$  the carrier of  $\mathbb{I}$ .
- (20) For all real numbers  $a, b$  such that  $a \neq b$  holds  $\overline{[a, b]} = [a, b]$ .
- (21) For all real numbers  $a, b$  such that  $a \neq b$  holds  $\overline{]a, b[} = [a, b]$ .
- (22) For every subset  $A$  of  $\mathbb{I}$  and for all real numbers  $a, b$  such that  $a < b$  and  $A = ]a, b[$  holds  $\overline{A} = [a, b]$ .
- (23) For every subset  $A$  of the carrier of  $\mathbb{I}$  and for all real numbers  $a, b$  such that  $a < b$  and  $A = ]a, b]$  holds  $\overline{A} = [a, b]$ .
- (24) For every subset  $A$  of the carrier of  $\mathbb{I}$  and for all real numbers  $a, b$  such that  $a < b$  and  $A = [a, b[$  holds  $\overline{A} = [a, b]$ .
- (25) For all real numbers  $a, b$  such that  $a < b$  holds  $[a, b] \neq ]a, b[$ .
- (26) For all real numbers  $a, b$  holds  $[a, b[$  misses  $\{b\}$  and  $]a, b]$  misses  $\{a\}$ .
- (27) For all real numbers  $a, b$  such that  $a \leq b$  holds  $[a, b] \setminus \{a\} = ]a, b]$ .
- (28) For all real numbers  $a, b$  such that  $a \leq b$  holds  $[a, b] \setminus \{b\} = [a, b[$ .
- (29) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $]a, b] \cap [b, c[ = \{b\}$ .
- (30) For all real numbers  $a, b, c$  holds  $[a, b[$  misses  $[b, c]$  and  $]a, b]$  misses  $]b, c[$ .
- (31) For all real numbers  $a, b, c$  such that  $a \leq b$  and  $b \leq c$  holds  $[a, c] \setminus \{b\} = [a, b[ \cup ]b, c]$ .
- (32) Let  $A$  be a subset of the carrier of  $\mathbb{I}$  and  $a, b$  be real numbers. If  $a \leq b$  and  $A = [a, b]$ , then  $0 \leq a$  and  $b \leq 1$ .
- (33) Let  $A, B$  be subsets of  $\mathbb{I}$  and  $a, b, c$  be real numbers. If  $a < b$  and  $b < c$  and  $A = [a, b[$  and  $B = ]b, c]$ , then  $A$  and  $B$  are separated.
- (34) For all real numbers  $a, b$  such that  $a \leq b$  holds  $[a, b] = [a, b[ \cup \{b\}$ .
- (35) For all real numbers  $a, b$  such that  $a \leq b$  holds  $[a, b] = \{a\} \cup ]a, b]$ .
- (36) For all real numbers  $a, b, c, d$  such that  $a \leq b$  and  $b < c$  and  $c \leq d$  holds  $[a, d] = [a, b] \cup ]b, c[ \cup [c, d]$ .
- (37) For all real numbers  $a, b, c, d$  such that  $a \leq b$  and  $b < c$  and  $c \leq d$  holds  $[a, d] \setminus ([a, b] \cup [c, d]) = ]b, c[$ .
- (38) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $]a, b] \cup ]b, c[ = ]a, c[$ .
- (39) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $[b, c[ \subseteq ]a, c[$ .
- (40) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $]a, b] \cup [b, c[ = ]a, c[$ .
- (41) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $]a, c[ \setminus ]a, b] = ]b, c[$ .
- (42) For all real numbers  $a, b, c$  such that  $a < b$  and  $b < c$  holds  $]a, c[ \setminus [b, c] = ]a, b[$ .

- (43) For all points  $p_1, p_2$  of  $\mathbb{I}$  holds  $[p_1, p_2]$  is a subset of  $\mathbb{I}$ .  
 (44) For all points  $a, b$  of  $\mathbb{I}$  holds  $]a, b[$  is a subset of  $\mathbb{I}$ .

## 2. DECOMPOSITIONS OF INTERVALS

The following propositions are true:

- (45) For every real number  $p$  holds  $\{p\}$  is a closed-interval subset of  $\mathbb{R}$ .  
 (46) Let  $A$  be a non empty connected subset of  $\mathbb{I}$  and  $a, b, c$  be points of  $\mathbb{I}$ . If  $a \leq b$  and  $b \leq c$  and  $a \in A$  and  $c \in A$ , then  $b \in A$ .  
 (47) For every non empty connected subset  $A$  of  $\mathbb{I}$  and for all real numbers  $a, b$  such that  $a \in A$  and  $b \in A$  holds  $[a, b] \subseteq A$ .  
 (48) For all real numbers  $a, b$  and for every subset  $A$  of  $\mathbb{I}$  such that  $a \leq b$  and  $A = [a, b]$  holds  $A$  is closed.  
 (49) For all points  $p_1, p_2$  of  $\mathbb{I}$  such that  $p_1 \leq p_2$  holds  $[p_1, p_2]$  is a non empty compact connected subset of  $\mathbb{I}$ .  
 (50) Let  $X$  be a subset of the carrier of  $\mathbb{I}$  and  $X'$  be a subset of  $\mathbb{R}$ . If  $X' = X$ , then  $X'$  is upper bounded and lower bounded.  
 (51) Let  $X$  be a subset of the carrier of  $\mathbb{I}$ ,  $X'$  be a subset of  $\mathbb{R}$ , and  $x$  be a real number. If  $x \in X'$  and  $X' = X$ , then  $\inf X' \leq x$  and  $x \leq \sup X'$ .  
 (52) For every subset  $A$  of  $\mathbb{R}$  and for every subset  $B$  of  $\mathbb{I}$  such that  $A = B$  holds  $A$  is closed iff  $B$  is closed.  
 (53) For every closed-interval subset  $C$  of  $\mathbb{R}$  holds  $\inf C \leq \sup C$ .  
 (54) Let  $C$  be a non empty compact connected subset of  $\mathbb{I}$  and  $C'$  be a subset of  $\mathbb{R}$ . If  $C = C'$  and  $[\inf C', \sup C'] \subseteq C'$ , then  $[\inf C', \sup C'] = C'$ .  
 (55) Every non empty compact connected subset of  $\mathbb{I}$  is a closed-interval subset of  $\mathbb{R}$ .  
 (56) For every non empty compact connected subset  $C$  of  $\mathbb{I}$  there exist points  $p_1, p_2$  of  $\mathbb{I}$  such that  $p_1 \leq p_2$  and  $C = [p_1, p_2]$ .

## 3. DECOMPOSITIONS OF SIMPLE CLOSED CURVES

The strict non empty subspace  $I(01)$  of  $\mathbb{I}$  is defined as follows:

(Def. 1) The carrier of  $I(01) = ]0, 1[$ .

One can prove the following propositions:

- (57) For every subset  $A$  of  $\mathbb{I}$  such that  $A =$  the carrier of  $I(01)$  holds  $I(01) = \mathbb{I} \upharpoonright A$ .  
 (58) The carrier of  $I(01) =$  (the carrier of  $\mathbb{I}) \setminus \{0, 1\}$ .  
 (59)  $I(01)$  is an open subspace of  $\mathbb{I}$ .

- (60) For every real number  $r$  holds  $r \in$  the carrier of  $I(01)$  iff  $0 < r$  and  $r < 1$ .
- (61) For all points  $a, b$  of  $\mathbb{I}$  such that  $a < b$  and  $b \neq 1$  holds  $]a, b]$  is a non empty subset of  $I(01)$ .
- (62) For all points  $a, b$  of  $\mathbb{I}$  such that  $a < b$  and  $a \neq 0$  holds  $[a, b[$  is a non empty subset of  $I(01)$ .
- (63) For every simple closed curve  $D$  holds  $(\mathcal{E}_T^2) \upharpoonright \square_{\mathcal{E}^2}$  and  $(\mathcal{E}_T^2) \upharpoonright D$  are homeomorphic.
- (64) Let  $D$  be a non empty subset of  $\mathcal{E}_T^2$  and  $p_1, p_2$  be points of  $\mathcal{E}_T^2$ . If  $D$  is an arc from  $p_1$  to  $p_2$ , then  $I(01)$  and  $(\mathcal{E}_T^2) \upharpoonright (D \setminus \{p_1, p_2\})$  are homeomorphic.
- (65) Let  $D$  be a subset of the carrier of  $\mathcal{E}_T^2$  and  $p_1, p_2$  be points of  $\mathcal{E}_T^2$ . If  $D$  is an arc from  $p_1$  to  $p_2$ , then  $\mathbb{I}$  and  $(\mathcal{E}_T^2) \upharpoonright D$  are homeomorphic.
- (66) For all points  $p_1, p_2$  of  $\mathcal{E}_T^2$  such that  $p_1 \neq p_2$  holds  $\mathbb{I}$  and  $(\mathcal{E}_T^2) \upharpoonright \mathcal{L}(p_1, p_2)$  are homeomorphic.
- (67) Let  $E$  be a subset of  $I(01)$ . Given points  $p_1, p_2$  of  $\mathbb{I}$  such that  $p_1 < p_2$  and  $E = [p_1, p_2]$ . Then  $\mathbb{I}$  and  $I(01) \upharpoonright E$  are homeomorphic.
- (68) Let  $A$  be a non empty subset of the carrier of  $\mathcal{E}_T^2$ ,  $p, q$  be points of  $\mathcal{E}_T^2$ , and  $a, b$  be points of  $\mathbb{I}$ . Suppose  $A$  is an arc from  $p$  to  $q$  and  $a < b$ . Then there exists a non empty subset  $E$  of  $\mathbb{I}$  and there exists a map  $f$  from  $\mathbb{I} \upharpoonright E$  into  $(\mathcal{E}_T^2) \upharpoonright A$  such that  $E = [a, b]$  and  $f$  is a homeomorphism and  $f(a) = p$  and  $f(b) = q$ .
- (69) Let  $A$  be a topological space,  $B$  be a non empty topological space,  $f$  be a map from  $A$  into  $B$ ,  $C$  be a topological space, and  $X$  be a subset of  $A$ . Suppose  $f$  is continuous and  $C$  is a subspace of  $B$ . Let  $h$  be a map from  $A \upharpoonright X$  into  $C$ . If  $h = f \upharpoonright X$ , then  $h$  is continuous.
- (70) For every subset  $X$  of  $\mathbb{I}$  and for all points  $a, b$  of  $\mathbb{I}$  such that  $a \leq b$  and  $X = ]a, b[$  holds  $X$  is open.
- (71) For every subset  $X$  of  $I(01)$  and for all points  $a, b$  of  $\mathbb{I}$  such that  $a \leq b$  and  $X = ]a, b[$  holds  $X$  is open.
- (72) For every non empty subset  $X$  of  $I(01)$  and for every point  $a$  of  $\mathbb{I}$  such that  $0 < a$  and  $X = ]0, a]$  holds  $X$  is closed.
- (73) For every non empty subset  $X$  of  $I(01)$  and for every point  $a$  of  $\mathbb{I}$  such that  $X = [a, 1[$  holds  $X$  is closed.
- (74) Let  $A$  be a non empty subset of the carrier of  $\mathcal{E}_T^2$ ,  $p, q$  be points of  $\mathcal{E}_T^2$ , and  $a, b$  be points of  $\mathbb{I}$ . Suppose  $A$  is an arc from  $p$  to  $q$  and  $a < b$  and  $b \neq 1$ . Then there exists a non empty subset  $E$  of  $I(01)$  and there exists a map  $f$  from  $I(01) \upharpoonright E$  into  $(\mathcal{E}_T^2) \upharpoonright (A \setminus \{p\})$  such that  $E = ]a, b]$  and  $f$  is a homeomorphism and  $f(b) = q$ .
- (75) Let  $A$  be a non empty subset of the carrier of  $\mathcal{E}_T^2$ ,  $p, q$  be points of  $\mathcal{E}_T^2$ , and  $a, b$  be points of  $\mathbb{I}$ . Suppose  $A$  is an arc from  $p$  to  $q$  and  $a < b$  and  $a \neq 0$ . Then there exists a non empty subset  $E$  of  $I(01)$  and there exists

a map  $f$  from  $I(01) \upharpoonright E$  into  $(\mathcal{E}_T^2) \upharpoonright (A \setminus \{q\})$  such that  $E = [a, b[$  and  $f$  is a homeomorphism and  $f(a) = p$ .

- (76) Let  $A, B$  be non empty subsets of the carrier of  $\mathcal{E}_T^2$  and  $p, q$  be points of  $\mathcal{E}_T^2$ . Suppose  $A$  is an arc from  $p$  to  $q$  and  $B$  is an arc from  $q$  to  $p$  and  $A \cap B = \{p, q\}$  and  $p \neq q$ . Then  $I(01)$  and  $(\mathcal{E}_T^2) \upharpoonright ((A \setminus \{p\}) \cup (B \setminus \{q\}))$  are homeomorphic.
- (77) For every simple closed curve  $D$  and for every point  $p$  of  $\mathcal{E}_T^2$  such that  $p \in D$  holds  $(\mathcal{E}_T^2) \upharpoonright (D \setminus \{p\})$  and  $I(01)$  are homeomorphic.
- (78) Let  $D$  be a simple closed curve and  $p, q$  be points of  $\mathcal{E}_T^2$ . If  $p \in D$  and  $q \in D$ , then  $(\mathcal{E}_T^2) \upharpoonright (D \setminus \{p\})$  and  $(\mathcal{E}_T^2) \upharpoonright (D \setminus \{q\})$  are homeomorphic.
- (79) Let  $C$  be a non empty subset of  $\mathcal{E}_T^2$  and  $E$  be a subset of  $I(01)$ . Suppose there exist points  $p_1, p_2$  of  $\mathbb{I}$  such that  $p_1 < p_2$  and  $E = [p_1, p_2]$  and  $I(01) \upharpoonright E$  and  $(\mathcal{E}_T^2) \upharpoonright C$  are homeomorphic. Then there exist points  $s_1, s_2$  of  $\mathcal{E}_T^2$  such that  $C$  is an arc from  $s_1$  to  $s_2$ .
- (80) Let  $D_1$  be a non empty subset of  $\mathcal{E}_T^2$ ,  $f$  be a map from  $(\mathcal{E}_T^2) \upharpoonright D_1$  into  $I(01)$ , and  $C$  be a non empty subset of  $\mathcal{E}_T^2$ . Suppose  $f$  is a homeomorphism and  $C \subseteq D_1$  and there exist points  $p_1, p_2$  of  $\mathbb{I}$  such that  $p_1 < p_2$  and  $f \circ C = [p_1, p_2]$ . Then there exist points  $s_1, s_2$  of  $\mathcal{E}_T^2$  such that  $C$  is an arc from  $s_1$  to  $s_2$ .
- (81) Let  $D$  be a simple closed curve and  $C$  be a non empty compact connected subset of  $\mathcal{E}_T^2$ . Suppose  $C \subseteq D$ . Then  $C = D$  or there exist points  $p_1, p_2$  of  $\mathcal{E}_T^2$  such that  $C$  is an arc from  $p_1$  to  $p_2$  or there exists a point  $p$  of  $\mathcal{E}_T^2$  such that  $C = \{p\}$ .

## REFERENCES

- [1] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(3):433–439, 1990.
- [2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [4] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [5] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in  $\mathcal{E}^2$ . *Formalized Mathematics*, 6(3):427–440, 1997.
- [6] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(2):383–386, 1990.
- [7] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257–261, 1990.
- [8] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [9] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces - fundamental concepts. *Formalized Mathematics*, 2(4):605–608, 1991.
- [10] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Arcs, line segments and special polygonal arcs. *Formalized Mathematics*, 2(5):617–621, 1991.
- [11] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Simple closed curves. *Formalized Mathematics*, 2(5):663–664, 1991.
- [12] Noboru Endou and Artur Korniłowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(1):93–102, 1999.

- [13] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [14] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(5):665–674, 1991.
- [15] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [16] Yatsuka Nakamura. Half open intervals in real numbers. *Formalized Mathematics*, 10(1):21–22, 2002.
- [17] Yatsuka Nakamura, Andrzej Trybulec, and Czesław Byliński. Bounded domains and unbounded domains. *Formalized Mathematics*, 8(1):1–13, 1999.
- [18] Beata Padlewska. Connected spaces. *Formalized Mathematics*, 1(1):239–244, 1990.
- [19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [20] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [22] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Formalized Mathematics*, 2(4):535–545, 1991.
- [23] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [24] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [25] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [26] Mariusz Żynel and Adam Guzowski.  $T_0$  topological spaces. *Formalized Mathematics*, 5(1):75–77, 1996.

*Received August 7, 2002*

---