

Order Sorted Algebras¹

Josef Urban
Charles University
Praha

Summary. Initial notions for order sorted algebras.

MML Identifier: OSALG.1.

The articles [9], [13], [14], [4], [15], [5], [8], [7], [2], [3], [1], [10], [12], [11], and [6] provide the notation and terminology for this paper.

1. PRELIMINARIES

In this paper i is a set.

Let I be a set, let f be a many sorted set indexed by I , and let p be a finite sequence of elements of I . One can check that $f \cdot p$ is finite sequence-like.

Let S be a non empty many sorted signature. A sort symbol of S is an element of S .

Let S be a non empty many sorted signature.

(Def. 1) An element of the operation symbols of S is said to be an operation symbol of S .

Let S be a non void non empty many sorted signature and let o be an operation symbol of S . Then the result sort of o is an element of S .

Let X be a set. Then Δ_X is an order in X . We introduce Δ_X^o as a synonym of Δ_X .

Let X be a set. Then Δ_X is an equivalence relation of X . We introduce Δ_X^r as a synonym of Δ_X .

¹This work was done during author's research visit in Białystok, funded by the CALCULEMUS grant HPRN-CT-2000-00102.

We introduce overloaded many sorted signatures which are extensions of many sorted signature and are systems

\langle a carrier, operation symbols, an overloading, an arity, a result sort \rangle ,

where the carrier is a set, the operation symbols constitute a set, the overloading is an equivalence relation of the operation symbols, the arity is a function from the operation symbols into the carrier*, and the result sort is a function from the operation symbols into the carrier.

We introduce relation sorted signatures which are extensions of many sorted signature and relational structure and are systems

\langle a carrier, an internal relation, operation symbols, an arity, a result sort \rangle ,

where the carrier is a set, the internal relation is a binary relation on the carrier, the operation symbols constitute a set, the arity is a function from the operation symbols into the carrier*, and the result sort is a function from the operation symbols into the carrier.

We consider overloaded relation sorted signatures as extensions of overloaded many sorted signature and relation sorted signature as systems

\langle a carrier, an internal relation, operation symbols, an overloading, an arity, a result sort \rangle ,

where the carrier is a set, the internal relation is a binary relation on the carrier, the operation symbols constitute a set, the overloading is an equivalence relation of the operation symbols, the arity is a function from the operation symbols into the carrier*, and the result sort is a function from the operation symbols into the carrier.

For simplicity, we use the following convention: A, O are non empty sets, R is an order in A , O_1 is an equivalence relation of O , f is a function from O into A^* , and g is a function from O into A .

One can prove the following proposition

- (1) $\langle A, R, O, O_1, f, g \rangle$ is non empty, non void, reflexive, transitive, and anti-symmetric.

Let us consider A, R, O, O_1, f, g . One can verify that $\langle A, R, O, O_1, f, g \rangle$ is strict, non empty, reflexive, transitive, and antisymmetric.

2. THE NOTIONS: ORDER-SORTED, DISCERNABLE, OP-DISCRETE

In the sequel S is an overloaded relation sorted signature.

Let us consider S . We say that S is order-sorted if and only if:

- (Def. 2) S is reflexive, transitive, and antisymmetric.

Let us note that every overloaded relation sorted signature which is order-sorted is also reflexive, transitive, and antisymmetric and there exists an overloaded relation sorted signature which is strict, non empty, non void, and order-sorted.

Let us observe that there exists an overloaded many sorted signature which is non empty and non void.

Let S be a non empty non void overloaded many sorted signature and let x, y be operation symbols of S . The predicate $x \cong y$ is defined by:

(Def. 3) $\langle x, y \rangle \in$ the overloading of S .

Let us notice that the predicate $x \cong y$ is reflexive and symmetric.

One can prove the following proposition

(2) Let S be a non empty non void overloaded many sorted signature and o, o_1, o_2 be operation symbols of S . If $o \cong o_1$ and $o_1 \cong o_2$, then $o \cong o_2$.

Let S be a non empty non void overloaded many sorted signature. We say that S is discernable if and only if the condition (Def. 4) is satisfied.

(Def. 4) Let x, y be operation symbols of S . Suppose $x \cong y$ and $\text{Arity}(x) = \text{Arity}(y)$ and the result sort of $x =$ the result sort of y . Then $x = y$.

We say that S is op-discrete if and only if:

(Def. 5) The overloading of $S = \Delta_{\text{the operation symbols of } S}^r$.

The following two propositions are true:

(3) Let S be a non empty non void overloaded many sorted signature. Then S is op-discrete if and only if for all operation symbols x, y of S such that $x \cong y$ holds $x = y$.

(4) For every non empty non void overloaded many sorted signature S such that S is op-discrete holds S is discernable.

3. ORDER SORTED SIGNATURE

In the sequel S_0 is a non empty non void many sorted signature.

Let us consider S_0 . The functor $\text{OSSign } S_0$ yields a strict non empty non void order-sorted overloaded relation sorted signature and is defined by the conditions (Def. 6).

- (Def. 6)(i) The carrier of $S_0 =$ the carrier of $\text{OSSign } S_0$,
- (ii) $\Delta_{\text{the carrier of } S_0} =$ the internal relation of $\text{OSSign } S_0$,
- (iii) the operation symbols of $S_0 =$ the operation symbols of $\text{OSSign } S_0$,
- (iv) $\Delta_{\text{the operation symbols of } S_0} =$ the overloading of $\text{OSSign } S_0$,
- (v) the arity of $S_0 =$ the arity of $\text{OSSign } S_0$, and
- (vi) the result sort of $S_0 =$ the result sort of $\text{OSSign } S_0$.

Next we state the proposition

(5) $\text{OSSign } S_0$ is discrete and op-discrete.

Let us mention that there exists a strict non empty non void order-sorted overloaded relation sorted signature which is discrete, op-discrete, and discernable.

Let us observe that every non empty non void overloaded relation sorted signature which is op-discrete is also discernable.

Let us consider S_0 . Observe that $\text{OSSign } S_0$ is discrete and op-discrete.

An order sorted signature is a discernable non empty non void order-sorted overloaded relation sorted signature.

We use the following convention: S is a non empty poset, s_1, s_2 are elements of S , and w_1, w_2 are elements of $(\text{the carrier of } S)^*$.

Let us consider S and let w_1, w_2 be elements of $(\text{the carrier of } S)^*$. The predicate $w_1 \leq w_2$ is defined as follows:

- (Def. 7) $\text{len } w_1 = \text{len } w_2$ and for every set i such that $i \in \text{dom } w_1$ and for all s_1, s_2 such that $s_1 = w_1(i)$ and $s_2 = w_2(i)$ holds $s_1 \leq s_2$.

Let us note that the predicate $w_1 \leq w_2$ is reflexive.

We now state two propositions:

- (6) For all elements w_1, w_2 of $(\text{the carrier of } S)^*$ such that $w_1 \leq w_2$ and $w_2 \leq w_1$ holds $w_1 = w_2$.
- (7) If S is discrete and $w_1 \leq w_2$, then $w_1 = w_2$.

We follow the rules: S is an order sorted signature, o, o_1, o_2 are operation symbols of S , and w_1 is an element of $(\text{the carrier of } S)^*$.

One can prove the following proposition

- (8) If S is discrete and $o_1 \cong o_2$ and $\text{Arity}(o_1) \leq \text{Arity}(o_2)$ and the result sort of $o_1 \leq$ the result sort of o_2 , then $o_1 = o_2$.

Let us consider S and let us consider o . We say that o is monotone if and only if:

- (Def. 8) For every o_2 such that $o \cong o_2$ and $\text{Arity}(o) \leq \text{Arity}(o_2)$ holds the result sort of $o \leq$ the result sort of o_2 .

Let us consider S . We say that S is monotone if and only if:

- (Def. 9) Every operation symbol of S is monotone.

The following proposition is true

- (9) If S is op-discrete, then S is monotone.

Let us observe that there exists an order sorted signature which is monotone.

Let S be a monotone order sorted signature. Observe that there exists an operation symbol of S which is monotone.

Let S be a monotone order sorted signature. One can check that every operation symbol of S is monotone.

One can check that every order sorted signature which is op-discrete is also monotone.

We now state the proposition

- (10) If S is monotone and $\text{Arity}(o_1) = \emptyset$ and $o_1 \cong o_2$ and $\text{Arity}(o_2) = \emptyset$, then $o_1 = o_2$.

Let us consider S, o, o_1, w_1 . We say that o_1 has least args for o, w_1 if and only if:

- (Def. 10) $o \cong o_1$ and $w_1 \leq \text{Arity}(o_1)$ and for every o_2 such that $o \cong o_2$ and $w_1 \leq \text{Arity}(o_2)$ holds $\text{Arity}(o_1) \leq \text{Arity}(o_2)$.

We say that o_1 has least sort for o, w_1 if and only if:

- (Def. 11) $o \cong o_1$ and $w_1 \leq \text{Arity}(o_1)$ and for every o_2 such that $o \cong o_2$ and $w_1 \leq \text{Arity}(o_2)$ holds the result sort of $o_1 \leq$ the result sort of o_2 .

Let us consider S, o, o_1, w_1 . We say that o_1 has least rank for o, w_1 if and only if:

- (Def. 12) o_1 has least args for o, w_1 and least sort for o, w_1 .

Let us consider S, o . We say that o is regular if and only if:

- (Def. 13) o is monotone and for every w_1 such that $w_1 \leq \text{Arity}(o)$ holds there exists o_1 which has least args for o, w_1 .

Let S_1 be a monotone order sorted signature. We say that S_1 is regular if and only if:

- (Def. 14) Every operation symbol of S_1 is regular.

In the sequel S_1 is a monotone order sorted signature, o, o_1 are operation symbols of S_1 , and w_1 is an element of (the carrier of S_1)*.

We now state two propositions:

- (11) S_1 is regular iff for all o, w_1 such that $w_1 \leq \text{Arity}(o)$ holds there exists o_1 which has least rank for o, w_1 .
- (12) For every monotone order sorted signature S_1 such that S_1 is op-discrete holds S_1 is regular.

One can verify that there exists a monotone order sorted signature which is regular.

Let us mention that every monotone order sorted signature which is op-discrete is also regular.

Let S_2 be a regular monotone order sorted signature. One can verify that every operation symbol of S_2 is regular.

We adopt the following rules: S_2 is a regular monotone order sorted signature, o, o_3, o_4 are operation symbols of S_2 , and w_1 is an element of (the carrier of S_2)*.

One can prove the following proposition

- (13) If $w_1 \leq \text{Arity}(o)$ and o_3 has least args for o, w_1 and o_4 has least args for o, w_1 , then $o_3 = o_4$.

Let us consider S_2, o, w_1 . Let us assume that $w_1 \leq \text{Arity}(o)$. The functor $\text{LBound}(o, w_1)$ yields an operation symbol of S_2 and is defined as follows:

- (Def. 15) $\text{LBound}(o, w_1)$ has least args for o, w_1 .

One can prove the following proposition

- (14) For every w_1 such that $w_1 \leq \text{Arity}(o)$ holds $\text{LBound}(o, w_1)$ has least rank for o, w_1 .

In the sequel R denotes a non empty poset and z denotes a non empty set.

Let us consider R, z . The functor $\text{ConstOSSet}(R, z)$ yielding a many sorted set indexed by the carrier of R is defined by:

- (Def. 16) $\text{ConstOSSet}(R, z) = (\text{the carrier of } R) \mapsto z$.

The following proposition is true

- (15) $\text{ConstOSSet}(R, z)$ is non-empty and for all elements s_1, s_2 of R such that $s_1 \leq s_2$ holds $(\text{ConstOSSet}(R, z))(s_1) \subseteq (\text{ConstOSSet}(R, z))(s_2)$.

Let C be a 1-sorted structure.

- (Def. 17) A many sorted set indexed by the carrier of C is said to be a many sorted set indexed by C .

Let us consider R, z . Then $\text{ConstOSSet}(R, z)$ is a many sorted set indexed by R .

Let us consider R and let M be a many sorted set indexed by R . We say that M is order-sorted if and only if:

- (Def. 18) For all elements s_1, s_2 of R such that $s_1 \leq s_2$ holds $M(s_1) \subseteq M(s_2)$.

Next we state the proposition

- (16) $\text{ConstOSSet}(R, z)$ is order-sorted.

Let us consider R . Observe that there exists a many sorted set indexed by R which is order-sorted.

Let us consider R, z . Then $\text{ConstOSSet}(R, z)$ is an order-sorted many sorted set indexed by R .

Let R be a non empty poset. An order sorted set of R is an order-sorted many sorted set indexed by R .

Let R be a non empty poset. Observe that there exists an order sorted set of R which is non-empty.

We adopt the following convention: s_1, s_2 denote sort symbols of S , o, o_1, o_2, o_3 denote operation symbols of S , and w_1, w_2 denote elements of $(\text{the carrier of } S)^*$.

Let us consider S and let M be an algebra over S . We say that M is order-sorted if and only if:

- (Def. 19) For all s_1, s_2 such that $s_1 \leq s_2$ holds $(\text{the sorts of } M)(s_1) \subseteq (\text{the sorts of } M)(s_2)$.

The following proposition is true

- (17) For every algebra M over S holds M is order-sorted iff the sorts of M are an order sorted set of S .

In the sequel C_1 denotes a many sorted function from $(\text{ConstOSSet}(S, z))^\#$ · the arity of S into $\text{ConstOSSet}(S, z)$ · the result sort of S .

Let us consider S, z, C_1 . The functor $\text{ConstOSA}(S, z, C_1)$ yielding a strict non-empty algebra over S is defined by:

(Def. 20) The sorts of $\text{ConstOSA}(S, z, C_1) = \text{ConstOSSet}(S, z)$ and the characteristics of $\text{ConstOSA}(S, z, C_1) = C_1$.

One can prove the following proposition

(18) $\text{ConstOSA}(S, z, C_1)$ is order-sorted.

Let us consider S . One can check that there exists an algebra over S which is strict, non-empty, and order-sorted.

Let us consider S, z, C_1 . One can verify that $\text{ConstOSA}(S, z, C_1)$ is order-sorted.

Let us consider S . An order sorted algebra of S is an order-sorted algebra over S .

Next we state the proposition

(19) For every discrete order sorted signature S holds every algebra over S is order-sorted.

Let S be a discrete order sorted signature. Observe that every algebra over S is order-sorted.

In the sequel A denotes an order sorted algebra of S .

We now state the proposition

(20) If $w_1 \leq w_2$, then $(\text{the sorts of } A)^\#(w_1) \subseteq (\text{the sorts of } A)^\#(w_2)$.

In the sequel M is an algebra over S_0 .

Let us consider S_0, M . The functor $\text{OSAlg } M$ yielding a strict order sorted algebra of $\text{OSSign } S_0$ is defined as follows:

(Def. 21) The sorts of $\text{OSAlg } M = \text{the sorts of } M$ and the characteristics of $\text{OSAlg } M = \text{the characteristics of } M$.

In the sequel A denotes an order sorted algebra of S .

We now state the proposition

(21) For all elements w_1, w_2, w_3 of $(\text{the carrier of } S)^*$ such that $w_1 \leq w_2$ and $w_2 \leq w_3$ holds $w_1 \leq w_3$.

Let us consider S, o_1, o_2 . The predicate $o_1 \leq o_2$ is defined as follows:

(Def. 22) $o_1 \cong o_2$ and $\text{Arity}(o_1) \leq \text{Arity}(o_2)$ and the result sort of $o_1 \leq$ the result sort of o_2 .

Let us note that the predicate $o_1 \leq o_2$ is reflexive.

We now state several propositions:

(22) If $o_1 \leq o_2$ and $o_2 \leq o_1$, then $o_1 = o_2$.

(23) If $o_1 \leq o_2$ and $o_2 \leq o_3$, then $o_1 \leq o_3$.

(24) If the result sort of $o_1 \leq$ the result sort of o_2 , then $\text{Result}(o_1, A) \subseteq \text{Result}(o_2, A)$.

(25) If $\text{Arity}(o_1) \leq \text{Arity}(o_2)$, then $\text{Args}(o_1, A) \subseteq \text{Args}(o_2, A)$.

- (26) If $o_1 \leq o_2$, then $\text{Args}(o_1, A) \subseteq \text{Args}(o_2, A)$ and $\text{Result}(o_1, A) \subseteq \text{Result}(o_2, A)$.

Let us consider S, A . We say that A is monotone if and only if:

- (Def. 23) For all o_1, o_2 such that $o_1 \leq o_2$ holds $\text{Den}(o_2, A) \upharpoonright \text{Args}(o_1, A) = \text{Den}(o_1, A)$.

We now state two propositions:

- (27) Let A be a non-empty order sorted algebra of S . Then A is monotone if and only if for all o_1, o_2 such that $o_1 \leq o_2$ holds $\text{Den}(o_1, A) \subseteq \text{Den}(o_2, A)$.
- (28) If S is discrete and op-discrete, then A is monotone.

Let us consider S, z and let z_1 be an element of z . The functor $\text{TrivialOSA}(S, z, z_1)$ yielding a strict order sorted algebra of S is defined by:

- (Def. 24) The sorts of $\text{TrivialOSA}(S, z, z_1) = \text{ConstOSSet}(S, z)$ and for every o holds $\text{Den}(o, \text{TrivialOSA}(S, z, z_1)) = \text{Args}(o, \text{TrivialOSA}(S, z, z_1)) \mapsto z_1$.

Next we state the proposition

- (29) For every element z_1 of z holds $\text{TrivialOSA}(S, z, z_1)$ is non-empty and $\text{TrivialOSA}(S, z, z_1)$ is monotone.

Let us consider S . Note that there exists an order sorted algebra of S which is monotone, strict, and non-empty.

Let us consider S, z and let z_1 be an element of z . One can check that $\text{TrivialOSA}(S, z, z_1)$ is monotone and non-empty.

In the sequel o_5, o_6 are operation symbols of S .

Let us consider S . The functor $\text{OperNames } S$ yields a non empty family of subsets of the operation symbols of S and is defined as follows:

- (Def. 25) $\text{OperNames } S = \text{Classes}$ (the overloading of S).

Let us consider S . One can check that every element of $\text{OperNames } S$ is non empty.

Let us consider S . An OperName of S is an element of $\text{OperNames } S$.

Let us consider S, o_5 . The functor $\text{Name } o_5$ yields an OperName of S and is defined by:

- (Def. 26) $\text{Name } o_5 = [o_5]_{\text{the overloading of } S}$.

Next we state three propositions:

- (30) $o_5 \cong o_6$ iff $o_6 \in [o_5]_{\text{the overloading of } S}$.
- (31) $o_5 \cong o_6$ iff $\text{Name } o_5 = \text{Name } o_6$.
- (32) For every set X holds X is an OperName of S iff there exists o_5 such that $X = \text{Name } o_5$.

Let us consider S and let o be an OperName of S . We see that the element of o is an operation symbol of S .

Next we state two propositions:

- (33) For every OperName o_8 of S and for every operation symbol o_7 of S holds o_7 is an element of o_8 iff $\text{Name } o_7 = o_8$.
- (34) Let S_2 be a regular monotone order sorted signature, o_5, o_6 be operation symbols of S_2 , and w be an element of (the carrier of S_2)^{*}. If $o_5 \cong o_6$ and $\text{len Arity}(o_5) = \text{len Arity}(o_6)$ and $w \leq \text{Arity}(o_5)$ and $w \leq \text{Arity}(o_6)$, then $\text{LBound}(o_5, w) = \text{LBound}(o_6, w)$.

Let S_2 be a regular monotone order sorted signature, let o_8 be an OperName of S_2 , and let w be an element of (the carrier of S_2)^{*}. Let us assume that there exists an element o_7 of o_8 such that $w \leq \text{Arity}(o_7)$. The functor $\text{LBound}(o_8, w)$ yields an element of o_8 and is defined as follows:

- (Def. 27) For every element o_7 of o_8 such that $w \leq \text{Arity}(o_7)$ holds $\text{LBound}(o_8, w) = \text{LBound}(o_7, w)$.

Next we state the proposition

- (35) Let S be a regular monotone order sorted signature, o be an operation symbol of S , and w_1 be an element of (the carrier of S)^{*}. If $w_1 \leq \text{Arity}(o)$, then $\text{LBound}(o, w_1) \leq o$.

ACKNOWLEDGMENTS

Thanks to Joseph Goguen, for providing me with his articles on osas, and Andrzej Trybulec, for suggesting and funding this work in Białystok.

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(3):589–593, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Adam Grabowski. On the category of posets. *Formalized Mathematics*, 5(4):501–505, 1996.
- [7] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [8] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Formalized Mathematics*, 1(3):441–444, 1990.
- [9] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [10] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [11] Andrzej Trybulec. Many sorted algebras. *Formalized Mathematics*, 5(1):37–42, 1996.
- [12] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(2):313–319, 1990.
- [13] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [14] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

- [15] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received September 19, 2002
