

Order Sorted Quotient Algebra¹

Josef Urban
Charles University
Praha

MML Identifier: OSALG_4.

The terminology and notation used in this paper are introduced in the following papers: [7], [16], [21], [24], [4], [25], [6], [15], [9], [19], [8], [14], [5], [3], [1], [20], [17], [2], [12], [18], [10], [13], [23], [22], and [11].

1. PRELIMINARIES

Let R be a non empty poset. One can verify that there exists an order sorted set of R which is binary relation yielding.

Let R be a non empty poset, let A, B be many sorted sets indexed by the carrier of R , and let I_1 be a many sorted relation between A and B . We say that I_1 is os-compatible if and only if the condition (Def. 1) is satisfied.

(Def. 1) Let s_1, s_2 be elements of the carrier of R . Suppose $s_1 \leq s_2$. Let x, y be sets. If $x \in A(s_1)$ and $y \in B(s_1)$, then $\langle x, y \rangle \in I_1(s_1)$ iff $\langle x, y \rangle \in I_1(s_2)$.

Let R be a non empty poset and let A, B be many sorted sets indexed by the carrier of R . A many sorted relation between A and B is said to be an order sorted relation of A, B if:

(Def. 2) It is os-compatible.

The following proposition is true

- (1) Let R be a non empty poset, A, B be many sorted sets indexed by the carrier of R , and O_1 be a many sorted relation between A and B . If O_1 is os-compatible, then O_1 is an order sorted set of R .

¹This work was done during author's research visit in Białystok, funded by the CALCULEMUS grant HPRN-CT-2000-00102.

Let R be a non empty poset and let A be a many sorted set indexed by the carrier of R . An order sorted relation of A is an order sorted relation of A, A .

Let S be an order sorted signature and let U_1 be an order sorted algebra of S . A many sorted relation indexed by U_1 is said to be an order sorted relation of U_1 if:

(Def. 3) It is os-compatible.

Let S be an order sorted signature and let U_1 be an order sorted algebra of S . One can check that there exists an order sorted relation of U_1 which is equivalence.

Let S be an order sorted signature and let U_1 be a non-empty order sorted algebra of S . Note that there exists an equivalence order sorted relation of U_1 which is MSCongruence-like.

Let S be an order sorted signature and let U_1 be a non-empty order sorted algebra of S . An order sorted congruence of U_1 is a MSCongruence-like equivalence order sorted relation of U_1 .

Let R be a non empty poset. The functor $\text{PathRel } R$ yields an equivalence relation of the carrier of R and is defined by the condition (Def. 4).

(Def. 4) Let x, y be sets. Then $\langle x, y \rangle \in \text{PathRel } R$ if and only if the following conditions are satisfied:

- (i) $x \in$ the carrier of R ,
- (ii) $y \in$ the carrier of R , and
- (iii) there exists a finite sequence p of elements of the carrier of R such that $1 < \text{len } p$ and $p(1) = x$ and $p(\text{len } p) = y$ and for every natural number n such that $2 \leq n$ and $n \leq \text{len } p$ holds $\langle p(n), p(n-1) \rangle \in$ the internal relation of R or $\langle p(n-1), p(n) \rangle \in$ the internal relation of R .

One can prove the following proposition

- (2) For every non empty poset R and for all elements s_1, s_2 of the carrier of R such that $s_1 \leq s_2$ holds $\langle s_1, s_2 \rangle \in \text{PathRel } R$.

Let R be a non empty poset and let s_1, s_2 be elements of the carrier of R . The predicate $s_1 \cong s_2$ is defined as follows:

(Def. 5) $\langle s_1, s_2 \rangle \in \text{PathRel } R$.

Let us notice that the predicate $s_1 \cong s_2$ is reflexive and symmetric.

One can prove the following proposition

- (3) For every non empty poset R and for all elements s_1, s_2, s_3 of the carrier of R such that $s_1 \cong s_2$ and $s_2 \cong s_3$ holds $s_1 \cong s_3$.

Let R be a non empty poset. The functor $\text{Components } R$ yields a non empty family of subsets of the carrier of R and is defined by:

(Def. 6) $\text{Components } R = \text{Classes PathRel } R$.

Let R be a non empty poset. Note that every element of $\text{Components } R$ is non empty.

Let R be a non empty poset. A subset of R is called a component of R if:

(Def. 7) $It \in \text{Components } R$.

Let R be a non empty poset and let s_1 be an element of the carrier of R .

The functor $\cdot_{\text{CSp}} s_1$ yielding a component of R is defined by:

(Def. 8) $\cdot_{\text{CSp}} s_1 = [s_1]_{\text{PathRel } R}$.

The following two propositions are true:

- (4) For every non empty poset R and for every element s_1 of the carrier of R holds $s_1 \in \cdot_{\text{CSp}} s_1$.
- (5) For every non empty poset R and for all elements s_1, s_2 of the carrier of R such that $s_1 \leq s_2$ holds $\cdot_{\text{CSp}} s_1 = \cdot_{\text{CSp}} s_2$.

Let R be a non empty poset, let A be a many sorted set indexed by the carrier of R , and let C be a component of R . A -carrier of C is defined as follows:

(Def. 9) $A\text{-carrier of } C = \bigcup\{A(s); s \text{ ranges over elements of the carrier of } R; s \in C\}$.

We now state the proposition

- (6) Let R be a non empty poset, A be a many sorted set indexed by the carrier of R , s be an element of the carrier of R , and x be a set. If $x \in A(s)$, then $x \in A\text{-carrier of } \cdot_{\text{CSp}} s$.

Let R be a non empty poset. We say that R is locally directed if and only if:

(Def. 10) Every component of R is directed.

The following three propositions are true:

- (7) For every discrete non empty poset R and for all elements x, y of the carrier of R such that $\langle x, y \rangle \in \text{PathRel } R$ holds $x = y$.
- (8) Let R be a discrete non empty poset and C be a component of R . Then there exists an element x of the carrier of R such that $C = \{x\}$.
- (9) Every discrete non empty poset is locally directed.

Let us observe that there exists a non empty poset which is locally directed.

One can verify that there exists an order sorted signature which is locally directed.

Let us observe that every non empty poset which is discrete is also locally directed.

Let S be a locally directed non empty poset. Note that every component of S is directed.

One can prove the following proposition

- (10) \emptyset is an equivalence relation of \emptyset .

Let S be a locally directed order sorted signature, let A be an order sorted algebra of S , let E be an equivalence order sorted relation of A , and let C be a

component of S . The functor $\text{CompClass}(E, C)$ yielding an equivalence relation of (the sorts of A)-carrier of C is defined as follows:

- (Def. 11) For all sets x, y holds $\langle x, y \rangle \in \text{CompClass}(E, C)$ iff there exists an element s_1 of the carrier of S such that $s_1 \in C$ and $\langle x, y \rangle \in E(s_1)$.

Let S be a locally directed order sorted signature, let A be an order sorted algebra of S , let E be an equivalence order sorted relation of A , and let s_1 be an element of the carrier of S . The functor $\text{OSClass}(E, s_1)$ yielding a subset of Classes $\text{CompClass}(E, \cdot_{\text{CSP}} s_1)$ is defined by:

- (Def. 12) For every set z holds $z \in \text{OSClass}(E, s_1)$ iff there exists a set x such that $x \in (\text{the sorts of } A)(s_1)$ and $z = [x]_{\text{CompClass}(E, \cdot_{\text{CSP}} s_1)}$.

Let S be a locally directed order sorted signature, let A be a non-empty order sorted algebra of S , let E be an equivalence order sorted relation of A , and let s_1 be an element of the carrier of S . One can verify that $\text{OSClass}(E, s_1)$ is non empty.

The following proposition is true

- (11) Let S be a locally directed order sorted signature, A be an order sorted algebra of S , E be an equivalence order sorted relation of A , and s_1, s_2 be elements of the carrier of S . If $s_1 \leq s_2$, then $\text{OSClass}(E, s_1) \subseteq \text{OSClass}(E, s_2)$.

Let S be a locally directed order sorted signature, let A be an order sorted algebra of S , and let E be an equivalence order sorted relation of A . The functor $\text{OSClass } E$ yields an order sorted set of S and is defined as follows:

- (Def. 13) For every element s_1 of the carrier of S holds $(\text{OSClass } E)(s_1) = \text{OSClass}(E, s_1)$.

Let S be a locally directed order sorted signature, let A be a non-empty order sorted algebra of S , and let E be an equivalence order sorted relation of A . One can check that $\text{OSClass } E$ is non-empty.

Let S be a locally directed order sorted signature, let U_1 be a non-empty order sorted algebra of S , let E be an equivalence order sorted relation of U_1 , let s be an element of the carrier of S , and let x be an element of $(\text{the sorts of } U_1)(s)$. The functor $\text{OSClass}(E, x)$ yields an element of $\text{OSClass}(E, s)$ and is defined by:

- (Def. 14) $\text{OSClass}(E, x) = [x]_{\text{CompClass}(E, \cdot_{\text{CSP}} s)}$.

One can prove the following three propositions:

- (12) Let R be a locally directed non empty poset and x, y be elements of the carrier of R . Given an element z of the carrier of R such that $z \leq x$ and $z \leq y$. Then there exists an element u of R such that $x \leq u$ and $y \leq u$.
- (13) Let S be a locally directed order sorted signature, U_1 be a non-empty order sorted algebra of S , E be an equivalence order sorted relation of U_1 ,

- s be an element of the carrier of S , and x, y be elements of (the sorts of U_1)(s). Then $\text{OSClass}(E, x) = \text{OSClass}(E, y)$ if and only if $\langle x, y \rangle \in E(s)$.
- (14) Let S be a locally directed order sorted signature, U_1 be a non-empty order sorted algebra of S , E be an equivalence order sorted relation of U_1 , s_1, s_2 be elements of the carrier of S , and x be an element of (the sorts of U_1)(s_1). Suppose $s_1 \leq s_2$. Let y be an element of (the sorts of U_1)(s_2). If $y = x$, then $\text{OSClass}(E, x) = \text{OSClass}(E, y)$.

2. ORDER SORTED QUOTIENT ALGEBRA

In the sequel S denotes a locally directed order sorted signature and o denotes an element of the operation symbols of S .

Let us consider S, o , let A be a non-empty order sorted algebra of S , let R be an order sorted congruence of A , and let x be an element of $\text{Args}(o, A)$. The functor $Rosx$ yields an element of $\prod(\text{OSClass } R \cdot \text{Arity}(o))$ and is defined by the condition (Def. 15).

- (Def. 15) Let n be a natural number. Suppose $n \in \text{dom Arity}(o)$. Then there exists an element y of (the sorts of A)($\text{Arity}(o)_n$) such that $y = x(n)$ and $(Rosx)(n) = \text{OSClass}(R, y)$.

Let us consider S, o , let A be a non-empty order sorted algebra of S , and let R be an order sorted congruence of A . The functor $\text{OSQuotRes}(R, o)$ yielding a function from ((the sorts of A) · (the result sort of S))(o) into $(\text{OSClass } R \cdot \text{the result sort of } S)(o)$ is defined as follows:

- (Def. 16) For every element x of (the sorts of A)(the result sort of o) holds $(\text{OSQuotRes}(R, o))(x) = \text{OSClass}(R, x)$.

The functor $\text{OSQuotArgs}(R, o)$ yielding a function from ((the sorts of A)[#] · the arity of S)(o) into $(\text{OSClass } R$)[#] · the arity of S)(o) is defined by:

- (Def. 17) For every element x of $\text{Args}(o, A)$ holds $(\text{OSQuotArgs}(R, o))(x) = Rosx$.

Let us consider S , let A be a non-empty order sorted algebra of S , and let R be an order sorted congruence of A . The functor $\text{OSQuotRes } R$ yields a many sorted function from (the sorts of A) · (the result sort of S) into $\text{OSClass } R \cdot \text{the result sort of } S$ and is defined by:

- (Def. 18) For every operation symbol o of S holds $(\text{OSQuotRes } R)(o) = \text{OSQuotRes}(R, o)$.

The functor $\text{OSQuotArgs } R$ yields a many sorted function from (the sorts of A)[#] · the arity of S into $(\text{OSClass } R$)[#] · the arity of S and is defined as follows:

- (Def. 19) For every operation symbol o of S holds $(\text{OSQuotArgs } R)(o) = \text{OSQuotArgs}(R, o)$.

One can prove the following proposition

- (15) Let A be a non-empty order sorted algebra of S , R be an order sorted congruence of A , and x be a set. Suppose $x \in ((\text{OSClass } R)^\# \cdot \text{the arity of } S)(o)$. Then there exists an element a of $\text{Args}(o, A)$ such that $x = \text{Rosa}$.

Let us consider S , o , let A be a non-empty order sorted algebra of S , and let R be an order sorted congruence of A . The functor $\text{OSQuotCharact}(R, o)$ yielding a function from $((\text{OSClass } R)^\# \cdot \text{the arity of } S)(o)$ into $(\text{OSClass } R \cdot \text{the result sort of } S)(o)$ is defined as follows:

- (Def. 20) For every element a of $\text{Args}(o, A)$ such that $\text{Rosa} \in ((\text{OSClass } R)^\# \cdot \text{the arity of } S)(o)$ holds $(\text{OSQuotCharact}(R, o))(\text{Rosa}) = (\text{OSQuotRes}(R, o) \cdot \text{Den}(o, A))(a)$.

Let us consider S , let A be a non-empty order sorted algebra of S , and let R be an order sorted congruence of A . The functor $\text{OSQuotCharact } R$ yielding a many sorted function from $(\text{OSClass } R)^\# \cdot \text{the arity of } S$ into $\text{OSClass } R \cdot \text{the result sort of } S$ is defined as follows:

- (Def. 21) For every operation symbol o of S holds $(\text{OSQuotCharact } R)(o) = \text{OSQuotCharact}(R, o)$.

Let us consider S , let U_1 be a non-empty order sorted algebra of S , and let R be an order sorted congruence of U_1 . The functor $\text{QuotOSAlg}(U_1, R)$ yields an order sorted algebra of S and is defined by:

- (Def. 22) $\text{QuotOSAlg}(U_1, R) = \langle \text{OSClass } R, \text{OSQuotCharact } R \rangle$.

Let us consider S , let U_1 be a non-empty order sorted algebra of S , and let R be an order sorted congruence of U_1 . One can check that $\text{QuotOSAlg}(U_1, R)$ is strict and non-empty.

Let us consider S , let U_1 be a non-empty order sorted algebra of S , let R be an order sorted congruence of U_1 , and let s be an element of the carrier of S . The functor $\text{OSNatHom}(U_1, R, s)$ yielding a function from $(\text{the sorts of } U_1)(s)$ into $\text{OSClass}(R, s)$ is defined by:

- (Def. 23) For every element x of $(\text{the sorts of } U_1)(s)$ holds $(\text{OSNatHom}(U_1, R, s))(x) = \text{OSClass}(R, x)$.

Let us consider S , let U_1 be a non-empty order sorted algebra of S , and let R be an order sorted congruence of U_1 . The functor $\text{OSNatHom}(U_1, R)$ yielding a many sorted function from U_1 into $\text{QuotOSAlg}(U_1, R)$ is defined as follows:

- (Def. 24) For every element s of the carrier of S holds $(\text{OSNatHom}(U_1, R))(s) = \text{OSNatHom}(U_1, R, s)$.

Next we state two propositions:

- (16) Let U_1 be a non-empty order sorted algebra of S and R be an order sorted congruence of U_1 . Then $\text{OSNatHom}(U_1, R)$ is an epimorphism of U_1 onto $\text{QuotOSAlg}(U_1, R)$ and $\text{OSNatHom}(U_1, R)$ is order-sorted.
- (17) Let U_1, U_2 be non-empty order sorted algebras of S and F be a many sorted function from U_1 into U_2 . Suppose F is a homomorphism of U_1 into

U_2 and order-sorted. Then $\text{Congruence}(F)$ is an order sorted congruence of U_1 .

Let us consider S , let U_1, U_2 be non-empty order sorted algebras of S , and let F be a many sorted function from U_1 into U_2 . Let us assume that F is a homomorphism of U_1 into U_2 and order-sorted. The functor $\text{OSCng } F$ yielding an order sorted congruence of U_1 is defined as follows:

(Def. 25) $\text{OSCng } F = \text{Congruence}(F)$.

Let us consider S , let U_1, U_2 be non-empty order sorted algebras of S , let F be a many sorted function from U_1 into U_2 , and let s be an element of the carrier of S . Let us assume that F is a homomorphism of U_1 into U_2 and order-sorted. The functor $\text{OSHomQuot}(F, s)$ yields a function from (the sorts of $\text{QuotOSAlg}(U_1, \text{OSCng } F)$)(s) into (the sorts of U_2)(s) and is defined as follows:

(Def. 26) For every element x of (the sorts of U_1)(s) holds

$$(\text{OSHomQuot}(F, s))(\text{OSClass}(\text{OSCng } F, x)) = F(s)(x).$$

Let us consider S , let U_1, U_2 be non-empty order sorted algebras of S , and let F be a many sorted function from U_1 into U_2 . The functor $\text{OSHomQuot } F$ yields a many sorted function from $\text{QuotOSAlg}(U_1, \text{OSCng } F)$ into U_2 and is defined by:

(Def. 27) For every element s of the carrier of S holds $(\text{OSHomQuot } F)(s) = \text{OSHomQuot}(F, s)$.

The following three propositions are true:

- (18) Let U_1, U_2 be non-empty order sorted algebras of S and F be a many sorted function from U_1 into U_2 . Suppose F is a homomorphism of U_1 into U_2 and order-sorted. Then $\text{OSHomQuot } F$ is a monomorphism of $\text{QuotOSAlg}(U_1, \text{OSCng } F)$ into U_2 and $\text{OSHomQuot } F$ is order-sorted.
- (19) Let U_1, U_2 be non-empty order sorted algebras of S and F be a many sorted function from U_1 into U_2 . Suppose F is an epimorphism of U_1 onto U_2 and order-sorted. Then $\text{OSHomQuot } F$ is an isomorphism of $\text{QuotOSAlg}(U_1, \text{OSCng } F)$ and U_2 .
- (20) Let U_1, U_2 be non-empty order sorted algebras of S and F be a many sorted function from U_1 into U_2 . Suppose F is an epimorphism of U_1 onto U_2 and order-sorted. Then $\text{QuotOSAlg}(U_1, \text{OSCng } F)$ and U_2 are isomorphic.

Let S be an order sorted signature, let U_1 be a non-empty order sorted algebra of S , and let R be an equivalence order sorted relation of U_1 . We say that R is monotone if and only if the condition (Def. 28) is satisfied.

(Def. 28) Let o_1, o_2 be operation symbols of S . Suppose $o_1 \leq o_2$. Let x_1 be an element of $\text{Args}(o_1, U_1)$ and x_2 be an element of $\text{Args}(o_2, U_1)$. Suppose that for every natural number y such that $y \in \text{dom } x_1$ holds $\langle x_1(y), x_2(y) \rangle \in$

$R(\text{Arity}(o_2)_y)$. Then $\langle (\text{Den}(o_1, U_1))(x_1), (\text{Den}(o_2, U_1))(x_2) \rangle \in R$ (the result sort of o_2).

One can prove the following two propositions:

- (21) Let S be an order sorted signature and U_1 be a non-empty order sorted algebra of S . Then $\llbracket \text{the sorts of } U_1, \text{ the sorts of } U_1 \rrbracket$ is an order sorted congruence of U_1 .
- (22) Let S be an order sorted signature, U_1 be a non-empty order sorted algebra of S , and R be an order sorted congruence of U_1 . If $R = \llbracket \text{the sorts of } U_1, \text{ the sorts of } U_1 \rrbracket$, then R is monotone.

Let S be an order sorted signature and let U_1 be a non-empty order sorted algebra of S . One can verify that there exists an order sorted congruence of U_1 which is monotone.

Let S be an order sorted signature and let U_1 be a non-empty order sorted algebra of S . Note that there exists an equivalence order sorted relation of U_1 which is monotone.

The following proposition is true

- (23) Let S be an order sorted signature and U_1 be a non-empty order sorted algebra of S . Then every monotone equivalence order sorted relation of U_1 is MSCongruence-like.

Let S be an order sorted signature and let U_1 be a non-empty order sorted algebra of S . Observe that every equivalence order sorted relation of U_1 which is monotone is also MSCongruence-like.

We now state the proposition

- (24) Let S be an order sorted signature and U_1 be a monotone non-empty order sorted algebra of S . Then every order sorted congruence of U_1 is monotone.

Let S be an order sorted signature and let U_1 be a monotone non-empty order sorted algebra of S . Observe that every order sorted congruence of U_1 is monotone.

Let us consider S , let U_1 be a non-empty order sorted algebra of S , and let R be a monotone order sorted congruence of U_1 . Note that $\text{QuotOSAlg}(U_1, R)$ is monotone.

We now state two propositions:

- (25) Let given S , U_1 be a non-empty order sorted algebra of S , and R be a monotone order sorted congruence of U_1 . Then $\text{QuotOSAlg}(U_1, R)$ is a monotone order sorted algebra of S .
- (26) Let U_1 be a non-empty order sorted algebra of S , U_2 be a monotone non-empty order sorted algebra of S , and F be a many sorted function from U_1 into U_2 . Suppose F is a homomorphism of U_1 into U_2 and order-sorted. Then $\text{OSCng } F$ is monotone.

Let us consider S , let U_1, U_2 be non-empty order sorted algebras of S , let F be a many sorted function from U_1 into U_2 , let R be an order sorted congruence of U_1 , and let s be an element of the carrier of S . Let us assume that F is a homomorphism of U_1 into U_2 and order-sorted and $R \subseteq \text{OSCng } F$. The functor $\text{OSHomQuot}(F, R, s)$ yields a function from (the sorts of $\text{QuotOSAlg}(U_1, R)$)(s) into (the sorts of U_2)(s) and is defined as follows:

(Def. 29) For every element x of (the sorts of U_1)(s) holds
 $(\text{OSHomQuot}(F, R, s))(\text{OSClass}(R, x)) = F(s)(x)$.

Let us consider S , let U_1, U_2 be non-empty order sorted algebras of S , let F be a many sorted function from U_1 into U_2 , and let R be an order sorted congruence of U_1 . The functor $\text{OSHomQuot}(F, R)$ yields a many sorted function from $\text{QuotOSAlg}(U_1, R)$ into U_2 and is defined as follows:

(Def. 30) For every element s of the carrier of S holds $(\text{OSHomQuot}(F, R))(s) = \text{OSHomQuot}(F, R, s)$.

Next we state the proposition

(27) Let U_1, U_2 be non-empty order sorted algebras of S , F be a many sorted function from U_1 into U_2 , and R be an order sorted congruence of U_1 . Suppose F is a homomorphism of U_1 into U_2 and order-sorted and $R \subseteq \text{OSCng } F$. Then $\text{OSHomQuot}(F, R)$ is a homomorphism of $\text{QuotOSAlg}(U_1, R)$ into U_2 and $\text{OSHomQuot}(F, R)$ is order-sorted.

ACKNOWLEDGMENTS

Thanks to Joseph Goguen, for providing me with his articles on osas, and Andrzej Trybulec, for suggesting and funding this work in Białystok.

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(3):589–593, 1990.
- [2] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Formalized Mathematics*, 6(1):93–107, 1997.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [7] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [8] Adam Grabowski. On the category of posets. *Formalized Mathematics*, 5(4):501–505, 1996.
- [9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [10] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Formalized Mathematics*, 5(1):61–65, 1996.
- [11] Małgorzata Korolkiewicz. Many sorted quotient algebra. *Formalized Mathematics*, 5(1):79–84, 1996.

- [12] Beata Madras. Product of family of universal algebras. *Formalized Mathematics*, 4(1):103–108, 1993.
- [13] Beata Madras. Products of many sorted algebras. *Formalized Mathematics*, 5(1):55–60, 1996.
- [14] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [15] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Formalized Mathematics*, 1(3):441–444, 1990.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [17] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [18] Andrzej Trybulec. Many sorted algebras. *Formalized Mathematics*, 5(1):37–42, 1996.
- [19] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(2):313–319, 1990.
- [20] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [21] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [22] Josef Urban. Homomorphisms of order sorted algebras. *Formalized Mathematics*, 10(3):197–200, 2002.
- [23] Josef Urban. Order sorted algebras. *Formalized Mathematics*, 10(3):179–188, 2002.
- [24] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [25] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received September 19, 2002
