

# Improvement of Radix- $2^k$ Signed-Digit Number for High Speed Circuit

Masaaki Niimura  
Shinshu University  
Nagano

Yasushi Fuwa  
Shinshu University  
Nagano

**Summary.** In this article, a new radix- $2^k$  signed-digit number (Radix- $2^k$  sub signed-digit number) is defined and its properties for hardware realization are discussed.

Until now, high speed calculation method with Radix- $2^k$  signed-digit numbers is proposed, but this method used “Compares With 2” to calculate carry. “Compares with 2” is a very simple method, but it needs very complicated hardware especially when the value of  $k$  becomes large. In this article, we propose a subset of Radix- $2^k$  signed-digit, named Radix- $2^k$  sub signed-digit numbers. Radix- $2^k$  sub signed-digit was designed so that the carry calculation use “bit compare” to hardware-realization simplifies more.

In the first section of this article, we defined the concept of Radix- $2^k$  sub signed-digit numbers and proved some of their properties. In the second section, we defined the new carry calculation method in consideration of hardware-realization, and proved some of their properties. In the third section, we provide some functions for generating Radix- $2^k$  sub signed-digit numbers from Radix- $2^k$  signed-digit numbers. In the last section, we defined some functions for generation natural numbers from Radix- $2^k$  sub signed-digit, and we clarified its correctness.

MML Identifier: RADIX\_3.

The articles [11], [14], [8], [12], [1], [4], [3], [13], [10], [7], [2], [9], [5], and [6] provide the notation and terminology for this paper.

## 1. DEFINITION FOR RADIX- $2^k$ SUB SIGNED-DIGIT NUMBER

We adopt the following convention:  $i$ ,  $n$ ,  $m$ ,  $k$ ,  $x$  are natural numbers and  $i_1$ ,  $i_2$  are integers.

Next we state the proposition

$$(1) \quad ((\text{Radix } k)_{\mathbb{N}}^n) \cdot \text{Radix } k = (\text{Radix } k)_{\mathbb{N}}^{n+1}.$$

Let us consider  $k$ . The functor  $k\text{-SD\_Sub\_S}$  is defined as follows:

$$(\text{Def. 1}) \quad k\text{-SD\_Sub\_S} = \{e; e \text{ ranges over elements of } \mathbb{Z}: e \geq -\text{Radix}(k-1) \wedge e \leq \text{Radix}(k-1) - 1\}.$$

Let us consider  $k$ . The functor  $k\text{-SD\_Sub}$  is defined by:

$$(\text{Def. 2}) \quad k\text{-SD\_Sub} = \{e; e \text{ ranges over elements of } \mathbb{Z}: e \geq -\text{Radix}(k-1) - 1 \wedge e \leq \text{Radix}(k-1)\}.$$

The following propositions are true:

- (2) If  $i_1 \in k\text{-SD\_Sub}$ , then  $-\text{Radix}(k-1) - 1 \leq i_1$  and  $i_1 \leq \text{Radix}(k-1)$ .
- (3) For every natural number  $k$  holds  $k\text{-SD\_Sub\_S} \subseteq k\text{-SD\_Sub}$ .
- (4)  $k\text{-SD\_Sub\_S} \subseteq (k+1)\text{-SD\_Sub\_S}$ .
- (5) For every natural number  $k$  such that  $2 \leq k$  holds  $k\text{-SD\_Sub} \subseteq k\text{-SD}$ .
- (6)  $0 \in 0\text{-SD\_Sub\_S}$ .
- (7)  $0 \in k\text{-SD\_Sub\_S}$ .
- (8)  $0 \in k\text{-SD\_Sub}$ .
- (9) For every set  $e$  such that  $e \in k\text{-SD\_Sub}$  holds  $e$  is an integer.
- (10)  $k\text{-SD\_Sub} \subseteq \mathbb{Z}$ .
- (11)  $k\text{-SD\_Sub\_S} \subseteq \mathbb{Z}$ .

Let us consider  $k$ . One can verify that  $k\text{-SD\_Sub\_S}$  is non empty.

Let us consider  $k$ . Note that  $k\text{-SD\_Sub}$  is non empty.

Let us consider  $k$ . Then  $k\text{-SD\_Sub\_S}$  is a non empty subset of  $\mathbb{Z}$ .

Let us consider  $k$ . Then  $k\text{-SD\_Sub}$  is a non empty subset of  $\mathbb{Z}$ .

In the sequel  $a$  denotes a  $n$ -tuple of  $k\text{-SD}$  and  $a_1$  denotes a  $n$ -tuple of  $k\text{-SD\_Sub}$ .

One can prove the following proposition

$$(12) \quad \text{If } i \in \text{Seg } n, \text{ then } a_1(i) \text{ is an element of } k\text{-SD\_Sub}.$$

## 2. DEFINITION FOR NEW CARRY CALCULATION METHOD

Let  $x$  be an integer and let  $k$  be a natural number.

The functor  $\text{SDSubAddCarry}(x, k)$  yields an integer and is defined as follows:

$$(\text{Def. 3}) \quad \text{SDSubAddCarry}(x, k) = \begin{cases} 1, & \text{if } \text{Radix}(k-1) \leq x, \\ -1, & \text{if } x < -\text{Radix}(k-1), \\ 0, & \text{otherwise.} \end{cases}$$

Let  $x$  be an integer and let  $k$  be a natural number.

The functor  $\text{SDSubAddData}(x, k)$  yields an integer and is defined as follows:

$$(\text{Def. 4}) \quad \text{SDSubAddData}(x, k) = x - \text{Radix } k \cdot \text{SDSubAddCarry}(x, k).$$

One can prove the following propositions:

- (13) For every integer  $x$  and for every natural number  $k$  such that  $2 \leq k$  holds  $-1 \leq \text{SDSubAddCarry}(x, k)$  and  $\text{SDSubAddCarry}(x, k) \leq 1$ .
- (14) If  $2 \leq k$  and  $i_1 \in k\text{-SD}$ , then  $\text{SDSubAddData}(i_1, k) \geq -\text{Radix}(k-1)$  and  $\text{SDSubAddData}(i_1, k) \leq \text{Radix}(k-1) - 1$ .
- (15) For every integer  $x$  and for every natural number  $k$  such that  $2 \leq k$  holds  $\text{SDSubAddCarry}(x, k) \in k\text{-SD\_Sub\_S}$ .
- (16) If  $2 \leq k$  and  $i_1 \in k\text{-SD}$  and  $i_2 \in k\text{-SD}$ , then  $\text{SDSubAddData}(i_1, k) + \text{SDSubAddCarry}(i_2, k) \in k\text{-SD\_Sub}$ .
- (17) If  $2 \leq k$ , then  $\text{SDSubAddCarry}(0, k) = 0$ .

### 3. DEFINITION FOR TRANSLATION FROM RADIX- $2^k$ SIGNED-DIGIT NUMBER

Let  $i, k, n$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k\text{-SD\_Sub}$ . The functor  $\text{DigA\_SDSub}(x, i)$  yields an integer and is defined as follows:

- (Def. 5)(i)  $\text{DigA\_SDSub}(x, i) = x(i)$  if  $i \in \text{Seg } n$ ,  
(ii)  $\text{DigA\_SDSub}(x, i) = 0$  if  $i = 0$ .

Let  $i, k, n$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k\text{-SD}$ . The functor  $\text{SD2SDSubDigit}(x, i, k)$  yields an integer and is defined by:

$$(\text{Def. 6}) \quad \text{SD2SDSubDigit}(x, i, k) = \begin{cases} \text{(i) } \text{SDSubAddData}(\text{DigA}(x, i), k) + \\ \quad \text{SDSubAddCarry}(\text{DigA}(x, i-1), k), \\ \quad \text{if } i \in \text{Seg } n, \\ \text{(ii) } \text{SDSubAddCarry}(\text{DigA}(x, i-1), k), \\ \quad \text{if } i = n+1, \\ 0, \text{ otherwise.} \end{cases}$$

We now state the proposition

- (18) If  $2 \leq k$  and  $i \in \text{Seg}(n+1)$ , then  $\text{SD2SDSubDigit}(a, i, k)$  is an element of  $k\text{-SD\_Sub}$ .

Let  $i, k, n$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k\text{-SD}$ . Let us assume that  $2 \leq k$  and  $i \in \text{Seg}(n+1)$ . The functor  $\text{SD2SDSubDigitS}(x, i, k)$  yielding an element of  $k\text{-SD\_Sub}$  is defined by:

- (Def. 7)  $\text{SD2SDSubDigitS}(x, i, k) = \text{SD2SDSubDigit}(x, i, k)$ .

Let  $n, k$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k\text{-SD}$ . The functor  $\text{SD2SDSub } x$  yielding a  $n+1$ -tuple of  $k\text{-SD\_Sub}$  is defined by:

- (Def. 8) For every natural number  $i$  such that  $i \in \text{Seg}(n+1)$  holds  $\text{DigA\_SDSub}(\text{SD2SDSub } x, i) = \text{SD2SDSubDigitS}(x, i, k)$ .

Next we state two propositions:

- (19) If  $i \in \text{Seg } n$ , then  $\text{DigA\_SDSub}(a_1, i)$  is an element of  $k\text{-SD\_Sub}$ .
- (20) If  $2 \leq k$  and  $i_1 \in k\text{-SD}$  and  $i_2 \in k\text{-SD\_Sub}$ , then  $\text{SDSubAddData}(i_1 + i_2, k) \in k\text{-SD\_Sub\_S}$ .

4. DEFINITION FOR TRANSLATION FROM RADIX- $2^k$  SUB SIGNED-DIGIT  
NUMBER TO INT

Let  $i, k, n$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k$ -SD\_Sub. The functor  $\text{DigB\_SDSub}(x, i)$  yielding an element of  $\mathbb{Z}$  is defined by:

(Def. 9)  $\text{DigB\_SDSub}(x, i) = \text{DigA\_SDSub}(x, i)$ .

Let  $i, k, n$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k$ -SD\_Sub. The functor  $\text{SDSub2INTDigit}(x, i, k)$  yielding an element of  $\mathbb{Z}$  is defined as follows:

(Def. 10)  $\text{SDSub2INTDigit}(x, i, k) = ((\text{Radix } k)_{\mathbb{N}}^{i-1}) \cdot \text{DigB\_SDSub}(x, i)$ .

Let  $n, k$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k$ -SD\_Sub. The functor  $\text{SDSub2INT } x$  yields a  $n$ -tuple of  $\mathbb{Z}$  and is defined as follows:

(Def. 11) For every natural number  $i$  such that  $i \in \text{Seg } n$  holds  $(\text{SDSub2INT } x)_i = \text{SDSub2INTDigit}(x, i, k)$ .

Let  $n, k$  be natural numbers and let  $x$  be a  $n$ -tuple of  $k$ -SD\_Sub. The functor  $\text{SDSub2IntOut } x$  yields an integer and is defined as follows:

(Def. 12)  $\text{SDSub2IntOut } x = \sum \text{SDSub2INT } x$ .

Next we state two propositions:

(21) For every  $i$  such that  $i \in \text{Seg } n$  holds if  $2 \leq k$ , then

$$\begin{aligned} \text{DigA\_SDSub}(\text{SD2SDSub DecSD}(m, n+1, k), i) = \\ \text{DigA\_SDSub}(\text{SD2SDSub DecSD}(m \bmod (\text{Radix } k)_{\mathbb{N}}^n, n, k), i). \end{aligned}$$

(22) For every  $n$  such that  $n \geq 1$  and for all  $k, x$  such that  $k \geq 2$  and  $x$  is represented by  $n, k$  holds  $x = \text{SDSub2IntOut SD2SDSub DecSD}(x, n, k)$ .

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Joining of decorated trees. *Formalized Mathematics*, 4(1):77–82, 1993.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Marek Chmur. The lattice of natural numbers and the sublattice of it. The set of prime numbers. *Formalized Mathematics*, 2(4):453–459, 1991.
- [6] Yoshinori Fujisawa and Yasushi Fuwa. Definitions of radix- $2^k$  signed-digit number and its adder algorithm. *Formalized Mathematics*, 9(1):71–75, 2001.
- [7] Yoshinori Fujisawa, Yasushi Fuwa, and Hidetaka Shimizu. Euler's Theorem and small Fermat's Theorem. *Formalized Mathematics*, 7(1):123–126, 1998.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [9] Andrzej Kondracki. The Chinese Remainder Theorem. *Formalized Mathematics*, 6(4):573–577, 1997.
- [10] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [12] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.

- [13] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(**3**):575–579, 1990.
- [14] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.

*Received January 3, 2003*

---