

High Speed Adder Algorithm with Radix- 2^k Sub Signed-Digit Number

Masaaki Niimura
Shinshu University
Nagano

Yasushi Fuwa
Shinshu University
Nagano

Summary. In this article, a new adder algorithm using Radix- 2^k sub signed-digit numbers is defined and properties for the hardware-realization is discussed.

Until now, we proposed Radix- 2^k sub signed-digit numbers in consideration of the hardware realization. In this article, we proposed High Speed Adder Algorithm using this Radix- 2^k sub signed-digit numbers. This method has two ways to speed up at hardware-realization. One is 'bit compare' at carry calculation, it is proposed in another article. Other is carry calculation between two numbers. We proposed that n digits Radix- 2^k signed-digit numbers is expressed in $n + 1$ digits Radix- 2^k sub signed-digit numbers, and addition result of two $n + 1$ digits Radix- 2^k sub signed-digit numbers is expressed in $n + 1$ digits. In this way, carry operation between two Radix- 2^k sub signed-digit numbers can be processed at $n + 1$ digit adder circuit and additional circuit to operate carry is not needed.

In the first section of this article, we prepared some useful theorems for operation of Radix- 2^k numbers. In the second section, we proved some properties about carry on Radix- 2^k sub signed-digit numbers. In the last section, we defined the new addition operation using Radix- 2^k sub signed-digit numbers, and we clarified its correctness.

MML Identifier: RADIX.4.

The terminology and notation used here are introduced in the following articles: [11], [13], [12], [1], [4], [3], [10], [7], [2], [8], [5], [6], and [9].

1. PRELIMINARIES

In this paper i, n, m, k, x, y are natural numbers.

The following proposition is true

- (1) For every natural number k such that $2 \leq k$ holds $2 < \text{Radix } k$.

2. CARRY OPERATION AT $n + 1$ DIGITS RADIX- 2^k SUB SIGNED-DIGIT NUMBER

The following propositions are true:

- (2) For all integers x, y and for every natural number k such that $3 \leq k$ holds $\text{SDSubAddCarry}(\text{SDSubAddCarry}(x, k) + \text{SDSubAddCarry}(y, k), k) = 0$.
- (3) If $2 \leq k$, then $\text{DigA_SDSub}(\text{SD2SDSub DecSD}(m, n, k), n + 1) = \text{SDSubAddCarry}(\text{DigA}(\text{DecSD}(m, n, k), n), k)$.
- (4) If $2 \leq k$ and m is represented by $1, k$, then $\text{DigA_SDSub}(\text{SD2SDSub DecSD}(m, 1, k), 1 + 1) = \text{SDSubAddCarry}(m, k)$.
- (5) Let k, x, n be natural numbers. Suppose $n \geq 1$ and $k \geq 3$ and x is represented by $n + 1, k$. Then $\text{DigA_SDSub}(\text{SD2SDSub DecSD}(x \bmod (\text{Radix } k)_{\mathbb{N}}^n, n, k), n + 1) = \text{SDSubAddCarry}(\text{DigA}(\text{DecSD}(x, n, k), n), k)$.
- (6) If $2 \leq k$ and m is represented by $1, k$, then $\text{DigA_SDSub}(\text{SD2SDSub DecSD}(m, 1, k), 1) = m - \text{SDSubAddCarry}(m, k) \cdot \text{Radix } k$.
- (7) Let k, x, n be natural numbers. Suppose $n \geq 1$ and $k \geq 2$ and x is represented by $n + 1, k$. Then $((\text{Radix } k)_{\mathbb{N}}^n) \cdot \text{DigA_SDSub}(\text{SD2SDSub DecSD}(x, n + 1, k), n + 1) = (((\text{Radix } k)_{\mathbb{N}}^n) \cdot \text{DigA}(\text{DecSD}(x, n + 1, k), n + 1) - ((\text{Radix } k)_{\mathbb{N}}^{n+1}) \cdot \text{SDSubAddCarry}(\text{DigA}(\text{DecSD}(x, n + 1, k), n + 1), k)) + ((\text{Radix } k)_{\mathbb{N}}^n) \cdot \text{SDSubAddCarry}(\text{DigA}(\text{DecSD}(x, n + 1, k), n), k)$.

3. DEFINITION FOR ADDER OPERATION ON RADIX- 2^k SUB SIGNED-DIGIT NUMBER

Let i, n, k be natural numbers, let x be a n -tuple of k -SD_Sub, and let y be a n -tuple of k -SD_Sub. Let us assume that $i \in \text{Seg } n$ and $k \geq 2$. The functor $\text{SDSubAddDigit}(x, y, i, k)$ yields an element of k -SD_Sub and is defined as follows:

- (Def. 1) $\text{SDSubAddDigit}(x, y, i, k) = \text{SDSubAddData}(\text{DigA_SDSub}(x, i) + \text{DigA_SDSub}(y, i), k) + \text{SDSubAddCarry}(\text{DigA_SDSub}(x, i - 1) + \text{DigA_SDSub}(y, i - 1), k)$.

Let n, k be natural numbers and let x, y be n -tuples of k -SD_Sub. The functor $x' +' y$ yields a n -tuple of k -SD_Sub and is defined by:

- (Def. 2) For every i such that $i \in \text{Seg } n$ holds $\text{DigA_SDSub}(x' +' y, i) = \text{SDSubAddDigit}(x, y, i, k)$.

Next we state two propositions:

- (8) For every i such that $i \in \text{Seg } n$ holds if $2 \leq k$, then $\text{SDSubAddDigit}(\text{SD2SDSubDecSD}(x, n+1, k), \text{SD2SDSubDecSD}(y, n+1, k), i, k) = \text{SDSubAddDigit}(\text{SD2SDSubDecSD}(x \bmod (\text{Radix } k)_{\mathbb{N}}^n, n, k), \text{SD2SDSubDecSD}(y \bmod (\text{Radix } k)_{\mathbb{N}}^n, n, k), i, k)$.
- (9) Let given n . Suppose $n \geq 1$. Let given k, x, y . Suppose $k \geq 3$ and x is represented by n, k and y is represented by n, k . Then $x + y = \text{SDSub2IntOutSD2SDSubDecSD}(x, n, k)' + \text{SD2SDSubDecSD}(y, n, k)$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Joining of decorated trees. *Formalized Mathematics*, 4(1):77–82, 1993.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Marek Chmur. The lattice of natural numbers and the sublattice of it. The set of prime numbers. *Formalized Mathematics*, 2(4):453–459, 1991.
- [6] Yoshinori Fujisawa and Yasushi Fuwa. Definitions of radix- 2^k signed-digit number and its adder algorithm. *Formalized Mathematics*, 9(1):71–75, 2001.
- [7] Yoshinori Fujisawa, Yasushi Fuwa, and Hidetaka Shimizu. Euler's Theorem and small Fermat's Theorem. *Formalized Mathematics*, 7(1):123–126, 1998.
- [8] Andrzej Kondracki. The Chinese Remainder Theorem. *Formalized Mathematics*, 6(4):573–577, 1997.
- [9] Masaaki Niimura and Yasushi Fuwa. Improvement of radix- 2^k signed-digit number for high speed circuit. *Formalized Mathematics*, 11(2):133–137, 2003.
- [10] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [12] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [13] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received January 3, 2003
