

# Series on Complex Banach Algebra

Noboru Endou  
Gifu National College of Technology

**Summary.** This article is an extension of [20].

MML Identifier: CLOPBAN3.

The articles [22], [24], [25], [5], [6], [3], [2], [21], [11], [1], [23], [4], [15], [16], [17], [14], [12], [13], [19], [18], [10], [8], [9], [7], and [20] provide the notation and terminology for this paper.

## 1. BASIC PROPERTIES OF SEQUENCES OF NORM SPACE

Let  $X$  be a non empty complex normed space structure and let  $s_1$  be a sequence of  $X$ . The functor  $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}$  yielding a sequence of  $X$  is defined as follows:

(Def. 1)  $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(0) = s_1(0)$  and for every natural number  $n$  holds  $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(n+1) = (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(n) + s_1(n+1)$ .

One can prove the following proposition

(1) Let  $X$  be an add-associative right zeroed right complementable non empty complex normed space structure and  $s_1$  be a sequence of  $X$ . Suppose that for every natural number  $n$  holds  $s_1(n) = 0_X$ . Let  $m$  be a natural number. Then  $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(m) = 0_X$ .

Let  $X$  be a complex normed space and let  $s_1$  be a sequence of  $X$ . We say that  $s_1$  is summable if and only if:

(Def. 2)  $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}$  is convergent.

Let  $X$  be a complex normed space. One can verify that there exists a sequence of  $X$  which is summable.

Let  $X$  be a complex normed space and let  $s_1$  be a sequence of  $X$ . The functor  $\sum s_1$  yields an element of  $X$  and is defined by:

(Def. 3)  $\sum s_1 = \lim((\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}})$ .

Let  $X$  be a complex normed space and let  $s_1$  be a sequence of  $X$ . We say that  $s_1$  is norm-summable if and only if:

(Def. 4)  $\|s_1\|$  is summable.

The following propositions are true:

- (2) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  and for every natural number  $m$  holds  $0 \leq \|s_1\|(m)$ .
- (3) For every complex normed space  $X$  and for all elements  $x, y, z$  of  $X$  holds  $\|x - y\| = \|(x - z) + (z - y)\|$ .
- (4) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Suppose  $s_1$  is convergent. Let  $s$  be a real number. Suppose  $0 < s$ . Then there exists a natural number  $n$  such that for every natural number  $m$  if  $n \leq m$ , then  $\|s_1(m) - s_1(n)\| < s$ .
- (5) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Then  $s_1$  is Cauchy sequence by norm if and only if for every real number  $p$  such that  $p > 0$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $\|s_1(m) - s_1(n)\| < p$ .
- (6) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Suppose that for every natural number  $n$  holds  $s_1(n) = 0_X$ . Let  $m$  be a natural number. Then  $(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa \in \mathbb{N}}(m) = 0$ .

Let  $X$  be a complex normed space and let  $s_1$  be a sequence of  $X$ . Let us observe that  $s_1$  is constant if and only if:

(Def. 5) There exists an element  $r$  of  $X$  such that for every natural number  $n$  holds  $s_1(n) = r$ .

Let  $X$  be a complex normed space, let  $s_1$  be a sequence of  $X$ , and let  $k$  be a natural number. The functor  $s_1 \uparrow k$  yielding a sequence of  $X$  is defined as follows:

(Def. 6) For every natural number  $n$  holds  $(s_1 \uparrow k)(n) = s_1(n + k)$ .

Let  $X$  be a complex normed space and let  $s_1, s_2$  be sequences of  $X$ . We say that  $s_1$  is a subsequence of  $s_2$  if and only if:

(Def. 7) There exists an increasing sequence  $N_1$  of naturals such that  $s_1 = s_2 \cdot N_1$ .

Next we state a number of propositions:

- (7) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  holds  $s_1 \uparrow 0 = s_1$ .
- (8) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  and for all natural numbers  $k, m$  holds  $s_1 \uparrow k \uparrow m = s_1 \uparrow m \uparrow k$ .
- (9) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  and for all natural numbers  $k, m$  holds  $s_1 \uparrow k \uparrow m = s_1 \uparrow (k + m)$ .

- (10) Let  $X$  be a complex normed space and  $s_1, s_2$  be sequences of  $X$ . If  $s_2$  is a subsequence of  $s_1$  and  $s_1$  is convergent, then  $s_2$  is convergent.
- (11) Let  $X$  be a complex normed space and  $s_1, s_2$  be sequences of  $X$ . If  $s_2$  is a subsequence of  $s_1$  and  $s_1$  is convergent, then  $\lim s_2 = \lim s_1$ .
- (12) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $k$  be a natural number. Then  $s_1 \uparrow k$  is a subsequence of  $s_1$ .
- (13) Let  $X$  be a complex normed space,  $s_1, s_2$  be sequences of  $X$ , and  $k$  be a natural number. If  $s_1$  is convergent, then  $s_1 \uparrow k$  is convergent and  $\lim(s_1 \uparrow k) = \lim s_1$ .
- (14) Let  $X$  be a complex normed space and  $s_1, s_2$  be sequences of  $X$ . Suppose  $s_1$  is convergent and there exists a natural number  $k$  such that  $s_1 = s_2 \uparrow k$ . Then  $s_2$  is convergent.
- (15) Let  $X$  be a complex normed space and  $s_1, s_2$  be sequences of  $X$ . Suppose  $s_1$  is convergent and there exists a natural number  $k$  such that  $s_1 = s_2 \uparrow k$ . Then  $\lim s_2 = \lim s_1$ .
- (16) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  such that  $s_1$  is constant holds  $s_1$  is convergent.
- (17) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . If for every natural number  $n$  holds  $s_1(n) = 0_X$ , then  $s_1$  is norm-summable.

Let  $X$  be a complex normed space. Observe that there exists a sequence of  $X$  which is norm-summable.

The following three propositions are true:

- (18) Let  $X$  be a complex normed space and  $s$  be a sequence of  $X$ . If  $s$  is summable, then  $s$  is convergent and  $\lim s = 0_X$ .
- (19) For every complex normed space  $X$  and for all sequences  $s_3, s_4$  of  $X$  holds  $(\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa \in \mathbb{N}} + (\sum_{\alpha=0}^{\kappa}(s_4)(\alpha))_{\kappa \in \mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_3 + s_4)(\alpha))_{\kappa \in \mathbb{N}}$ .
- (20) For every complex normed space  $X$  and for all sequences  $s_3, s_4$  of  $X$  holds  $(\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa \in \mathbb{N}} - (\sum_{\alpha=0}^{\kappa}(s_4)(\alpha))_{\kappa \in \mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_3 - s_4)(\alpha))_{\kappa \in \mathbb{N}}$ .

Let  $X$  be a complex normed space and let  $s_1$  be a norm-summable sequence of  $X$ . Observe that  $\|s_1\|$  is summable.

Let  $X$  be a complex normed space. One can check that every sequence of  $X$  which is summable is also convergent.

The following two propositions are true:

- (21) Let  $X$  be a complex normed space and  $s_2, s_5$  be sequences of  $X$ . If  $s_2$  is summable and  $s_5$  is summable, then  $s_2 + s_5$  is summable and  $\sum(s_2 + s_5) = \sum s_2 + \sum s_5$ .
- (22) Let  $X$  be a complex normed space and  $s_2, s_5$  be sequences of  $X$ . If  $s_2$  is summable and  $s_5$  is summable, then  $s_2 - s_5$  is summable and  $\sum(s_2 - s_5) = \sum s_2 - \sum s_5$ .

Let  $X$  be a complex normed space and let  $s_2, s_5$  be summable sequences of  $X$ . One can check that  $s_2 + s_5$  is summable and  $s_2 - s_5$  is summable.

The following propositions are true:

- (23) For every complex normed space  $X$  and for every sequence  $s_1$  of  $X$  and for every complex number  $z$  holds  $(\sum_{\alpha=0}^{\kappa}(z \cdot s_1)(\alpha))_{\kappa \in \mathbb{N}} = z \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}$ .
- (24) Let  $X$  be a complex normed space,  $s_1$  be a summable sequence of  $X$ , and  $z$  be a complex number. Then  $z \cdot s_1$  is summable and  $\sum(z \cdot s_1) = z \cdot \sum s_1$ .

Let  $X$  be a complex normed space, let  $z$  be a complex number, and let  $s_1$  be a summable sequence of  $X$ . One can check that  $z \cdot s_1$  is summable.

Next we state two propositions:

- (25) Let  $X$  be a complex normed space and  $s, s_3$  be sequences of  $X$ . If for every natural number  $n$  holds  $s_3(n) = s(0)$ , then  $(\sum_{\alpha=0}^{\kappa}(s \uparrow 1)(\alpha))_{\kappa \in \mathbb{N}} = (\sum_{\alpha=0}^{\kappa} s(\alpha))_{\kappa \in \mathbb{N}} \uparrow 1 - s_3$ .
- (26) Let  $X$  be a complex normed space and  $s$  be a sequence of  $X$ . If  $s$  is summable, then for every natural number  $n$  holds  $s \uparrow n$  is summable.

Let  $X$  be a complex normed space, let  $s_1$  be a summable sequence of  $X$ , and let  $n$  be a natural number. Observe that  $s_1 \uparrow n$  is summable.

We now state the proposition

- (27) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Then  $(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa \in \mathbb{N}}$  is upper bounded if and only if  $s_1$  is norm-summable.
- Let  $X$  be a complex normed space and let  $s_1$  be a norm-summable sequence of  $X$ . Note that  $(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa \in \mathbb{N}}$  is upper bounded.

The following propositions are true:

- (28) Let  $X$  be a complex Banach space and  $s_1$  be a sequence of  $X$ . Then  $s_1$  is summable if and only if for every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $\|(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(n)\| < p$ .
- (29) Let  $X$  be a complex normed space,  $s$  be a sequence of  $X$ , and  $n, m$  be natural numbers. If  $n \leq m$ , then  $\|(\sum_{\alpha=0}^{\kappa} s(\alpha))_{\kappa \in \mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa} s(\alpha))_{\kappa \in \mathbb{N}}(n)\| \leq |(\sum_{\alpha=0}^{\kappa} \|s\|(\alpha))_{\kappa \in \mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa} \|s\|(\alpha))_{\kappa \in \mathbb{N}}(n)|$ .
- (30) For every complex Banach space  $X$  and for every sequence  $s_1$  of  $X$  such that  $s_1$  is norm-summable holds  $s_1$  is summable.
- (31) Let  $X$  be a complex normed space,  $r_1$  be a sequence of real numbers, and  $s_5$  be a sequence of  $X$ . Suppose  $r_1$  is summable and there exists a natural number  $m$  such that for every natural number  $n$  such that  $m \leq n$  holds  $\|s_5(n)\| \leq r_1(n)$ . Then  $s_5$  is norm-summable.
- (32) Let  $X$  be a complex normed space and  $s_2, s_5$  be sequences of  $X$ . Suppose for every natural number  $n$  holds  $0 \leq \|s_2\|(n)$  and  $\|s_2\|(n) \leq \|s_5\|(n)$  and  $s_5$  is norm-summable. Then  $s_2$  is norm-summable and  $\sum \|s_2\| \leq \sum \|s_5\|$ .

- (33) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Suppose that
- (i) for every natural number  $n$  holds  $\|s_1\|(n) > 0$ , and
  - (ii) there exists a natural number  $m$  such that for every natural number  $n$  such that  $n \geq m$  holds  $\frac{\|s_1\|(n+1)}{\|s_1\|(n)} \geq 1$ .
- Then  $s_1$  is not norm-summable.
- (34) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $r_1$  be a sequence of real numbers. Suppose for every natural number  $n$  holds  $r_1(n) = \sqrt[n]{\|s_1\|(n)}$  and  $r_1$  is convergent and  $\lim r_1 < 1$ . Then  $s_1$  is norm-summable.
- (35) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $r_1$  be a sequence of real numbers. Suppose that
- (i) for every natural number  $n$  holds  $r_1(n) = \sqrt[n]{\|s_1\|(n)}$ , and
  - (ii) there exists a natural number  $m$  such that for every natural number  $n$  such that  $m \leq n$  holds  $r_1(n) \geq 1$ .
- Then  $\|s_1\|$  is not summable.
- (36) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $r_1$  be a sequence of real numbers. Suppose for every natural number  $n$  holds  $r_1(n) = \sqrt[n]{\|s_1\|(n)}$  and  $r_1$  is convergent and  $\lim r_1 > 1$ . Then  $s_1$  is not norm-summable.
- (37) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $r_1$  be a sequence of real numbers. Suppose  $\|s_1\|$  is non-increasing and for every natural number  $n$  holds  $r_1(n) = 2^n \cdot \|s_1\|(2^n)$ . Then  $s_1$  is norm-summable if and only if  $r_1$  is summable.
- (38) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $p$  be a real number. Suppose  $p > 1$  and for every natural number  $n$  such that  $n \geq 1$  holds  $\|s_1\|(n) = \frac{1}{n^p}$ . Then  $s_1$  is norm-summable.
- (39) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $p$  be a real number. Suppose  $p \leq 1$  and for every natural number  $n$  such that  $n \geq 1$  holds  $\|s_1\|(n) = \frac{1}{n^p}$ . Then  $s_1$  is not norm-summable.
- (40) Let  $X$  be a complex normed space,  $s_1$  be a sequence of  $X$ , and  $r_1$  be a sequence of real numbers. Suppose for every natural number  $n$  holds  $s_1(n) \neq 0_X$  and  $r_1(n) = \frac{\|s_1\|(n+1)}{\|s_1\|(n)}$  and  $r_1$  is convergent and  $\lim r_1 < 1$ . Then  $s_1$  is norm-summable.
- (41) Let  $X$  be a complex normed space and  $s_1$  be a sequence of  $X$ . Suppose that
- (i) for every natural number  $n$  holds  $s_1(n) \neq 0_X$ , and
  - (ii) there exists a natural number  $m$  such that for every natural number  $n$  such that  $n \geq m$  holds  $\frac{\|s_1\|(n+1)}{\|s_1\|(n)} \geq 1$ .
- Then  $s_1$  is not norm-summable.

Let  $X$  be a complex Banach space. One can check that every sequence of  $X$  which is norm-summable is also summable.

## 2. BASIC PROPERTIES OF SEQUENCE OF BANACH ALGEBRA

The scheme *ExNCBCASeq* deals with a non empty normed complex algebra structure  $\mathcal{A}$  and a unary functor  $\mathcal{F}$  yielding a point of  $\mathcal{A}$ , and states that:

There exists a sequence  $S$  of  $\mathcal{A}$  such that for every natural number  $n$  holds  $S(n) = \mathcal{F}(n)$

for all values of the parameters.

We now state the proposition

- (42) Let  $X$  be a complex Banach algebra,  $x, y, z$  be elements of  $X$ , and  $a, b$  be complex numbers. Then  $x + y = y + x$  and  $(x + y) + z = x + (y + z)$  and  $x + 0_X = x$  and there exists an element  $t$  of  $X$  such that  $x + t = 0_X$  and  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$  and  $1_{\mathbb{C}} \cdot x = x$  and  $0_{\mathbb{C}} \cdot x = 0_X$  and  $a \cdot 0_X = 0_X$  and  $(-1_{\mathbb{C}}) \cdot x = -x$  and  $x \cdot 1_X = x$  and  $1_X \cdot x = x$  and  $x \cdot (y + z) = x \cdot y + x \cdot z$  and  $(y + z) \cdot x = y \cdot x + z \cdot x$  and  $a \cdot (x \cdot y) = (a \cdot x) \cdot y$  and  $a \cdot (x + y) = a \cdot x + a \cdot y$  and  $(a + b) \cdot x = a \cdot x + b \cdot x$  and  $(a \cdot b) \cdot x = a \cdot (b \cdot x)$  and  $(a \cdot b) \cdot (x \cdot y) = a \cdot x \cdot (b \cdot y)$  and  $a \cdot (x \cdot y) = x \cdot (a \cdot y)$  and  $0_X \cdot x = 0_X$  and  $x \cdot 0_X = 0_X$  and  $x \cdot (y - z) = x \cdot y - x \cdot z$  and  $(y - z) \cdot x = y \cdot x - z \cdot x$  and  $(x + y) - z = x + (y - z)$  and  $(x - y) + z = x - (y - z)$  and  $x - y - z = x - (y + z)$  and  $x + y = (x - z) + (z + y)$  and  $x - y = (x - z) + (z - y)$  and  $x = (x - y) + y$  and  $x = y - (y - x)$  and  $\|x\| = 0$  iff  $x = 0_X$  and  $\|a \cdot x\| = |a| \cdot \|x\|$  and  $\|x + y\| \leq \|x\| + \|y\|$  and  $\|x \cdot y\| \leq \|x\| \cdot \|y\|$  and  $\|1_X\| = 1$  and  $X$  is complete.

Let  $X$  be a non empty normed complex algebra structure, let  $S$  be a sequence of  $X$ , and let  $a$  be an element of  $X$ . The functor  $a \cdot S$  yields a sequence of  $X$  and is defined by:

- (Def. 8) For every natural number  $n$  holds  $(a \cdot S)(n) = a \cdot S(n)$ .

Let  $X$  be a non empty normed complex algebra structure, let  $S$  be a sequence of  $X$ , and let  $a$  be an element of  $X$ . The functor  $S \cdot a$  yields a sequence of  $X$  and is defined by:

- (Def. 9) For every natural number  $n$  holds  $(S \cdot a)(n) = S(n) \cdot a$ .

Let  $X$  be a non empty normed complex algebra structure and let  $s_2, s_5$  be sequences of  $X$ . The functor  $s_2 \cdot s_5$  yielding a sequence of  $X$  is defined by:

- (Def. 10) For every natural number  $n$  holds  $(s_2 \cdot s_5)(n) = s_2(n) \cdot s_5(n)$ .

Let  $X$  be a complex Banach algebra and let  $x$  be an element of  $X$ . Let us assume that  $x$  is invertible. The functor  $x^{-1}$  yields an element of  $X$  and is defined as follows:

- (Def. 11)  $x \cdot x^{-1} = 1_X$  and  $x^{-1} \cdot x = 1_X$ .

Let  $X$  be a complex Banach algebra and let  $z$  be an element of  $X$ . The functor  $(z^\kappa)_{\kappa \in \mathbb{N}}$  yielding a sequence of  $X$  is defined as follows:

(Def. 12)  $(z^\kappa)_{\kappa \in \mathbb{N}}(0) = \mathbf{1}_X$  and for every natural number  $n$  holds  $(z^\kappa)_{\kappa \in \mathbb{N}}(n+1) = (z^\kappa)_{\kappa \in \mathbb{N}}(n) \cdot z$ .

Let  $X$  be a complex Banach algebra, let  $z$  be an element of  $X$ , and let  $n$  be a natural number. The functor  $z_{\mathbb{N}}^n$  yielding an element of  $X$  is defined as follows:

(Def. 13)  $z_{\mathbb{N}}^n = (z^\kappa)_{\kappa \in \mathbb{N}}(n)$ .

The following propositions are true:

- (43) For every complex Banach algebra  $X$  and for every element  $z$  of  $X$  holds  $z_{\mathbb{N}}^0 = \mathbf{1}_X$ .
- (44) For every complex Banach algebra  $X$  and for every element  $z$  of  $X$  such that  $\|z\| < 1$  holds  $(z^\kappa)_{\kappa \in \mathbb{N}}$  is summable and norm-summable.
- (45) Let  $X$  be a complex Banach algebra and  $x$  be a point of  $X$ . If  $\|\mathbf{1}_X - x\| < 1$ , then  $((\mathbf{1}_X - x)^\kappa)_{\kappa \in \mathbb{N}}$  is summable and  $((\mathbf{1}_X - x)^\kappa)_{\kappa \in \mathbb{N}}$  is norm-summable.
- (46) For every complex Banach algebra  $X$  and for every point  $x$  of  $X$  such that  $\|\mathbf{1}_X - x\| < 1$  holds  $x$  is invertible and  $x^{-1} = \sum(((\mathbf{1}_X - x)^\kappa)_{\kappa \in \mathbb{N}})$ .

#### REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [4] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [7] Noboru Endou. Banach algebra of bounded complex linear operators. *Formalized Mathematics*, 12(3):237–242, 2004.
- [8] Noboru Endou. Banach space of absolute summable complex sequences. *Formalized Mathematics*, 12(2):191–194, 2004.
- [9] Noboru Endou. Complex Banach space of bounded linear operators. *Formalized Mathematics*, 12(2):201–209, 2004.
- [10] Noboru Endou. Complex linear space and complex normed space. *Formalized Mathematics*, 12(2):93–102, 2004.
- [11] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [12] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [13] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [14] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [15] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [16] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [17] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.

- [18] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(2):213–216, 1991.
- [19] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Formalized Mathematics*, 2(4):449–452, 1991.
- [20] Yasunari Shidama. The series on Banach algebra. *Formalized Mathematics*, 12(2):131–138, 2004.
- [21] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [22] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [23] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [24] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [25] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

*Received April 6, 2004*

---