

Eigenvalues of a Linear Transformation

Karol Pąk
Institute of Computer Science
University of Białystok
Poland

Summary. The article presents well known facts about eigenvalues of linear transformation of a vector space (see [13]). I formalize main dependencies between eigenvalues and the diagram of the matrix of a linear transformation over a finite-dimensional vector space. Finally, I formalize the subspace $\bigcup_{i=0}^{\infty} \text{Ker}(f - \lambda I)^i$ called a generalized eigenspace for the eigenvalue λ and show its basic properties.

MML identifier: VECTSP11, version: 7.9.03 4.108.1028

The articles [11], [33], [2], [3], [12], [34], [8], [10], [9], [5], [31], [27], [15], [7], [14], [32], [35], [25], [30], [29], [28], [26], [6], [22], [16], [23], [20], [1], [19], [4], [21], [17], [18], and [24] provide the notation and terminology for this paper.

1. PRELIMINARIES

We adopt the following convention: i, j, m, n denote natural numbers, K denotes a field, and a denotes an element of K .

Next we state several propositions:

- (1) Let A, B be matrices over K , n_1 be an element of \mathbb{N}^n , and m_1 be an element of \mathbb{N}^m . If $\text{rng } n_1 \times \text{rng } m_1 \subseteq$ the indices of A , then $\text{Segm}(A + B, n_1, m_1) = \text{Segm}(A, n_1, m_1) + \text{Segm}(B, n_1, m_1)$.
- (2) For every without zero finite subset P of \mathbb{N} such that $P \subseteq \text{Seg } n$ holds $\text{Segm}(I_K^{n \times n}, P, P) = I_K^{\text{card } P \times \text{card } P}$.
- (3) Let A, B be matrices over K and P, Q be without zero finite subsets of \mathbb{N} . If $P \times Q \subseteq$ the indices of A , then $\text{Segm}(A + B, P, Q) = \text{Segm}(A, P, Q) + \text{Segm}(B, P, Q)$.

- (4) For all square matrices A, B over K of dimension n such that $i, j \in \text{Seg } n$ holds $\text{Delete}(A + B, i, j) = \text{Delete}(A, i, j) + \text{Delete}(B, i, j)$.
- (5) For every square matrix A over K of dimension n such that $i, j \in \text{Seg } n$ holds $\text{Delete}(a \cdot A, i, j) = a \cdot \text{Delete}(A, i, j)$.
- (6) If $i \in \text{Seg } n$, then $\text{Delete}(I_K^{n \times n}, i, i) = I_K^{(n-1) \times (n-1)}$.
- (7) Let A, B be square matrices over K of dimension n . Then there exists a polynomial P of K such that $\text{len } P \leq n + 1$ and for every element x of K holds $\text{eval}(P, x) = \text{Det}(A + x \cdot B)$.
- (8) Let A be a square matrix over K of dimension n . Then there exists a polynomial P of K such that $\text{len } P = n + 1$ and for every element x of K holds $\text{eval}(P, x) = \text{Det}(A + x \cdot I_K^{n \times n})$.

Let us consider K . Observe that there exists a vector space over K which is non trivial and finite dimensional.

2. MAPS WITH EIGENVALUES

Let R be a non empty double loop structure, let V be a non empty vector space structure over R , and let I_1 be a function from V into V . We say that I_1 has eigenvalues if and only if:

- (Def. 1) There exists a vector v of V and there exists a scalar a of R such that $v \neq 0_V$ and $I_1(v) = a \cdot v$.

For simplicity, we follow the rules: V denotes a non trivial vector space over K , V_1, V_2 denote vector spaces over K , f denotes a linear transformation from V_1 to V_1 , v, w denote vectors of V , v_1 denotes a vector of V_1 , and L denotes a scalar of K .

Let us consider K, V . One can verify that there exists a linear transformation from V to V which has eigenvalues.

Let R be a non empty double loop structure, let V be a non empty vector space structure over R , and let f be a function from V into V . Let us assume that f has eigenvalues. An element of R is called an eigenvalue of f if:

- (Def. 2) There exists a vector v of V such that $v \neq 0_V$ and $f(v) = \text{it} \cdot v$.

Let R be a non empty double loop structure, let V be a non empty vector space structure over R , let f be a function from V into V , and let L be a scalar of R . Let us assume that f has eigenvalues and L is an eigenvalue of f . A vector of V is called an eigenvector of f and L if:

- (Def. 3) $f(\text{it}) = L \cdot \text{it}$.

We now state several propositions:

- (9) Let given a . Suppose $a \neq 0_K$. Let f be a function from V into V with eigenvalues and L be an eigenvalue of f . Then
 - (i) $a \cdot f$ has eigenvalues,

- (ii) $a \cdot L$ is an eigenvalue of $a \cdot f$, and
 - (iii) w is an eigenvector of f and L iff w is an eigenvector of $a \cdot f$ and $a \cdot L$.
- (10) Let f_1, f_2 be functions from V into V with eigenvalues and L_1, L_2 be scalars of K . Suppose that
- (i) L_1 is an eigenvalue of f_1 ,
 - (ii) L_2 is an eigenvalue of f_2 , and
 - (iii) there exists v such that v is an eigenvector of f_1 and L_1 and an eigenvector of f_2 and L_2 and $v \neq 0_V$.

Then

- (iv) $f_1 + f_2$ has eigenvalues,
 - (v) $L_1 + L_2$ is an eigenvalue of $f_1 + f_2$, and
 - (vi) for every w such that w is an eigenvector of f_1 and L_1 and an eigenvector of f_2 and L_2 holds w is an eigenvector of $f_1 + f_2$ and $L_1 + L_2$.
- (11) id_V has eigenvalues and $\mathbf{1}_K$ is an eigenvalue of id_V and every v is an eigenvector of id_V and $\mathbf{1}_K$.
- (12) For every eigenvalue L of id_V holds $L = \mathbf{1}_K$.
- (13) If $\ker f$ is non trivial, then f has eigenvalues and 0_K is an eigenvalue of f .
- (14) f has eigenvalues and L is an eigenvalue of f iff $\ker f + (-L) \cdot \text{id}_{(V_1)}$ is non trivial.
- (15) Let V_1 be a finite dimensional vector space over K , b_1, b'_1 be ordered bases of V_1 , and f be a linear transformation from V_1 to V_1 . Then f has eigenvalues and L is an eigenvalue of f if and only if $\text{Det AutEqMt}(f + (-L) \cdot \text{id}_{(V_1)}, b_1, b'_1) = 0_K$.
- (16) Let K be an algebraic-closed field and V_1 be a non trivial finite dimensional vector space over K . Then every linear transformation from V_1 to V_1 has eigenvalues.
- (17) Let given f, L . Suppose f has eigenvalues and L is an eigenvalue of f . Then v_1 is an eigenvector of f and L if and only if $v_1 \in \ker f + (-L) \cdot \text{id}_{(V_1)}$.

Let S be a 1-sorted structure, let F be a function from S into S , and let n be a natural number. The functor F^n yields a function from S into S and is defined as follows:

- (Def. 4) For every element F' of the semigroup of functions onto the carrier of S such that $F' = F$ holds $F^n = \prod(n \mapsto F')$.

In the sequel S denotes a 1-sorted structure and F denotes a function from S into S .

Next we state several propositions:

- (18) $F^0 = \text{id}_S$.
- (19) $F^1 = F$.
- (20) $F^{i+j} = F^i \cdot F^j$.

- (21) For all elements s_1, s_2 of S and for all n, m such that $F^m(s_1) = s_2$ and $F^n(s_2) = s_2$ holds $F^{m+i \cdot n}(s_1) = s_2$.
- (22) Let K be an add-associative right zeroed right complementable Abelian associative well unital distributive non empty double loop structure, V_1 be an Abelian add-associative right zeroed right complementable vector space-like non empty vector space structure over K , W be a subspace of V_1 , f be a function from V_1 into V_1 , and f_3 be a function from W into W . If $f_3 = f \upharpoonright W$, then $f^n \upharpoonright W = f_3^n$.

Let us consider K, V_1 , let f be a linear transformation from V_1 to V_1 , and let n be a natural number. Then f^n is a linear transformation from V_1 to V_1 .

We now state the proposition

- (23) If $f^i(v_1) = 0_{(V_1)}$, then $f^{i+j}(v_1) = 0_{(V_1)}$.

3. GENERALIZED EIGENSPACE OF A LINEAR TRANSFORMATION

Let us consider K, V_1, f . The functor $\text{UnionKers } f$ yielding a strict subspace of V_1 is defined by:

- (Def. 5) The carrier of $\text{UnionKers } f = \{v; v \text{ ranges over vectors of } V_1: \bigvee_n f^n(v) = 0_{(V_1)}\}$.

We now state a number of propositions:

- (24) $v_1 \in \text{UnionKers } f$ iff there exists n such that $f^n(v_1) = 0_{(V_1)}$.
- (25) $\ker f^i$ is a subspace of $\text{UnionKers } f$.
- (26) $\ker f^i$ is a subspace of $\ker f^{i+j}$.
- (27) Let V be a finite dimensional vector space over K and f be a linear transformation from V to V . Then there exists n such that $\text{UnionKers } f = \ker f^n$.
- (28) $f \upharpoonright \ker f^n$ is a linear transformation from $\ker f^n$ to $\ker f^n$.
- (29) $f \upharpoonright \ker (f + L \cdot \text{id}_{(V_1)})^n$ is a linear transformation from $\ker (f + L \cdot \text{id}_{(V_1)})^n$ to $\ker (f + L \cdot \text{id}_{(V_1)})^n$.
- (30) $f \upharpoonright \text{UnionKers } f$ is a linear transformation from $\text{UnionKers } f$ to $\text{UnionKers } f$.
- (31) $f \upharpoonright \text{UnionKers}(f + L \cdot \text{id}_{(V_1)})$ is a linear transformation from $\text{UnionKers}(f + L \cdot \text{id}_{(V_1)})$ to $\text{UnionKers}(f + L \cdot \text{id}_{(V_1)})$.
- (32) $f \upharpoonright \text{im}(f^n)$ is a linear transformation from $\text{im}(f^n)$ to $\text{im}(f^n)$.
- (33) $f \upharpoonright \text{im}((f + L \cdot \text{id}_{(V_1)})^n)$ is a linear transformation from $\text{im}((f + L \cdot \text{id}_{(V_1)})^n)$ to $\text{im}((f + L \cdot \text{id}_{(V_1)})^n)$.
- (34) If $\text{UnionKers } f = \ker f^n$, then $\ker f^n \cap \text{im}(f^n) = \mathbf{0}_{(V_1)}$.

- (35) Let V be a finite dimensional vector space over K , f be a linear transformation from V to V , and given n . If $\text{UnionKers } f = \ker f^n$, then V is the direct sum of $\ker f^n$ and $\text{im}(f^n)$.
- (36) For every linear complement I of $\text{UnionKers } f$ holds $f|I$ is one-to-one.
- (37) Let I be a linear complement of $\text{UnionKers}(f + (-L) \cdot \text{id}_{(V_1)})$ and f_4 be a linear transformation from I to I . If $f_4 = f|I$, then for every vector v of I such that $f_4(v) = L \cdot v$ holds $v = 0_{(V_1)}$.
- (38) Suppose $n \geq 1$. Then there exists a linear transformation h from V_1 to V_1 such that $(f + L \cdot \text{id}_{(V_1)})^n = f \cdot h + (L \cdot \text{id}_{(V_1)})^n$ and for every i holds $f^i \cdot h = h \cdot f^i$.
- (39) Let L_1, L_2 be scalars of K . Suppose f has eigenvalues and $L_1 \neq L_2$ and L_1 is an eigenvalue of f and L_2 is an eigenvalue of f . Let I be a linear complement of $\text{UnionKers}(f + (-L_1) \cdot \text{id}_{(V_1)})$ and f_4 be a linear transformation from I to I . Suppose $f_4 = f|I$. Then f_4 has eigenvalues and L_1 is not an eigenvalue of f_4 and L_2 is an eigenvalue of f_4 and $\text{UnionKers}(f + (-L_2) \cdot \text{id}_{(V_1)})$ is a subspace of I .
- (40) Let U be a finite subset of V_1 . Suppose U is linearly independent. Let u be a vector of V_1 . Suppose $u \in U$. Let L be a linear combination of $U \setminus \{u\}$. Then $\overline{U} = \overline{(U \setminus \{u\}) \cup \{u + \sum L\}}$ and $(U \setminus \{u\}) \cup \{u + \sum L\}$ is linearly independent.
- (41) Let A be a subset of V_1 , L be a linear combination of V_2 , and f be a linear transformation from V_1 to V_2 . Suppose the support of $L \subseteq f^\circ A$. Then there exists a linear combination M of A such that $f(\sum M) = \sum L$.
- (42) Let f be a linear transformation from V_1 to V_2 , A be a subset of V_1 , and B be a subset of V_2 . If $f^\circ A = B$, then $f^\circ(\text{the carrier of } \text{Lin}(A)) = \text{the carrier of } \text{Lin}(B)$.
- (43) Let L be a linear combination of V_1 , F be a finite sequence of elements of V_1 , and f be a linear transformation from V_1 to V_2 . Suppose $f|((\text{the support of } L) \cap \text{rng } F)$ is one-to-one and $\text{rng } F \subseteq \text{the support of } L$. Then there exists a linear combination L_3 of V_2 such that
 - (i) the support of $L_3 = f^\circ((\text{the support of } L) \cap \text{rng } F)$,
 - (ii) $f \cdot (L F) = L_3 (f \cdot F)$, and
 - (iii) for every v_1 such that $v_1 \in (\text{the support of } L) \cap \text{rng } F$ holds $L(v_1) = L_3(f(v_1))$.
- (44) Let A, B be subsets of V_1 and L be a linear combination of V_1 . Suppose the support of $L \subseteq A \cup B$ and $\sum L = 0_{(V_1)}$. Let f be a linear function from V_1 into V_2 . Suppose $f|B$ is one-to-one and $f^\circ B$ is a linearly independent subset of V_2 and $f^\circ A \subseteq \{0_{(V_2)}\}$. Then the support of $L \subseteq A$.

REFERENCES

- [1] Jesse Alama. The rank+nullity theorem. *Formalized Mathematics*, 15(3):137–142, 2007.
- [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [4] Grzegorz Bancerek. Monoids. *Formalized Mathematics*, 3(2):213–225, 1992.
- [5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [6] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [7] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [8] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [9] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [10] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [13] I.N. Herstein and David J. Winter. *Matrix Theory and Linear Algebra*. Macmillan, 1988.
- [14] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Formalized Mathematics*, 2(4):475–480, 1991.
- [15] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [16] Robert Milewski. Associated matrix of linear map. *Formalized Mathematics*, 5(3):339–345, 1996.
- [17] Robert Milewski. The evaluation of polynomials. *Formalized Mathematics*, 9(2):391–395, 2001.
- [18] Robert Milewski. Fundamental theorem of algebra. *Formalized Mathematics*, 9(3):461–470, 2001.
- [19] Robert Milewski. The ring of polynomials. *Formalized Mathematics*, 9(2):339–346, 2001.
- [20] Michał Muzalewski. Rings and modules – part II. *Formalized Mathematics*, 2(4):579–585, 1991.
- [21] Michał Muzalewski and Lesław W. Szczerba. Construction of finite sequences over ring and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(1):97–104, 1991.
- [22] Karol Pak. Basic properties of the rank of matrices over a field. *Formalized Mathematics*, 15(4):199–211, 2007.
- [23] Karol Pak and Andrzej Trybulec. Laplace expansion. *Formalized Mathematics*, 15(3):143–150, 2007.
- [24] Karol Pak. Linear map of matrices. *Formalized Mathematics*, 16(3):269–275, 2008.
- [25] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [26] Wojciech A. Trybulec. Basis of vector space. *Formalized Mathematics*, 1(5):883–885, 1990.
- [27] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [28] Wojciech A. Trybulec. Linear combinations in vector space. *Formalized Mathematics*, 1(5):877–882, 1990.
- [29] Wojciech A. Trybulec. Operations on subspaces in vector space. *Formalized Mathematics*, 1(5):871–876, 1990.
- [30] Wojciech A. Trybulec. Subspaces and cosets of subspaces in vector space. *Formalized Mathematics*, 1(5):865–870, 1990.
- [31] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [32] Wojciech A. Trybulec. Lattice of subgroups of a group. Frattini subgroup. *Formalized Mathematics*, 2(1):41–47, 1991.
- [33] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [34] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

- [35] Katarzyna Zawadzka. The product and the determinant of matrices with entries in a field. *Formalized Mathematics*, 4(1):1–8, 1993.

Received July 11, 2008
