

# The Definition of Topological Manifolds

Marco Riccardi  
Via del Pero 102  
54038 Montignoso, Italy

**Summary.** This article introduces the definition of  $n$ -locally Euclidean topological spaces and topological manifolds [13].

MML identifier: MFOLD\_1, version: 7.11.07 4.156.1112

The papers [8], [1], [6], [15], [7], [18], [3], [4], [17], [2], [16], [9], [19], [20], [11], [12], [10], [14], and [5] provide the terminology and notation for this paper.

## 1. PRELIMINARIES

Let  $x, y$  be sets. Observe that  $\{\langle x, y \rangle\}$  is one-to-one.

In the sequel  $n$  denotes a natural number.

One can prove the following two propositions:

- (1) For every non empty topological space  $T$  holds  $T$  and  $T|\Omega_T$  are homeomorphic.
- (2) Let  $X$  be a non empty subspace of  $\mathcal{E}_T^n$  and  $f$  be a function from  $X$  into  $\mathbb{R}^1$ . Suppose  $f$  is continuous. Then there exists a function  $g$  from  $X$  into  $\mathcal{E}_T^n$  such that
  - (i) for every point  $a$  of  $X$  and for every point  $b$  of  $\mathcal{E}_T^n$  and for every real number  $r$  such that  $a = b$  and  $f(a) = r$  holds  $g(b) = r \cdot b$ , and
  - (ii)  $g$  is continuous.

Let us consider  $n$  and let  $S$  be a subset of  $\mathcal{E}_T^n$ . We say that  $S$  is ball if and only if:

- (Def. 1) There exists a point  $p$  of  $\mathcal{E}_T^n$  and there exists a real number  $r$  such that  $S = \text{Ball}(p, r)$ .

Let us consider  $n$ . Observe that there exists a subset of  $\mathcal{E}_T^n$  which is ball and every subset of  $\mathcal{E}_T^n$  which is ball is also open.

Let us consider  $n$ . One can verify that there exists a subset of  $\mathcal{E}_T^n$  which is non empty and ball.

In the sequel  $p$  denotes a point of  $\mathcal{E}_T^n$  and  $r$  denotes a real number.

The following proposition is true

- (3) For every open subset  $S$  of  $\mathcal{E}_T^n$  such that  $p \in S$  there exists ball subset  $B$  of  $\mathcal{E}_T^n$  such that  $B \subseteq S$  and  $p \in B$ .

Let us consider  $n, p, r$ . The functor  $\mathbb{B}_r(p)$  yields a subspace of  $\mathcal{E}_T^n$  and is defined as follows:

(Def. 2)  $\mathbb{B}_r(p) = \mathcal{E}_T^n \upharpoonright \text{Ball}(p, r)$ .

Let us consider  $n$ . The functor  $\mathbb{B}^n$  yields a subspace of  $\mathcal{E}_T^n$  and is defined as follows:

(Def. 3)  $\mathbb{B}^n = \mathbb{B}_1(0_{\mathcal{E}_T^n})$ .

Let us consider  $n$ . One can verify that  $\mathbb{B}^n$  is non empty. Let us consider  $p$  and let  $s$  be a positive real number. Observe that  $\mathbb{B}_s(p)$  is non empty.

The following propositions are true:

- (4) The carrier of  $\mathbb{B}_r(p) = \text{Ball}(p, r)$ .
- (5) If  $n \neq 0$  and  $p$  is a point of  $\mathbb{B}^n$ , then  $|p| < 1$ .
- (6) Let  $f$  be a function from  $\mathbb{B}^n$  into  $\mathcal{E}_T^n$ . Suppose  $n \neq 0$  and for every point  $a$  of  $\mathbb{B}^n$  and for every point  $b$  of  $\mathcal{E}_T^n$  such that  $a = b$  holds  $f(a) = \frac{1}{1-|b| \cdot |b|} \cdot b$ . Then  $f$  is homeomorphism.
- (7) Let  $r$  be a positive real number and  $f$  be a function from  $\mathbb{B}^n$  into  $\mathbb{B}_r(p)$ . Suppose  $n \neq 0$  and for every point  $a$  of  $\mathbb{B}^n$  and for every point  $b$  of  $\mathcal{E}_T^n$  such that  $a = b$  holds  $f(a) = r \cdot b + p$ . Then  $f$  is homeomorphism.
- (8)  $\mathbb{B}^n$  and  $\mathcal{E}_T^n$  are homeomorphic.

In the sequel  $q$  denotes a point of  $\mathcal{E}_T^n$ .

We now state three propositions:

- (9) For all positive real numbers  $r, s$  holds  $\mathbb{B}_r(p)$  and  $\mathbb{B}_s(q)$  are homeomorphic.
- (10) For every non empty ball subset  $B$  of  $\mathcal{E}_T^n$  holds  $B$  and  $\Omega_{\mathcal{E}_T^n}$  are homeomorphic.
- (11) Let  $M, N$  be non empty topological spaces,  $p$  be a point of  $M$ ,  $U$  be a neighbourhood of  $p$ , and  $B$  be an open subset of  $N$ . Suppose  $U$  and  $B$  are homeomorphic. Then there exists an open subset  $V$  of  $M$  and there exists an open subset  $S$  of  $N$  such that  $V \subseteq U$  and  $p \in V$  and  $V$  and  $S$  are homeomorphic.

## 2. MANIFOLD

In the sequel  $M$  is a non empty topological space.

Let us consider  $n, M$ . We say that  $M$  is  $n$ -locally Euclidean if and only if the condition (Def. 4) is satisfied.

(Def. 4) Let  $p$  be a point of  $M$ . Then there exists a neighbourhood  $U$  of  $p$  and there exists an open subset  $S$  of  $\mathcal{E}_T^n$  such that  $U$  and  $S$  are homeomorphic.

Let us consider  $n$ . Observe that  $\mathcal{E}_T^n$  is  $n$ -locally Euclidean.

Let us consider  $n$ . Observe that there exists a non empty topological space which is  $n$ -locally Euclidean.

We now state two propositions:

(12)  $M$  is  $n$ -locally Euclidean if and only if for every point  $p$  of  $M$  there exists a neighbourhood  $U$  of  $p$  and there exists ball subset  $B$  of  $\mathcal{E}_T^n$  such that  $U$  and  $B$  are homeomorphic.

(13)  $M$  is  $n$ -locally Euclidean if and only if for every point  $p$  of  $M$  there exists a neighbourhood  $U$  of  $p$  such that  $U$  and  $\Omega_{\mathcal{E}_T^n}$  are homeomorphic.

Let us consider  $n$ . Observe that every non empty topological space which is  $n$ -locally Euclidean is also first-countable.

Let us note that every non empty topological space which is 0-locally Euclidean is also discrete and every non empty topological space which is discrete is also 0-locally Euclidean.

Let us consider  $n$ . One can verify that  $\mathcal{E}_T^n$  is second-countable.

Let us consider  $n$ . Note that there exists a non empty topological space which is second-countable, Hausdorff, and  $n$ -locally Euclidean.

Let us consider  $n, M$ . We say that  $M$  is  $n$ -manifold if and only if:

(Def. 5)  $M$  is second-countable, Hausdorff, and  $n$ -locally Euclidean.

Let us consider  $M$ . We say that  $M$  is manifold-like if and only if:

(Def. 6) There exists  $n$  such that  $M$  is  $n$ -manifold.

Let us consider  $n$ . Observe that there exists a non empty topological space which is  $n$ -manifold.

Let us consider  $n$ . One can check the following observations:

- \* every non empty topological space which is  $n$ -manifold is also second-countable, Hausdorff, and  $n$ -locally Euclidean,
- \* every non empty topological space which is second-countable, Hausdorff, and  $n$ -locally Euclidean is also  $n$ -manifold, and
- \* every non empty topological space which is  $n$ -manifold is also manifold-like.

Let us note that every non empty topological space which is second-countable and discrete is also 0-manifold.

Let us consider  $n$  and let  $M$  be an  $n$ -manifold non empty topological space. One can verify that every non empty subspace of  $M$  which is open is also  $n$ -manifold.

Let us note that there exists a non empty topological space which is manifold-like.

A manifold is a manifold-like non empty topological space.

#### REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [6] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(2):383–386, 1990.
- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Adam Grabowski. Properties of the product of compact topological spaces. *Formalized Mathematics*, 8(1):55–59, 1999.
- [9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [10] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(5):665–674, 1991.
- [11] Zbigniew Karno. The lattice of domains of an extremally disconnected space. *Formalized Mathematics*, 3(2):143–149, 1992.
- [12] Artur Korniłowicz and Yasunari Shidama. Intersections of intervals and balls in  $\mathcal{E}_T^n$ . *Formalized Mathematics*, 12(3):301–306, 2004.
- [13] John M. Lee. *Introduction to Topological Manifolds*. Springer-Verlag, New York Berlin Heidelberg, 2000.
- [14] Robert Milewski. Bases of continuous lattices. *Formalized Mathematics*, 7(2):285–294, 1998.
- [15] Beata Padlewska. Locally connected spaces. *Formalized Mathematics*, 2(1):93–96, 1991.
- [16] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [17] Karol Pał. Basic properties of metrizable topological spaces. *Formalized Mathematics*, 17(3):201–205, 2009, doi: 10.2478/v10037-009-0024-8.
- [18] Bartłomiej Skorulski. First-countable, sequential, and Frechet spaces. *Formalized Mathematics*, 7(1):81–86, 1998.
- [19] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [20] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received August 17, 2010

---