

# Euler's Partition Theorem<sup>1</sup>

Karol Pał  
Institute of Informatics  
University of Białystok  
Ciołkowskiego 1M, 15-245 Białystok  
Poland

**Summary.** In this article we prove the Euler's Partition Theorem which states that the number of integer partitions with odd parts equals the number of partitions with distinct parts. The formalization follows H.S. Wilf's lecture notes [28] (see also [1]).

Euler's Partition Theorem is listed as item #45 from the "Formalizing 100 Theorems" list maintained by Freek Wiedijk at <http://www.cs.ru.nl/F.Wiedijk/100/> [27].

MSC: 05A17 03B35

Keywords: partition theorem

MML identifier: EULRPART, version: 8.1.04 5.32.1237

The notation and terminology used in this paper have been introduced in the following articles: [22], [2], [3], [17], [7], [16], [19], [14], [15], [23], [9], [10], [24], [5], [18], [6], [11], [29], [12], [26], and [13].

## 1. PRELIMINARIES

From now on  $x, y$  denote objects and  $i, j, k, m, n$  denote natural numbers.

Let  $r$  be an extended real number. One can verify that  $\langle r \rangle$  is extended real-valued and  $\langle r \rangle$  is decreasing, increasing, non-decreasing, and non-increasing.

Now we state the proposition:

---

<sup>1</sup>This work has been financed by the resources of the Polish National Science Centre granted by decision no. DEC-2012/07/N/ST6/02147.

- (1) Let us consider non-decreasing, extended real-valued finite sequences  $f, g$ . If  $f(\text{len } f) \leq g(1)$ , then  $f \wedge g$  is non-decreasing.

PROOF: Set  $f_3 = f \wedge g$ . For every extended reals  $e_1, e_2$  such that  $e_1, e_2 \in \text{dom } f_3$  and  $e_1 \leq e_2$  holds  $f_3(e_1) \leq f_3(e_2)$  by [7, (25)], [25, (25)].  $\square$

Let  $R$  be a binary relation. We say that  $R$  is odd-valued if and only if

(Def. 1)  $\text{rng } R \subseteq \mathbb{N}_{\text{odd}}$ .

- (2)  $n \in \mathbb{N}_{\text{odd}}$  if and only if  $n$  is odd.

Let us note that every binary relation which is odd-valued is also non-zero and natural-valued.

Let  $F$  be a function. Observe that  $F$  is odd-valued if and only if the condition

(Def. 2) is satisfied.

(Def. 2) for every  $x$  such that  $x \in \text{dom } F$  holds  $F(x)$  is an odd natural number.

One can check that every binary relation which is empty is also odd-valued.

Let  $i$  be an odd natural number. Let us observe that  $\langle i \rangle$  is odd-valued.

Let  $f, g$  be odd-valued finite sequences. Note that  $f \wedge g$  is odd-valued and every binary relation which is  $\mathbb{N}_{\text{odd}}$ -valued is also odd-valued.

Let  $n$  be a natural number. A partition of  $n$  is a non-zero, non-decreasing, natural-valued finite sequence and is defined by

(Def. 3)  $\sum it = n$ .

Now we state the proposition:

- (3)  $\emptyset$  is a partition of 0.

Let  $n$  be a natural number. Observe that there exists a partition of  $n$  which is odd-valued and there exists a partition of  $n$  which is one-to-one.

Let us observe that sethood property holds for partitions of  $n$ .

Let  $f$  be an odd-valued finite sequence.

An odd organization of  $f$  is a valued reorganization of  $f$  and is defined by

(Def. 4)  $2 \cdot n - 1 = f(it_{n,1})$  and ... and  $2 \cdot n - 1 = f(it_{n,\text{len}(it(n))})$ .

- (4) Let us consider an odd-valued finite sequence  $f$ , and a double reorganization  $o$  of  $\text{dom } f$ . Suppose for every  $n$ ,  $2 \cdot n - 1 = f(o_{n,1})$  and ... and  $2 \cdot n - 1 = f(o_{n,\text{len}(o(n))})$ . Then  $o$  is an odd organization of  $f$ .

PROOF: For every  $n$ , there exists  $x$  such that  $x = f(o_{n,1})$  and ... and  $x = f(o_{n,\text{len}(o(n))})$ . For every natural numbers  $n_1, n_2, i_1, i_2$  such that  $i_1 \in \text{dom}(o(n_1))$  and  $i_2 \in \text{dom}(o(n_2))$  and  $f(o_{n_1,i_1}) = f(o_{n_2,i_2})$  holds  $n_1 = n_2$  by [25, (25)].  $\square$

- (5) Let us consider an odd-valued finite sequence  $f$ , a complex-valued finite sequence  $g$ , and double reorganizations  $o_1, o_2$  of  $\text{dom } g$ . Suppose  $o_1$  is an odd organization of  $f$  and  $o_2$  is an odd organization of  $f$ . Then  $(\sum(g \odot o_1))(i) = (\sum(g \odot o_2))(i)$ .

PROOF: For every double reorganizations  $o_1, o_2$  of  $\text{dom } g$  such that  $o_1$  is an odd organization of  $f$  and  $o_2$  is an odd organization of  $f$  holds  $\text{rng}((f \odot o_1)(n)) \subseteq \text{rng}((f \odot o_2)(n))$  by [19, (49), (1)], [25, (29), (25)].  $\square$

- (6) Let us consider a partition  $p$  of  $n$ . Then there exists an odd-valued finite sequence  $O$  and there exists a natural-valued finite sequence  $a$  such that  $\text{len } O = \text{len } p = \text{len } a$  and  $p = O \cdot 2^a$  and  $p(1) = O(1) \cdot 2^{a(1)}$  and ... and  $p(\text{len } p) = O(\text{len } p) \cdot 2^{a(\text{len } p)}$ .

PROOF: Define  $\mathcal{P}[\text{object, object}] \equiv$  for every  $i$  and  $j$  such that  $p(\$_1) = 2^i \cdot (2 \cdot j + 1)$  holds  $\$2 = \langle 2 \cdot j + 1, i \rangle$ . For every  $k$  such that  $k \in \text{Seg len } p$  there exists  $x$  such that  $\mathcal{P}[k, x]$  by [20, (1)], [4, (4)]. Consider  $O_3$  being a finite sequence such that  $\text{dom } O_3 = \text{Seg len } p$  and for every  $k$  such that  $k \in \text{Seg len } p$  holds  $\mathcal{P}[k, O_3(k)]$  from [7, Sch. 1]. Define  $\mathcal{Q}(\text{object}) = O_3(\$1)_1$ . Consider  $O$  being a finite sequence such that  $\text{len } O = \text{len } p$  and for every  $k$  such that  $k \in \text{dom } O$  holds  $O(k) = \mathcal{Q}(k)$  from [7, Sch. 2]. For every  $x$  such that  $x \in \text{dom } O$  holds  $O(x)$  is an odd natural number by [20, (1)]. Define  $\mathcal{T}(\text{object}) = O_3(\$1)_2$ . Consider  $A$  being a finite sequence such that  $\text{len } A = \text{len } p$  and for every  $k$  such that  $k \in \text{dom } A$  holds  $A(k) = \mathcal{T}(k)$  from [7, Sch. 2]. For every  $x$  such that  $x \in \text{dom } A$  holds  $A(x)$  is natural by [20, (1)]. Set  $O_2 = O \cdot 2^A$ .  $p(1) = O(1) \cdot 2^{A(1)}$  and ... and  $p(\text{len } p) = O(\text{len } p) \cdot 2^{A(\text{len } p)}$  by [25, (25)], [20, (1)]. For every  $i$  such that  $i \in \text{dom } p$  holds  $p(i) = O_2(i)$  by [25, (25)].  $\square$

- (7) Let us consider a finite set  $D$ , and a function  $f$  from  $D$  into  $\mathbb{N}$ . Then there exists a finite sequence  $K$  of elements of  $D$  such that for every element  $d$  of  $D$ ,  $\overline{\text{Coim}(K, d)} = f(d)$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every finite set  $D$  such that  $\overline{D} = \$1$  for every function  $f$  from  $D$  into  $\mathbb{N}$ , there exists a finite sequence  $K$  of elements of  $D$  such that for every element  $d$  of  $D$ ,  $\overline{\text{Coim}(K, d)} = f(d)$ .  $\mathcal{P}[0]$ . If  $\mathcal{P}[i]$ , then  $\mathcal{P}[i + 1]$  by [21, (55)], [8, (63)], [25, (57)], [13, (56)].  $\mathcal{P}[i]$  from [5, Sch. 2].  $\square$

- (8) Let us consider complex-valued finite sequences  $f_1, f_2, g_1, g_2$ . Suppose  $\text{len } f_1 = \text{len } g_1$ . Then  $(f_1 \wedge f_2) \cdot (g_1 \wedge g_2) = (f_1 \cdot g_1) \wedge (f_2 \cdot g_2)$ .

- (9) Let us consider natural-valued finite sequences  $f, K$ . Suppose for every  $i$ ,  $\overline{\text{Coim}(K, i)} = f(i)$ . Then  $\sum K = 1 \cdot f(1) + 2 \cdot f(2) + ((\text{id}_{\text{dom } f} \cdot f), 3) + \dots$

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every natural-valued finite sequences  $f, K$  such that  $\text{len } f = \$1$  and for every  $i$ ,  $\overline{\text{Coim}(K, i)} = f(i)$  holds  $\sum K = ((\text{id}_{\text{dom } f} \cdot f), 1) + \dots$   $\mathcal{P}[0]$  by [25, (25)], [9, (72)], [19, (20), (22)]. If  $\mathcal{P}[i]$ , then  $\mathcal{P}[i + 1]$  by [25, (55)], [5, (13)], [7, (59)], [8, (51)].  $\mathcal{P}[i]$  from [5, Sch. 2].  $\square$

- (10) Let us consider a natural-valued finite sequence  $g$ , and a double reorgani-

zation  $s_1$  of  $\text{dom } g$ . Then there exists a  $(2 \cdot \text{len } s_1)$ -element finite sequence  $K$  of elements of  $\mathbb{N}$  such that for every  $j$ ,  $K(2 \cdot j) = 0$  and  $K(2 \cdot j - 1) = g(s_{1j,1}) + ((g \odot s_1)(j), 2) + \dots$ . PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  if  $\$1 = 2 \cdot j - 1$ , then  $\$2 = g(s_{1j,1}) + ((g \odot s_1)(j), 2) + \dots$  and if  $\$1 = 2 \cdot j$ , then  $\$2 = 0$ . Set  $S = \text{Seg}(2 \cdot \text{len } s_1)$ . For every  $k$  such that  $k \in S$  there exists  $x$  such that  $\mathcal{P}[k, x]$  by [22, (9)]. Consider  $f$  being a finite sequence such that  $\text{dom } f = S$  and for every  $i$  such that  $i \in S$  holds  $\mathcal{P}[i, f(i)]$  from [7, Sch. 1].  $\text{rng } f \subseteq \mathbb{N}$  by [22, (9)].  $f(2 \cdot i) = 0$ .  $f(2 \cdot i - 1) = g(s_{1i,1}) + ((g \odot s_1)(i), 2) + \dots$  by [25, (25)], [5, (13)], [19, (15)].  $\square$

## 2. EULER TRANSFORMATION

Now we state the proposition:

- (11) Let us consider a one-to-one partition  $d$  of  $n$ . Then there exists an odd-valued partition  $e$  of  $n$  such that for every natural number  $j$  for every odd-valued finite sequence  $O_1$  for every natural-valued finite sequence  $a_1$  such that  $\text{len } O_1 = \text{len } d = \text{len } a_1$  and  $d = O_1 \cdot 2^{a_1}$  for every double reorganization  $s_1$  of  $\text{dom } d$  such that  $1 = O_1(s_{11,1})$  and ... and  $1 = O_1(s_{11,\text{len}(s_1(1))})$  and  $3 = O_1(s_{12,1})$  and ... and  $3 = O_1(s_{12,\text{len}(s_1(2))})$  and  $5 = O_1(s_{13,1})$  and ... and  $5 = O_1(s_{13,\text{len}(s_1(3))})$  and for every  $i$ ,  $2 \cdot i - 1 = O_1(s_{1i,1})$  and ... and  $2 \cdot i - 1 = O_1(s_{1i,\text{len}(s_1(i))})$  holds  $\overline{\text{Coim}(e, 1)} = 2^{a_1}(s_{11,1}) + ((2^{a_1} \odot s_1)(1), 2) + \dots$  and  $\overline{\text{Coim}(e, 3)} = 2^{a_1}(s_{12,1}) + ((2^{a_1} \odot s_1)(2), 2) + \dots$  and  $\overline{\text{Coim}(e, 5)} = 2^{a_1}(s_{13,1}) + ((2^{a_1} \odot s_1)(3), 2) + \dots$  and  $\overline{\text{Coim}(e, j \cdot 2 - 1)} = 2^{a_1}(s_{1j,1}) + ((2^{a_1} \odot s_1)(j), 2) + \dots$

PROOF: Consider  $O$  being an odd-valued finite sequence,  $a$  being a natural-valued finite sequence such that  $\text{len } O = \text{len } d = \text{len } a$  and  $d = O \cdot 2^a$  and  $d(1) = O(1) \cdot 2^{a(1)}$  and ... and  $d(\text{len } d) = O(\text{len } d) \cdot 2^{a(\text{len } d)}$ .  $n = d(1) + ((d, 2) + \dots + (d, \text{len } d))$  by [19, (22)].  $n = 2^{a(1)} \cdot O(1) + 2^{a(2)} \cdot O(2) + ((O \cdot 2^a, 3) + \dots + (O \cdot 2^a, \text{len } d))$  by [19, (20)], [25, (25)]. Reconsider  $s_1 =$  the odd organization of  $O$  as a double reorganization of  $\text{dom } 2^a$ . Consider  $\mu$  being a  $(2 \cdot \text{len } s_1)$ -element finite sequence of elements of  $\mathbb{N}$  such that for every  $j$ ,  $\mu(2 \cdot j) = 0$  and  $\mu(2 \cdot j - 1) = 2^a(s_{1j,1}) + ((2^a \odot s_1)(j), 2) + \dots$ . Set  $\alpha = a \cdot s_1(1)$ . Set  $\beta = a \cdot s_1(2)$ . Set  $\gamma = a \cdot s_1(3)$ .  $n = (2^\alpha(1) + (2^\alpha, 2) + \dots) \cdot 1 + (2^\beta(1) + (2^\beta, 2) + \dots) \cdot 3 + (2^\gamma(1) + (2^\gamma, 2) + \dots) \cdot 5 + ((\text{id}_{\text{dom } \mu} \cdot \mu), 7) + \dots$  by [25, (29)], [19, (41)], [25, (25)], [9, (12)].  $n = \mu(1) \cdot 1 + \mu(3) \cdot 3 + \mu(5) \cdot 5 + ((\text{id}_{\text{dom } \mu} \cdot \mu), 7) + \dots$  by [19, (42), (41), (25)]. Consider  $K$  being an odd-valued finite sequence such that  $K$  is non-decreasing and for every  $i$ ,  $\overline{\text{Coim}(K, i)} = \mu(i)$ .  $n = \overline{\text{Coim}(K, 1)} \cdot 1 + \overline{\text{Coim}(K, 3)} \cdot 3 + \overline{\text{Coim}(K, 5)} \cdot 5 + ((\text{id}_{\text{dom } \mu} \cdot \mu), 7) + \dots$ .  $n = \sum K$  by [19, (20)], (9). For every  $j$  such

that  $1 \leq j \leq \text{len } d$  holds  $O(j) = O_1(j)$  and  $a(j) = a_1(j)$  by [25, (25)], [22, (9)], [4, (4)]. For every  $j$ ,  $\overline{\text{Coim}(K, j \cdot 2 - 1)} = 2^{a_1}(\text{sort}1_{j,1}) + ((2^{a_1} \odot \text{sort}1)(j), 2) + \dots$  by [19, (42)], [25, (29)], [9, (72)], [19, (22)].  $\square$

Let  $n$  be a natural number and  $p$  be a one-to-one partition of  $n$ . The Euler transformation  $p$  yielding an odd-valued partition of  $n$  is defined by

(Def. 5) for every odd-valued finite sequence  $O$  and for every natural-valued finite sequence  $a$  such that  $\text{len } O = \text{len } p = \text{len } a$  and  $p = O \cdot 2^a$  for every double reorganization  $s_1$  of  $\text{dom } p$  such that  $1 = O(s_{11,1})$  and ... and  $1 = O(s_{11, \text{len}(s_1(1))})$  and  $3 = O(s_{12,1})$  and ... and  $3 = O(s_{12, \text{len}(s_1(2))})$  and  $5 = O(s_{13,1})$  and ... and  $5 = O(s_{13, \text{len}(s_1(3))})$  and for every  $i$ ,  $2 \cdot i - 1 = O(s_{1i,1})$  and ... and  $2 \cdot i - 1 = O(s_{1i, \text{len}(s_1(i))})$  holds  $\overline{\text{Coim}(it, 1)} = 2^a(s_{11,1}) + ((2^a \odot s_1)(1), 2) + \dots$  and  $\overline{\text{Coim}(it, 3)} = 2^a(s_{12,1}) + ((2^a \odot s_1)(2), 2) + \dots$  and  $\overline{\text{Coim}(it, 5)} = 2^a(s_{13,1}) + ((2^a \odot s_1)(3), 2) + \dots$  and  $\overline{\text{Coim}(it, j \cdot 2 - 1)} = 2^a(s_{1j,1}) + ((2^a \odot s_1)(j), 2) + \dots$

Now we state the proposition:

(12) Let us consider a natural number  $n$ , a one-to-one partition  $p$  of  $n$ , and an odd-valued partition  $e$  of  $n$ . Then  $e =$  the Euler transformation  $p$  if and only if for every odd-valued finite sequence  $O$  and for every natural-valued finite sequence  $a$  and for every odd organization  $s_1$  of  $O$  such that  $\text{len } O = \text{len } p = \text{len } a$  and  $p = O \cdot 2^a$  for every  $j$ ,  $\overline{\text{Coim}(e, j \cdot 2 - 1)} = ((2^a \odot s_1)(j), 1) + \dots$

PROOF: If  $e =$  the Euler transformation  $p$ , then for every odd-valued finite sequence  $O$  and for every natural-valued finite sequence  $a$  and for every odd organization  $s_1$  of  $O$  such that  $\text{len } O = \text{len } p = \text{len } a$  and  $p = O \cdot 2^a$  for every  $j$ ,  $\overline{\text{Coim}(e, j \cdot 2 - 1)} = ((2^a \odot s_1)(j), 1) + \dots$  by [25, (29)], [19, (42), (20)]. For every  $j$  and for every odd-valued finite sequence  $O$  and for every natural-valued finite sequence  $a$  such that  $\text{len } O = \text{len } p = \text{len } a$  and  $p = O \cdot 2^a$  for every double reorganization  $s_1$  of  $\text{dom } p$  such that  $1 = O(s_{11,1})$  and ... and  $1 = O(s_{11, \text{len}(s_1(1))})$  and  $3 = O(s_{12,1})$  and ... and  $3 = O(s_{12, \text{len}(s_1(2))})$  and  $5 = O(s_{13,1})$  and ... and  $5 = O(s_{13, \text{len}(s_1(3))})$  and for every  $i$ ,  $2 \cdot i - 1 = O(s_{1i,1})$  and ... and  $2 \cdot i - 1 = O(s_{1i, \text{len}(s_1(i))})$  holds  $\overline{\text{Coim}(e, 1)} = 2^a(s_{11,1}) + ((2^a \odot s_1)(1), 2) + \dots$  and  $\overline{\text{Coim}(e, 3)} = 2^a(s_{12,1}) + ((2^a \odot s_1)(2), 2) + \dots$  and  $\overline{\text{Coim}(e, 5)} = 2^a(s_{13,1}) + ((2^a \odot s_1)(3), 2) + \dots$  and  $\overline{\text{Coim}(e, j \cdot 2 - 1)} = 2^a(s_{1j,1}) + ((2^a \odot s_1)(j), 2) + \dots$  by [25, (29)], (4), [19, (42), (20)].  $\square$

One can verify that every real-valued function which is one-to-one and non-decreasing is also increasing.

- (13) Let us consider an odd-valued finite sequence  $O$ , a natural-valued finite sequence  $a$ , and an odd organization  $s$  of  $O$ . Suppose  $\text{len } O = \text{len } a$  and  $O \cdot 2^a$  is one-to-one. Then  $(a \odot s)(i)$  is one-to-one.

PROOF:  $(a \odot s)(i)$  is one-to-one by [9, (11), (12)], [25, (25)].  $\square$

- (14) Let us consider one-to-one partitions  $p_1, p_2$  of  $n$ . Suppose the Euler transformation  $p_1 =$  the Euler transformation  $p_2$ . Then  $p_1 = p_2$ .
- (15) Let us consider an odd-valued partition  $e$  of  $n$ . Then there exists a one-to-one partition  $p$  of  $n$  such that  $e =$  the Euler transformation  $p$ .

PROOF: Define  $\mathcal{K}(\text{object}) = \overline{\text{Coim}(e, \$_1)}$ . Consider  $H$  being a finite sequence such that  $\text{len } H = n$  and for every  $k$  such that  $k \in \text{dom } H$  holds  $H(k) = \mathcal{K}(k)$  from [7, Sch. 2].  $\text{rng } H \subseteq \mathbb{N}$ .  $\sum e = \sum(\text{idseq}(n) \cdot H)$  by [25, (25)], [5, (14)], [9, (72)], [30, (5)]. Define  $\mathcal{F}[\text{natural number, object}] \equiv$  there exists an increasing, natural-valued finite sequence  $f$  such that  $H(\$_1) = 2^f(1) + (2^f, 2) + \dots$  and  $\$_2 = \$_1 \cdot 2^f$ . There exists a finite sequence  $p$  of elements of  $\mathbb{N}^*$  such that  $\text{dom } p = \text{Seg len } H$  and for every  $k$  such that  $k \in \text{Seg len } H$  holds  $\mathcal{F}[k, p(k)]$  by [19, (31)]. Consider  $p$  being a finite sequence of elements of  $\mathbb{N}^*$  such that  $\text{dom } p = \text{Seg len } H$  and for every  $k$  such that  $k \in \text{Seg len } H$  holds  $\mathcal{F}[k, p(k)]$ . For every  $k$  such that  $p(k) \neq \emptyset$  holds  $k$  is odd by [18, (83)], [12, (85)], [19, (22)], [9, (72)]. Set  $N =$  the concatenation of  $\mathbb{N}$ . Set  $n_3 = N \odot p$ . Set  $s_2 = \text{sort}_a n_3$ .  $s_2$  is a one-to-one partition of  $n$  by [19, (1)], [25, (25)], [12, (45)], [18, (83)]. For every odd-valued finite sequence  $O$  and for every natural-valued finite sequence  $a$  and for every odd organization  $s_1$  of  $O$  such that  $\text{len } O = \text{len } s_2 = \text{len } a$  and  $s_2 = O \cdot 2^a$  for every  $j$ ,  $\overline{\text{Coim}(e, j \cdot 2 - 1)} = ((2^a \odot s_1)(j), 1) + \dots$  by [25, (29)], [5, (14)], [9, (72)], [25, (25)].  $\square$

### 3. MAIN THEOREM

Now we state the proposition:

- (16) EULER'S PARTITION THEOREM:

the set of all  $p$  where  $p$  is an odd-valued partition of  $n =$

the set of all  $p$  where  $p$  is a one-to-one partition of  $n$ . The theorem is a consequence of (15) and (14).

### REFERENCES

- [1] George E. Andrews and Kimmo Eriksson. *Integer Partitions*. ISBN 9780521600903.  
 [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.  
 [3] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(3):589–593, 1990.

- [4] Grzegorz Bancerek. Countable sets and Hessenberg's theorem. *Formalized Mathematics*, 2(1):65–69, 1991.
- [5] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [6] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [7] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [8] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [11] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [12] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [13] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [14] Marco B. Caminati. Preliminaries to classical first order model theory. *Formalized Mathematics*, 19(3):155–167, 2011. doi:10.2478/v10037-011-0025-2.
- [15] Marco B. Caminati. First order languages: Further syntax and semantics. *Formalized Mathematics*, 19(3):179–192, 2011. doi:10.2478/v10037-011-0027-0.
- [16] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [17] Magdalena Jastrzębska and Adam Grabowski. Some properties of Fibonacci numbers. *Formalized Mathematics*, 12(3):307–313, 2004.
- [18] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [19] Karol Pąk. Flexary operations. *Formalized Mathematics*, 23(2):81–92, 2015. doi:10.1515/forma-2015-0008.
- [20] Karol Pąk. The Nagata-Smirnov theorem. Part II. *Formalized Mathematics*, 12(3):385–389, 2004.
- [21] Karol Pąk. Stirling numbers of the second kind. *Formalized Mathematics*, 13(2):337–345, 2005.
- [22] Piotr Rudnicki and Andrzej Trybulec. Abian's fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [23] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [24] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [25] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [26] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [27] Freek Wiedijk. Formalizing 100 theorems.
- [28] Herbert S. Wilf. Lectures on integer partitions.
- [29] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [30] Bo Zhang and Yatsuka Nakamura. The definition of finite sequences and matrices of probability, and addition of matrices of real elements. *Formalized Mathematics*, 14(3):101–108, 2006. doi:10.2478/v10037-006-0012-1.

*Received March 26, 2015*

---