

# Leibniz Series for $\pi^1$

Karol Pałk  
Institute of Informatics  
University of Białystok  
Ciołkowskiego 1M, 15-245 Białystok  
Poland

**Summary.** In this article we prove the Leibniz series for  $\pi$  which states that

$$\frac{\pi}{4} = \sum_{n=0}^{\infty} \frac{(-1)^n}{2 \cdot n + 1}.$$

The formalization follows K. Knopp [8], [1] and [6]. *Leibniz's Series for Pi* is item #26 from the “Formalizing 100 Theorems” list maintained by Freek Wiedijk at <http://www.cs.ru.nl/F.Wiedijk/100/>.

MSC: 40G99 03B35

Keywords:  $\pi$  approximation; Leibniz theorem; Leibniz series

MML identifier: LEIBNIZ1, version: 8.1.05 5.39.1282

## 1. PRELIMINARIES

From now on  $i, n, m$  denote natural numbers,  $r, s$  denote real numbers, and  $A$  denotes a non empty, closed interval subset of  $\mathbb{R}$ .

Now we state the proposition:

(1)  $\text{rng}(\text{the function } \tan \upharpoonright ]-\frac{\pi}{2}, \frac{\pi}{2}[) = \mathbb{R}$ .

PROOF: Set  $P = \frac{\pi}{2}$ . Set  $I = ]-P, P[$ .  $\mathbb{R} \subseteq \text{rng}(\text{the function } \tan \upharpoonright I)$  by [4, (50)], [20, (30)], [14, (15)], [16, (1)].  $\square$

---

<sup>1</sup>This work has been financed by the resources of the Polish National Science Centre granted by decision no. DEC-2012/07/N/ST6/02147.

One can verify that the function arctan is total and the function arctan is differentiable.

Now we state the propositions:

(2) (The function arctan)'( $r$ ) =  $\frac{1}{1+r^2}$ .

(3) Let us consider an open subset  $Z$  of  $\mathbb{R}$ . Then

(i) the function arctan is differentiable on  $Z$ , and

(ii) for every  $r$  such that  $r \in Z$  holds (the function arctan)' $_{|Z}(r) = \frac{1}{1+r^2}$ .

The theorem is a consequence of (2).

Let us consider  $n$ . One can verify that  $\square^n$  is continuous.

Now we state the propositions:

(4) (i)  $\text{dom}(\frac{\square^n}{\square^0 + \square^2}) = \mathbb{R}$ , and

(ii)  $\frac{\square^n}{\square^0 + \square^2}$  is continuous, and

(iii)  $(\frac{\square^n}{\square^0 + \square^2})(r) = \frac{r^n}{1+r^2}$ .

(5)  $\int_A (\frac{\square^0}{\square^0 + \square^2})(x) dx =$

(the function arctan)(sup  $A$ ) – (the function arctan)(inf  $A$ ).

PROOF: Set  $Z_0 = \square^0$ . Set  $Z_2 = \square^2$ . Set  $f = \frac{Z_0}{Z_0 + Z_2}$ .  $\text{dom } f = \mathbb{R}$ .  $f$  is continuous. If  $r \in \mathbb{R}$ , then  $f(r) = \frac{1}{1+r^2}$  by [13, (4)], (4). For every element  $x$  of  $\mathbb{R}$  such that  $x \in \text{dom}(\text{the function arctan})'_{|\mathbb{R}}$  holds (the function arctan)' $_{|\mathbb{R}}(x) = f(x)$ .  $\square$

(6)  $\int_A ((-1)^i \cdot (\frac{\square^{2 \cdot n}}{\square^0 + \square^2}))(x) dx = (-1)^i \cdot (\frac{1}{2 \cdot n + 1}) \cdot (\text{sup } A)^{2 \cdot n + 1} - (\frac{1}{2 \cdot n + 1}) \cdot (\text{inf } A)^{2 \cdot n + 1} + \int_A ((-1)^{i+1} \cdot (\frac{\square^{2 \cdot (n+1)}}{\square^0 + \square^2}))(x) dx.$

PROOF: Set  $I_1 = (-1)^i$ . Set  $i_1 = i + 1$ . Set  $n_1 = n + 1$ . Set  $I_2 = (-1)^{i_1}$ . Set  $Z_0 = \square^0$ . Set  $Z_2 = \square^2$ . Set  $Z_{2n} = \square^{2 \cdot n}$ . Set  $f = I_1 \cdot Z_{2n}$ . Set  $g = I_2 \cdot (\frac{\square^{2 \cdot n_1}}{Z_0 + Z_2})$ .  $\text{dom } g = \mathbb{R}$ . For every element  $x$  of  $\mathbb{R}$ ,  $(I_1 \cdot (\frac{Z_{2n}}{Z_0 + Z_2}))(x) = (f + g)(x)$  by [13, (6)], [17, (36)], (4).  $f + g = I_1 \cdot (\frac{Z_{2n}}{Z_0 + Z_2}) \cdot \frac{\square^{2 \cdot n_1}}{Z_0 + Z_2}$  is continuous.  $\square$

(7) Suppose  $A = [0, r]$  and  $r \geq 0$ . Then  $|\int_A (\frac{\square^{2 \cdot n}}{\square^0 + \square^2})(x) dx| \leq (\frac{1}{2 \cdot n + 1}) \cdot r^{2 \cdot n + 1}.$

PROOF: Set  $Z_0 = \square^0$ . Set  $Z_2 = \square^2$ . Set  $N = 2 \cdot n$ . Set  $Z_n = \square^N$ . Set  $f = \frac{Z_n}{Z_0 + Z_2}$ .  $f$  is continuous and  $\text{dom } f = \mathbb{R}$ . Reconsider  $f_1 = f \upharpoonright A$  as a function from  $A$  into  $\mathbb{R}$ . Reconsider  $Z_1 = Z_n \upharpoonright A$  as a function from  $A$  into  $\mathbb{R}$ . For every  $r$  such that  $r \in A$  holds  $f_1(r) \leq Z_1(r)$  by [4, (49)], [17,

(36)], [18, (3)], (4). For every object  $x$  such that  $x \in \mathbb{R}$  holds  $f(x) = |f|(x)$  by [13, (8)], (4).  $\square$

## 2. EULER TRANSFORMATION

Let  $a$  be a sequence of real numbers. The alternating series of  $a$  yielding a sequence of real numbers is defined by

(Def. 1)  $it(i) = (-1)^i \cdot a(i)$ .

Now we state the proposition:

- (8) Let us consider a sequence  $a$  of real numbers. Suppose  $a$  is non-negative yielding, non-increasing, and convergent and  $\lim a = 0$ . Then
- (i) the alternating series of  $a$  is summable, and
  - (ii) for every  $n$ ,  $(\sum_{\alpha=0}^{\kappa}(\text{the alternating series of } a)(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n) \geq \sum(\text{the alternating series of } a) \geq (\sum_{\alpha=0}^{\kappa}(\text{the alternating series of } a)(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n + 1)$ .

PROOF: Set  $A =$  the alternating series of  $a$ . Set  $P = (\sum_{\alpha=0}^{\kappa} A(\alpha))_{\kappa \in \mathbb{N}}$ . Define  $\mathcal{T}[\text{natural number, object}] \equiv \$_2 = P(2 \cdot \$_1)$ . Define  $\mathcal{S}[\text{natural number, object}] \equiv \$_2 = P(2 \cdot \$_1 + 1)$ . Consider  $T$  being a function from  $\mathbb{N}$  into  $\mathbb{R}$  such that for every element  $x$  of  $\mathbb{N}$ ,  $\mathcal{T}[x, T(x)]$  from [5, Sch. 3].

Consider  $S$  being a function from  $\mathbb{N}$  into  $\mathbb{R}$  such that for every element  $x$  of  $\mathbb{N}$ ,  $\mathcal{S}[x, S(x)]$  from [5, Sch. 3]. For every natural number  $n$ ,  $S(n) \leq S(n+1)$ . For every natural number  $n$ ,  $T(n) \geq T(n+1)$ . For every natural number  $n$ ,  $T(n) \geq S(n)$ . For every natural number  $n$ ,  $T(n) > S(0) - 1$  by [10, (6)]. For every natural number  $n$ ,  $S(n) < T(0) + 1$  by [10, (8)].

Define  $\mathcal{D}(\text{natural number}) = 2 \cdot \$_1 + 1$ . Consider  $D$  being a function from  $\mathbb{N}$  into  $\mathbb{N}$  such that for every element  $x$  of  $\mathbb{N}$ ,  $\mathcal{D}(x) = D(x)$  from [5, Sch. 8]. Reconsider  $D_1 = D$  as a many sorted set indexed by  $\mathbb{N}$ . For every natural number  $n$ ,  $D(n) < D(n+1)$  by [2, (13)]. Reconsider  $a_2 = a \cdot D_1$  as a sequence of real numbers.

For every object  $x$  such that  $x \in \mathbb{N}$  holds  $a_2(x) = (T - S)(x)$  by [4, (12)]. For every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|P(m) - \lim T| < p$  by [19, (9)].  $\square$

3. MAIN THEOREM

Let us consider  $r$ . The Leibniz series of  $r$  yielding a sequence of real numbers is defined by

(Def. 2)  $it(n) = \frac{(-1)^n \cdot r^{2 \cdot n + 1}}{2 \cdot n + 1}$ .

The Leibniz series yielding a sequence of real numbers is defined by the term

(Def. 3) the Leibniz series of 1.

Now we state the propositions:

(9) Suppose  $r \in [-1, 1]$ . Then

(i) |the Leibniz series of  $r$ | is non-negative yielding, non-increasing, and convergent, and

(ii)  $\lim$ |the Leibniz series of  $r$ | = 0.

PROOF: Set  $r_1 =$  the Leibniz series of  $r$ . Set  $A = |r_1|$ .  $A(n) = \frac{|r|^{2 \cdot n + 1}}{2 \cdot n + 1}$  by [15, (1)], [3, (67), (65)].  $A(n) \geq A(n + 1)$  by [3, (46)], [15, (1)], [13, (6)], [2, (13)]. Set  $C = \{0\}_{n \in \mathbb{N}}$ . Define  $\mathcal{F}$ (natural number) =  $\frac{1}{\mathbb{S}_1 + \frac{1}{2}}$ . Consider  $f$  being a sequence of real numbers such that  $f(n) = \mathcal{F}(n)$  from [11, Sch. 1].  $C(n) \leq A(n) \leq f(n)$  by [11, (57)], [3, (46)], [13, (11)], [2, (11)].  $\square$

(10) (i) if  $r \geq 0$ , then the alternating series of |the Leibniz series of  $r$ | = the Leibniz series of  $r$ , and

(ii) if  $r < 0$ , then  $(-1) \cdot$  (the alternating series of |the Leibniz series of  $r$ |) = the Leibniz series of  $r$ .

PROOF: Set  $r_1 =$  the Leibniz series of  $r$ . Set  $A = |r_1|$ . Set  $a_1 =$  the alternating series of  $A$ .  $a_1(n) = (-1)^n \cdot (\frac{|r|^{2 \cdot n + 1}}{2 \cdot n + 1})$  by [15, (1)], [3, (67), (65)]. If  $r \geq 0$ , then  $a_1 = r_1$ .  $\square$

(11) If  $r \in [-1, 1]$ , then the Leibniz series of  $r$  is summable. The theorem is a consequence of (9), (8), and (10).

(12) Suppose  $A = [0, r]$  and  $r \geq 0$ . Then (the function arctan)( $r$ ) =  $(\sum_{\alpha=0}^{\kappa}$ (the Leibniz series of  $r$ )( $\alpha$ )) $_{\kappa \in \mathbb{N}}$ ( $n$ ) +  $\int_A ((-1)^{n+1} \cdot (\frac{\square^{2 \cdot (n+1)}}{\square^0 + \square^2}))(x)dx$ .

PROOF: Set  $Z_0 = \square^0$ . Set  $Z_2 = \square^2$ . Set  $r_1 =$  the Leibniz series of  $r$ . Define  $\mathcal{P}$ [natural number]  $\equiv$  (the function arctan)( $r$ ) =  $(\sum_{\alpha=0}^{\kappa} r_1(\alpha))_{\kappa \in \mathbb{N}}$ ( $\mathbb{S}_1$ ) +  $\int_A ((-1)^{\mathbb{S}_1 + 1} \cdot (\frac{\square^{2 \cdot (\mathbb{S}_1 + 1)}}{Z_0 + Z_2}))(x)dx$ .  $\mathcal{P}[0]$  by (5), [14, (43)], [13, (4)], [9, (21)].

If  $\mathcal{P}[i]$ , then  $\mathcal{P}[i + 1]$  by [13, (11)], [2, (11)], (6).  $\mathcal{P}[i]$  from [2, Sch. 2].  $\square$

(13) If  $0 \leq r \leq 1$ , then (the function arctan)( $r$ ) =  $\sum$ (the Leibniz series of  $r$ ).

PROOF: Set  $r_1 =$  the Leibniz series of  $r$ . Set  $P = (\sum_{\alpha=0}^{\kappa} r_1(\alpha))_{\kappa \in \mathbb{N}}$ . Set  $A =$  (the function  $\arctan$ )( $r$ ). Define  $\mathcal{I}$ (natural number)  $= \frac{\square^{2 \cdot s_1}}{\square^0 + \square^2}$ .  $P$  is convergent. For every  $s$  such that  $0 < s$  there exists  $n$  such that for every  $m$  such that  $n \leq m$  holds  $|P(m) - A| < s$  by [12, (3)], (4), [7, (11), (10)].  
□

(14) LEIBNIZ SERIES FOR  $\pi$ :

$$\frac{\pi}{4} = \sum(\text{the Leibniz series}).$$

(15)  $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n + 1) \leq \sum(\text{the Leibniz series}) \leq (\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n)$ . The theorem is a consequence of (9), (10), and (8).

(16) (i)  $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(1) = \frac{2}{3}$ , and

(ii) if  $n$  is odd, then  $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(n+2) = (\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(n) + \frac{2}{4 \cdot n^2 + 16 \cdot n + 15}$ .

(17)  $\pi$  APPROXIMATION:

$$\frac{313}{100} < \pi < \frac{315}{100}. \text{ The theorem is a consequence of (16), (14), and (15).}$$

## REFERENCES

- [1] George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*. Cambridge University Press, 1999.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Lokenath Debnath. *The Legacy of Leonhard Euler: A Tricentennial Tribute*. World Scientific, 2010.
- [7] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and integrability for continuous functions. *Formalized Mathematics*, 9(2):281–284, 2001.
- [8] Konrad Knopp. *Infinite Sequences and Series*. Dover Publications, 1956. ISBN 978-0-486-60153-3.
- [9] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(4):703–709, 1990.
- [10] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [11] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [12] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [13] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [14] Xiquan Liang and Bing Xie. Inverse trigonometric functions  $\arctan$  and  $\text{arccot}$ . *Formalized Mathematics*, 16(2):147–158, 2008. doi:10.2478/v10037-008-0021-3.
- [15] Akira Nishino and Yasunari Shidama. The Maclaurin expansions. *Formalized Mathematics*, 13(3):421–425, 2005.

- [16] Chanapat Pacharapokin, Kanchun, and Hiroshi Yamazaki. Formulas and identities of trigonometric functions. *Formalized Mathematics*, 12(2):139–141, 2004.
- [17] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(1):125–130, 1991.
- [18] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(2):213–216, 1991.
- [19] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [20] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(2):255–263, 1998.

*Received October 18, 2016*

---



The English version of this volume of *Formalized Mathematics* was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.