

Kleene Algebra of Partial Predicates

Artur Kornilowicz
Institute of Informatics
University of Białystok
Poland

Ievgen Ivanov
Taras Shevchenko National University
Kyiv, Ukraine

Mykola Nikitchenko
Taras Shevchenko National University
Kyiv, Ukraine

Summary. We show that the set of all partial predicates over a set D together with the disjunction, conjunction, and negation operations (denoted in the paper as “ or_{PP} ”, “ and_{PP} ”, and “ not_{PP} ”), defined in accordance with the truth tables of S.C. Kleene’s strong logic of indeterminacy [17], forms a Kleene algebra. A Kleene algebra is a De Morgan algebra [3] (also called quasi-Boolean algebra) which satisfies the condition $x \wedge \neg x \leq y \vee \neg y$ (sometimes called the normality axiom). We use the formalization of De Morgan algebras from [8].

The term “Kleene algebra” was introduced by A. Monteiro and D. Brignole in [3]. A similar notion of a “normal i-lattice” had been previously studied by J.A. Kalman [16]. More details about the origin of this notion and its relation to other notions can be found in [24, 4, 1, 2]. It should be noted that there is a different widely known class of algebras, also called Kleene algebras [22, 6], which generalize the algebra of regular expressions, however, the term “Kleene algebra” used in this paper does not refer to them.

Algebras of partial predicates naturally arise in computability theory in the study on partial recursive predicates. They were studied in connection with non-classical logics [17, 5, 18, 32, 29, 30]. A partial predicate also corresponds to the notion of a partial set [26] on a given domain, which represents a (partial) property which for any given element of this domain may hold, not hold, or neither hold nor not hold. The field of all partial sets on a given domain is an algebra with generalized operations of union, intersection, complement, and three constants (0, 1, n which is the fixed point of complement) which can be generalized to an equational class of algebras called DMF-algebras (De Morgan algebras with a single fixed point of involution) [25]. In [27] partial sets and DMF-algebras were considered as a basis for unification of set-theoretic and linguistic approaches to probability.

Partial predicates over classes of mathematical models of data were used for formalizing semantics of computer programs in the composition-nominative approach to program formalization [31, 28, 33, 15], for formalizing extensions of the Floyd-Hoare logic [7, 9] which allow reasoning about properties of programs in the case of partial pre- and postconditions [23, 20, 19, 21], for formalizing dynamical models with partial behaviors in the context of the mathematical systems theory [11, 13, 14, 12, 10].

MSC: 03B70 03G25 03B35

Keywords: partial predicate; Kleene algebra

MML identifier: PARTPR.1, version: 8.1.07 5.47.1318

1. PARTIAL PREDICATES

From now on x denotes an object and D denotes a set.

Let us consider D . The functor $\text{Pr}(D)$ yielding a set is defined by the term

(Def. 1) $D \dot{\rightarrow} \text{Boolean}$.

Observe that $\text{Pr}(D)$ is non empty and functional.

A partial predicate of D is a partial function from D to *Boolean*. From now on p denotes a partial predicate of D .

Now we state the propositions:

- (1) If $x \in \text{Pr}(D)$, then x is a partial predicate of D .
- (2) $p \in \text{Pr}(D)$.
- (3) (i) $p(x) = \text{true}$, or
(ii) $p(x) = \text{false}$.

Let us consider D . The functor $\text{PPneg}(D)$ yielding a function from $\text{Pr}(D)$ into $\text{Pr}(D)$ is defined by

(Def. 2) for every partial predicate p of D , $\text{dom}(it(p)) = \text{dom } p$ and for every object d , if $d \in \text{dom } p$ and $p(d) = \text{true}$, then $it(p)(d) = \text{false}$ and if $d \in \text{dom } p$ and $p(d) = \text{false}$, then $it(p)(d) = \text{true}$.

Let us consider p . The functor $\text{not}_{\text{PP}}(p)$ yielding a partial predicate of D is defined by the term

(Def. 3) $(\text{PPneg}(D))(p)$.

Note that the functor is involutive.

Now we state the propositions:

- (4) If $x \in \text{dom } p$ and $(\text{not}_{\text{PP}}(p))(x) = \text{false}$, then $p(x) = \text{true}$. The theorem is a consequence of (3).

- (5) If $x \in \text{dom } p$ and $(\text{not}_{\text{PP}}(p))(x) = \text{true}$, then $p(x) = \text{false}$. The theorem is a consequence of (3).
- (6) If $x \in \text{dom not}_{\text{PP}}(p)$ and $(\text{not}_{\text{PP}}(p))(x) = \text{false}$, then $x \in \text{dom } p$ and $p(x) = \text{true}$. The theorem is a consequence of (3).
- (7) If $x \in \text{dom not}_{\text{PP}}(p)$ and $(\text{not}_{\text{PP}}(p))(x) = \text{true}$, then $x \in \text{dom } p$ and $p(x) = \text{false}$. The theorem is a consequence of (3).

In the sequel D denotes a non empty set and p, q, r denote partial predicates of D .

Let us consider D . The functor $\text{PPdisj}(D)$ yielding a function from $\text{Pr}(D) \times \text{Pr}(D)$ into $\text{Pr}(D)$ is defined by

- (Def. 4) for every partial predicates p, q of D , $\text{dom } it(p, q) = \{d, \text{ where } d \text{ is an element of } D : d \in \text{dom } p \text{ and } p(d) = \text{true} \text{ or } d \in \text{dom } q \text{ and } q(d) = \text{true} \text{ or } d \in \text{dom } p \text{ and } p(d) = \text{false} \text{ and } d \in \text{dom } q \text{ and } q(d) = \text{false}\}$ and for every object d , if $d \in \text{dom } p$ and $p(d) = \text{true}$ or $d \in \text{dom } q$ and $q(d) = \text{true}$, then $it(p, q)(d) = \text{true}$ and if $d \in \text{dom } p$ and $p(d) = \text{false}$ and $d \in \text{dom } q$ and $q(d) = \text{false}$, then $it(p, q)(d) = \text{false}$.

Let us consider p and q . The functor $\text{or}_{\text{PP}}(p, q)$ yielding a partial predicate of D is defined by the term

- (Def. 5) $(\text{PPdisj}(D))(p, q)$.

Let us note that the functor is commutative and idempotent.

Now we state the propositions:

- (8) Suppose $x \in \text{dom or}_{\text{PP}}(p, q)$. Then
- (i) $x \in \text{dom } p$ and $p(x) = \text{true}$, or
 - (ii) $x \in \text{dom } q$ and $q(x) = \text{true}$, or
 - (iii) $x \in \text{dom } p$ and $p(x) = \text{false}$ and $x \in \text{dom } q$ and $q(x) = \text{false}$.
- (9) If $x \in \text{dom } p$ and $x \in \text{dom } q$ and $(\text{or}_{\text{PP}}(p, q))(x) = \text{true}$, then $p(x) = \text{true}$ or $q(x) = \text{true}$. The theorem is a consequence of (3).
- (10) Suppose $x \in \text{dom or}_{\text{PP}}(p, q)$ and $(\text{or}_{\text{PP}}(p, q))(x) = \text{true}$. Then
- (i) $x \in \text{dom } p$ and $p(x) = \text{true}$, or
 - (ii) $x \in \text{dom } q$ and $q(x) = \text{true}$.

The theorem is a consequence of (8) and (9).

- (11) If $x \in \text{dom } p$ and $(\text{or}_{\text{PP}}(p, q))(x) = \text{false}$, then $p(x) = \text{false}$. The theorem is a consequence of (3).
- (12) If $x \in \text{dom } q$ and $(\text{or}_{\text{PP}}(p, q))(x) = \text{false}$, then $q(x) = \text{false}$. The theorem is a consequence of (3).
- (13) Suppose $x \in \text{dom or}_{\text{PP}}(p, q)$ and $(\text{or}_{\text{PP}}(p, q))(x) = \text{false}$. Then

- (i) $x \in \text{dom } p$, and
- (ii) $p(x) = \text{false}$, and
- (iii) $x \in \text{dom } q$, and
- (iv) $q(x) = \text{false}$.

The theorem is a consequence of (8) and (12).

Now we state the propositions:

(14) ASSOCIATIVITY LAW:

$\text{or}_{\text{PP}}(p, \text{or}_{\text{PP}}(q, r)) = \text{or}_{\text{PP}}(\text{or}_{\text{PP}}(p, q), r)$. The theorem is a consequence of (8) and (11).

(15) $\text{or}_{\text{PP}}(\text{or}_{\text{PP}}(p, q), \text{or}_{\text{PP}}(p, r)) = \text{or}_{\text{PP}}(\text{or}_{\text{PP}}(p, q), r)$. The theorem is a consequence of (14).

Let us consider D , p , and q . The functor $\text{and}_{\text{PP}}(p, q)$ yielding a partial predicate of D is defined by the term

(Def. 6) $\text{not}_{\text{PP}}(\text{or}_{\text{PP}}(\text{not}_{\text{PP}}(p), \text{not}_{\text{PP}}(q)))$.

Let us note that the functor is commutative and idempotent. The functor $\text{impl}_{\text{PP}}(p, q)$ yielding a partial predicate of D is defined by the term

(Def. 7) $\text{or}_{\text{PP}}(\text{not}_{\text{PP}}(p), q)$.

Now we state the propositions:

(16) $\text{dom } \text{and}_{\text{PP}}(p, q) = \{d, \text{ where } d \text{ is an element of } D : d \in \text{dom } p \text{ and } p(d) = \text{false} \text{ or } d \in \text{dom } q \text{ and } q(d) = \text{false} \text{ or } d \in \text{dom } p \text{ and } p(d) = \text{true} \text{ and } d \in \text{dom } q \text{ and } q(d) = \text{true}\}$. The theorem is a consequence of (5) and (4).

(17) Suppose $x \in \text{dom } \text{and}_{\text{PP}}(p, q)$. Then

- (i) $x \in \text{dom } p$ and $p(x) = \text{false}$, or
- (ii) $x \in \text{dom } q$ and $q(x) = \text{false}$, or
- (iii) $x \in \text{dom } p$ and $p(x) = \text{true}$ and $x \in \text{dom } q$ and $q(x) = \text{true}$.

The theorem is a consequence of (16).

(18) If $x \in \text{dom } p$ and $p(x) = \text{true}$ and $x \in \text{dom } q$ and $q(x) = \text{true}$, then $(\text{and}_{\text{PP}}(p, q))(x) = \text{true}$.

(19) If $x \in \text{dom } p$ and $p(x) = \text{false}$, then $(\text{and}_{\text{PP}}(p, q))(x) = \text{false}$.

(20) If $x \in \text{dom } q$ and $q(x) = \text{false}$, then $(\text{and}_{\text{PP}}(p, q))(x) = \text{false}$.

(21) If $x \in \text{dom } p$ and $(\text{and}_{\text{PP}}(p, q))(x) = \text{true}$, then $p(x) = \text{true}$.

(22) If $x \in \text{dom } q$ and $(\text{and}_{\text{PP}}(p, q))(x) = \text{true}$, then $q(x) = \text{true}$.

(23) Suppose $x \in \text{dom } \text{and}_{\text{PP}}(p, q)$ and $(\text{and}_{\text{PP}}(p, q))(x) = \text{true}$. Then

- (i) $x \in \text{dom } p$, and

- (ii) $p(x) = true$, and
- (iii) $x \in \text{dom } q$, and
- (iv) $q(x) = true$.

The theorem is a consequence of (17) and (19).

- (24) If $x \in \text{dom } p$ and $x \in \text{dom } q$ and $(\text{and}_{\text{PP}}(p, q))(x) = false$, then $p(x) = false$ or $q(x) = false$. The theorem is a consequence of (18) and (3).
- (25) Suppose $x \in \text{dom } \text{and}_{\text{PP}}(p, q)$ and $(\text{and}_{\text{PP}}(p, q))(x) = false$. Then
- (i) $x \in \text{dom } p$ and $p(x) = false$, or
 - (ii) $x \in \text{dom } q$ and $q(x) = false$.

The theorem is a consequence of (17) and (24).

Now we state the propositions:

- (26) ASSOCIATIVITY LAW:
 $\text{and}_{\text{PP}}(p, \text{and}_{\text{PP}}(q, r)) = \text{and}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), r)$.
- (27) $\text{and}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), \text{and}_{\text{PP}}(p, r)) = \text{and}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), r)$.

Now we state the proposition:

- (28) MEET-ABSORBING LAW:
 $\text{or}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), q) = q$. The theorem is a consequence of (16), (8), (17), (19), and (3).

Now we state the proposition:

- (29) JOIN-ABSORBING LAW:
 $\text{and}_{\text{PP}}(p, \text{or}_{\text{PP}}(p, q)) = p$. The theorem is a consequence of (16), (17), (8), (3), (19), and (18).

Now we state the propositions:

- (30) DISTRIBUTIVITY LAW:
 $\text{and}_{\text{PP}}(p, \text{or}_{\text{PP}}(q, r)) = \text{or}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), \text{and}_{\text{PP}}(p, r))$. The theorem is a consequence of (16), (17), (19), (13), (10), (18), (8), (23), and (25).
- (31) $\text{dom } \text{impl}_{\text{PP}}(p, q) = \{d, \text{ where } d \text{ is an element of } D : d \in \text{dom } p \text{ and } p(d) = false \text{ or } d \in \text{dom } q \text{ and } q(d) = true \text{ or } d \in \text{dom } p \text{ and } p(d) = true \text{ and } d \in \text{dom } q \text{ and } q(d) = false\}$. The theorem is a consequence of (5) and (4).
- (32) Suppose $x \in \text{dom } \text{impl}_{\text{PP}}(p, q)$. Then
- (i) $x \in \text{dom } p$ and $p(x) = false$, or
 - (ii) $x \in \text{dom } q$ and $q(x) = true$, or
 - (iii) $x \in \text{dom } p$ and $p(x) = true$ and $x \in \text{dom } q$ and $q(x) = false$.

The theorem is a consequence of (31).

- (33) If $x \in \text{dom } p$ and $p(x) = \text{false}$, then $(\text{impl}_{\text{PP}}(p, q))(x) = \text{true}$.
- (34) If $x \in \text{dom } q$ and $q(x) = \text{true}$, then $(\text{impl}_{\text{PP}}(p, q))(x) = \text{true}$.
- (35) If $x \in \text{dom } p$ and $p(x) = \text{true}$ and $x \in \text{dom } q$ and $q(x) = \text{false}$, then $(\text{impl}_{\text{PP}}(p, q))(x) = \text{false}$.
- (36) If $x \in \text{dom } p$ and $x \in \text{dom } q$ and $(\text{impl}_{\text{PP}}(p, q))(x) = \text{true}$, then $p(x) = \text{false}$ or $q(x) = \text{true}$. The theorem is a consequence of (35) and (3).
- (37) If $x \in \text{dom } p$ and $(\text{impl}_{\text{PP}}(p, q))(x) = \text{false}$, then $p(x) = \text{true}$.
- (38) If $x \in \text{dom } q$ and $(\text{impl}_{\text{PP}}(p, q))(x) = \text{false}$, then $q(x) = \text{false}$.
- (39) Suppose $x \in \text{dom } \text{impl}_{\text{PP}}(p, q)$ and $(\text{impl}_{\text{PP}}(p, q))(x) = \text{false}$. Then
- (i) $x \in \text{dom } p$, and
 - (ii) $p(x) = \text{true}$, and
 - (iii) $x \in \text{dom } q$, and
 - (iv) $q(x) = \text{false}$.

The theorem is a consequence of (32) and (33).

- (40) Suppose $x \in \text{dom } \text{impl}_{\text{PP}}(p, q)$ and $(\text{impl}_{\text{PP}}(p, q))(x) = \text{true}$. Then
- (i) $x \in \text{dom } p$ and $p(x) = \text{false}$, or
 - (ii) $x \in \text{dom } q$ and $q(x) = \text{true}$.

The theorem is a consequence of (32) and (35).

- (41) $\text{and}_{\text{PP}}(\text{impl}_{\text{PP}}(p, r), \text{impl}_{\text{PP}}(q, r)) = \text{impl}_{\text{PP}}(\text{or}_{\text{PP}}(p, q), r)$. The theorem is a consequence of (30).
- (42) $\text{or}_{\text{PP}}(\text{impl}_{\text{PP}}(p, r), \text{impl}_{\text{PP}}(q, r)) = \text{impl}_{\text{PP}}(\text{and}_{\text{PP}}(p, q), r)$. The theorem is a consequence of (15) and (14).

Let D be a set. The functor $\text{true}_{\text{PP}}(D)$ yielding a partial predicate of D is defined by the term

(Def. 8) $D \mapsto \text{true}$.

Let D be a set. The functor $\text{false}_{\text{PP}}(D)$ yielding a partial predicate of D is defined by the term

(Def. 9) $D \mapsto \text{false}$.

Let us consider a set D . Now we state the propositions:

- (43) $\text{not}_{\text{PP}}(\text{false}_{\text{PP}}(D)) = \text{true}_{\text{PP}}(D)$.
- (44) $\text{not}_{\text{PP}}(\text{true}_{\text{PP}}(D)) = \text{false}_{\text{PP}}(D)$. The theorem is a consequence of (43).

Now we state the propositions:

- (45) $\text{or}_{\text{PP}}(p, \text{true}_{\text{PP}}(D)) = \text{true}_{\text{PP}}(D)$.
- (46) $\text{or}_{\text{PP}}(\text{true}_{\text{PP}}(D), p) = \text{true}_{\text{PP}}(D)$.
- (47) $\text{and}_{\text{PP}}(p, \text{false}_{\text{PP}}(D)) = \text{false}_{\text{PP}}(D)$.

$$(48) \quad \text{and}_{\text{PP}}(\text{false}_{\text{PP}}(D), p) = \text{false}_{\text{PP}}(D).$$

$$(49) \quad \text{or}_{\text{PP}}(p, \text{not}_{\text{PP}}(p)) = \text{true}_{\text{PP}}(D) \upharpoonright \text{dom } p. \text{ The theorem is a consequence of (8) and (3).}$$

$$(50) \quad \text{or}_{\text{PP}}(\text{not}_{\text{PP}}(p), p) = \text{true}_{\text{PP}}(D) \upharpoonright \text{dom } p.$$

$$(51) \quad \text{and}_{\text{PP}}(p, \text{not}_{\text{PP}}(p)) = \text{false}_{\text{PP}}(D) \upharpoonright \text{dom } p. \text{ The theorem is a consequence of (16), (17), (3), and (19).}$$

$$(52) \quad \text{and}_{\text{PP}}(\text{not}_{\text{PP}}(p), p) = \text{false}_{\text{PP}}(D) \upharpoonright \text{dom } p.$$

$$(53) \quad \text{impl}_{\text{PP}}(\text{false}_{\text{PP}}(D), p) = \text{true}_{\text{PP}}(D). \text{ The theorem is a consequence of (43) and (45).}$$

$$(54) \quad \text{impl}_{\text{PP}}(p, \text{true}_{\text{PP}}(D)) = \text{true}_{\text{PP}}(D).$$

$$(55) \quad \text{or}_{\text{PP}}(\text{false}_{\text{PP}}(D) \upharpoonright \text{dom } p, \text{true}_{\text{PP}}(D) \upharpoonright \text{dom } q) = \text{true}_{\text{PP}}(D) \upharpoonright \text{dom } q.$$

Let D be a set. The functor $\perp_{\text{PP}}(D)$ yielding a partial predicate of D is defined by the term

$$\text{(Def. 10)} \quad \emptyset.$$

Now we state the propositions:

$$(56) \quad \text{Let us consider a set } D. \text{ Then } \text{not}_{\text{PP}}(\perp_{\text{PP}}(D)) = \perp_{\text{PP}}(D).$$

$$(57) \quad \text{or}_{\text{PP}}(\perp_{\text{PP}}(D), \text{true}_{\text{PP}}(D)) = \text{true}_{\text{PP}}(D).$$

$$(58) \quad \text{and}_{\text{PP}}(\perp_{\text{PP}}(D), \text{false}_{\text{PP}}(D)) = \text{false}_{\text{PP}}(D).$$

$$(59) \quad \text{impl}_{\text{PP}}(\perp_{\text{PP}}(D), \text{true}_{\text{PP}}(D)) = \text{true}_{\text{PP}}(D). \text{ The theorem is a consequence of (56) and (57).}$$

2. ALGEBRA OF PARTIAL CONNECTIVES WITH (STRONG) KLEENE LOGICAL CONNECTIVES

Let us consider D . The functors: $\text{PartPredConnMeet}(D)$ and $\text{PartPredConnJoin}(D)$ yielding binary operations on $\text{Pr}(D)$ are defined by conditions

$$\text{(Def. 11)} \quad \text{for every partial predicates } p, q \text{ of } D, \text{ PartPredConnMeet}(D)(p, q) = \text{and}_{\text{PP}}(p, q),$$

$$\text{(Def. 12)} \quad \text{for every partial predicates } p, q \text{ of } D, \text{ PartPredConnJoin}(D)(p, q) = \text{or}_{\text{PP}}(p, q),$$

respectively. The functor $\text{PartPredConnCompl}(D)$ yielding a unary operation on $\text{Pr}(D)$ is defined by

$$\text{(Def. 13)} \quad \text{for every partial predicate } p \text{ of } D, \text{ it}(p) = \text{not}_{\text{PP}}(p).$$

The functor $\text{PartPredConnLatt}(D)$ yielding a strict ortholattice structure is defined by the term

$$\text{(Def. 14)} \quad \langle \text{Pr}(D), \text{PartPredConnJoin}(D), \text{PartPredConnMeet}(D), \text{PartPredConnCompl}(D) \rangle$$

Let D be a non empty set, f, g be binary operations on D , and h be a unary operation on D . Let us observe that $\langle D, f, g, h \rangle$ is non empty.

Let us consider D . Note that $\text{PartPredConnLatt}(D)$ is non empty and constituted functions and there exists a lattice structure which is non empty and constituted functions and there exists an ortholattice structure which is strict, non empty, and constituted functions.

Let us consider D . Let us observe that $\text{PartPredConnLatt}(D)$ is lattice-like and $\text{PartPredConnLatt}(D)$ is bounded and $\text{PartPredConnLatt}(D)$ is de Morgan and join-idempotent and has idempotent element.

Now we state the propositions:

$$(60) \quad \top_{\text{PartPredConnLatt}(D)} = \text{true}_{\text{PP}}(D).$$

$$(61) \quad \perp_{\text{PartPredConnLatt}(D)} = \text{false}_{\text{PP}}(D).$$

Let L be a non empty, constituted functions lattice structure and a, b be elements of L . We say that a is a partial complement of b if and only if

$$\text{(Def. 15)} \quad a \sqcup b = \top_L \upharpoonright \text{dom } b \text{ and } b \sqcup a = \top_L \upharpoonright \text{dom } b \text{ and } a \sqcap b = \perp_L \upharpoonright \text{dom } b \text{ and } b \sqcap a = \perp_L \upharpoonright \text{dom } b.$$

We say that L is partially complemented if and only if

$$\text{(Def. 16)} \quad \text{for every element } b \text{ of } L, \text{ there exists an element } a \text{ of } L \text{ such that } a \text{ is a partial complement of } b.$$

Let I be a constituted functions, non empty lattice structure. We say that I is partially Boolean if and only if

$$\text{(Def. 17)} \quad I \text{ is bounded, partially complemented, and distributive.}$$

Let us observe that every constituted functions, non empty lattice structure which is partially Boolean is also bounded, partially complemented, and distributive and every constituted functions, non empty lattice structure which is bounded, partially complemented, and distributive is also partially Boolean.

Now we state the proposition:

$$(62) \quad \text{Let us consider elements } a, b \text{ of } \text{PartPredConnLatt}(D). \text{ If } a = p \text{ and } b = \text{not}_{\text{PP}}(p), \text{ then } b \text{ is a partial complement of } a. \text{ The theorem is a consequence of (60), (49), (61), and (51).}$$

Let us consider D . One can check that $\text{PartPredConnLatt}(D)$ is partially Boolean.

Now we state the proposition:

$$(63) \quad \text{DISTRIBUTIVITY LAW:}$$

$$\text{or}_{\text{PP}}(p, \text{and}_{\text{PP}}(q, r)) = \text{and}_{\text{PP}}(\text{or}_{\text{PP}}(p, q), \text{or}_{\text{PP}}(p, r)).$$

Let L be a non empty ortholattice structure. We say that L is Kleene if and only if

$$\text{(Def. 18)} \quad \text{for every elements } x, y \text{ of } L, x \sqcap x^c \sqsubseteq y \sqcup y^c.$$

Observe that every meet-absorbing, join-absorbing, meet-commutative, non empty ortholattice structure which is Boolean and well-complemented is also Kleene.

Let us consider D . Let us note that $\text{PartPredConnLatt}(D)$ is Kleene and there exists a non empty, constituted functions lattice structure which is partially Boolean, join-idempotent, and lattice-like and there exists a non empty ortholattice structure which is Kleene, de Morgan, join-idempotent, lattice-like, and strict and has idempotent element and there exists a non empty, constituted functions ortholattice structure which is partially Boolean, Kleene, de Morgan, join-idempotent, lattice-like, and strict and has idempotent element.

REFERENCES

- [1] Raymond Balbes and Philip Dwinger. *Distributive Lattices*. University of Missouri Press, 1975.
- [2] T.S. Blyth and J. Varlet. *Ockham Algebras*. Oxford science publications. Oxford University Press, 1994.
- [3] Diana Brignole and Antonio Monteiro. *Caractérisation des algèbres de Nelson par des égalités*. Instituto de Matemática, Universidad Nacional del Sur, Argentina, 1964.
- [4] Roberto Cignoli. Injective de Morgan and Kleene algebras. *Proceedings of the American Mathematical Society*, 47(2):269–278, 1975.
- [5] J.P. Cleave. *A Study of Logics*. Oxford logic guides. Clarendon Press, 1991. ISBN 9780198532118.
- [6] J.H. Conway. *Regular algebra and finite machines*. Chapman and Hall mathematics series. Chapman and Hall, 1971.
- [7] R.W. Floyd. Assigning meanings to programs. *Mathematical aspects of computer science*, 19(19–32), 1967.
- [8] Adam Grabowski. Robbins algebras vs. Boolean algebras. *Formalized Mathematics*, 9(4): 681–690, 2001.
- [9] C.A.R. Hoare. An axiomatic basis for computer programming. *Commun. ACM*, 12(10): 576–580, 1969.
- [10] Ievgen Ivanov. On the underapproximation of reach sets of abstract continuous-time systems. In Erika Ábrahám and Sergiy Bogomolov, editors, *Proceedings 3rd International Workshop on Symbolic and Numerical Methods for Reachability Analysis, SNR@ETAPS 2017, Uppsala, Sweden, 22nd April 2017*, volume 247 of *EPTCS*, pages 46–51, 2017. doi:10.4204/EPTCS.247.4.
- [11] Ievgen Ivanov. On representations of abstract systems with partial inputs and outputs. In T. V. Gopal, Manindra Agrawal, Angsheng Li, and S. Barry Cooper, editors, *Theory and Applications of Models of Computation – 11th Annual Conference, TAMC 2014, Chennai, India, April 11–13, 2014. Proceedings*, volume 8402 of *Lecture Notes in Computer Science*, pages 104–123. Springer, 2014. ISBN 978-3-319-06088-0. doi:10.1007/978-3-319-06089-7_8.
- [12] Ievgen Ivanov. On local characterization of global timed bisimulation for abstract continuous-time systems. In Ichiro Hasuo, editor, *Coalgebraic Methods in Computer Science – 13th IFIP WG 1.3 International Workshop, CMCS 2016, Colocated with ETAPS 2016, Eindhoven, The Netherlands, April 2–3, 2016, Revised Selected Papers*, volume 9608 of *Lecture Notes in Computer Science*, pages 216–234. Springer, 2016. ISBN 978-3-319-40369-4. doi:10.1007/978-3-319-40370-0_13.
- [13] Ievgen Ivanov, Mykola Nikitchenko, and Uri Abraham. *On a Decidable Formal Theory for Abstract Continuous-Time Dynamical Systems*, pages 78–99. Springer International Publishing, 2014. ISBN 978-3-319-13206-8. doi:10.1007/978-3-319-13206-8_4.
- [14] Ievgen Ivanov, Mykola Nikitchenko, and Uri Abraham. Event-based proof of the mutual exclusion property of Peterson’s algorithm. *Formalized Mathematics*, 23(4):325–331, 2015.

doi:10.1515/forma-2015-0026.

- [15] Ievgen Ivanov, Mykola Nikitchenko, and Volodymyr G. Skobelev. Proving properties of programs on hierarchical nominative data. *The Computer Science Journal of Moldova*, 24(3):371–398, 2016.
- [16] J. A. Kalman. Lattices with involution. *Transactions of the American Mathematical Society*, 87(2):485–485, February 1958. doi:10.1090/s0002-9947-1958-0095135-x.
- [17] S.C. Kleene. *Introduction to Metamathematics*. North-Holland Publishing Co., Amsterdam, and P. Noordhoff, Groningen, 1952.
- [18] S. Körner. *Experience and Theory: An Essay in the Philosophy of Science*. International library of philosophy and scientific method. Routledge & Kegan Paul, 1966.
- [19] Artur Kornilowicz, Andrii Kryvolap, Mykola Nikitchenko, and Ievgen Ivanov. Formalization of the algebra of nominative data in Mizar. In Maria Ganzha, Leszek A. Maciaszek, and Marcin Paprzycki, editors, *Proceedings of the 2017 Federated Conference on Computer Science and Information Systems, FedCSIS 2017, Prague, Czech Republic, September 3–6, 2017.*, pages 237–244, 2017. ISBN 978-83-946253-7-5. doi:10.15439/2017F301.
- [20] Artur Kornilowicz, Andrii Kryvolap, Mykola Nikitchenko, and Ievgen Ivanov. An approach to formalization of an extension of Floyd-Hoare logic. In Vadim Ermolayev, Nick Bassiliades, Hans-Georg Fill, Vitaliy Yakovyna, Heinrich C. Mayr, Vyacheslav Kharchenko, Vladimir Peschanenko, Mariya Shyshkina, Mykola Nikitchenko, and Aleksander Spivakovsky, editors, *Proceedings of the 13th International Conference on ICT in Education, Research and Industrial Applications. Integration, Harmonization and Knowledge Transfer, Kyiv, Ukraine, May 15–18, 2017*, volume 1844 of *CEUR Workshop Proceedings*, pages 504–523. CEUR-WS.org, 2017.
- [21] Artur Kornilowicz, Andrii Kryvolap, Mykola Nikitchenko, and Ievgen Ivanov. *Formalization of the Nominative Algorithmic Algebra in Mizar*, pages 176–186. Springer International Publishing, 2018. ISBN 978-3-319-67229-8. doi:10.1007/978-3-319-67229-8_16.
- [22] Dexter Kozen. *On Kleene algebras and closed semirings*, pages 26–47. Springer Berlin Heidelberg, 1990. doi:10.1007/BFb0029594.
- [23] Andrii Kryvolap, Mykola Nikitchenko, and Wolfgang Schreiner. *Extending Floyd-Hoare Logic for Partial Pre- and Postconditions*, pages 355–378. Springer International Publishing, 2013. ISBN 978-3-319-03998-5. doi:10.1007/978-3-319-03998-5_18.
- [24] Antonio Monteiro and Luiz Monteiro. Axiomes indépendants pour les algèbres de Nelson, de Łukasiewicz trivalentes, de de Morgan et de Kleene. *Notas de lógica matemática*, (40): 1–11, 1996.
- [25] Maurizio Negri. Three valued semantics and DMF-algebras. *Boll. Un. Mat. Ital. B (7)*, 10(3):733–760, 1996.
- [26] Maurizio Negri. DMF-algebras: representation and topological characterization. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8)*, 1(2):369–390, 1998.
- [27] Maurizio Negri. Partial probability and Kleene logic, 2013.
- [28] M.S. Nikitchenko and S.S. Shkilniak. *Mathematical logic and theory of algorithms*. Publishing house of Taras Shevchenko National University of Kyiv, Ukraine (in Ukrainian), 2008.
- [29] M.S. Nikitchenko and S.S. Shkilniak. *Applied logic*. Publishing house of Taras Shevchenko National University of Kyiv, Ukraine (in Ukrainian), 2013.
- [30] Mykola Nikitchenko and Stepan Shkilniak. Algebras and logics of partial quasiary predicates. *Algebra and Discrete Mathematics*, 23(2):263–278, 2017.
- [31] Nikolaj S. Nikitchenko. A composition nominative approach to program semantics. Technical Report IT-TR 1998-020, Department of Information Technology, Technical University of Denmark, 1998.
- [32] Helena Rasiowa. *An Algebraic Approach to Non-Classical Logics*. North Holland, 1974.
- [33] Volodymyr G. Skobelev, Mykola Nikitchenko, and Ievgen Ivanov. On algebraic properties of nominative data and functions. In Vadim Ermolayev, Heinrich C. Mayr, Mykola Nikitchenko, Aleksander Spivakovsky, and Grygoriy Zholtkevych, editors, *Information and Communication Technologies in Education, Research, and Industrial Applications – 10th International Conference, ICTERI 2014, Kherson, Ukraine, June 9–12, 2014, Revised Selected Papers*, volume 469 of *Communications in Computer and Information Science*, pages 117–138. Springer, 2014. ISBN 978-3-319-13205-1. doi:10.1007/978-3-319-13206-8_6.

Received March 27, 2018
