

# Finite Sums of Vectors in Vector Space

Wojciech A. Trybulec  
Warsaw University

**Summary.** We define the sum of finite sequences of vectors in vector space. Theorems concerning those sums are proved.

MML Identifier: VECTSP\_3.

The terminology and notation used here have been introduced in the following papers: [7], [2], [3], [5], [6], [4], and [1]. Let  $F$  be a field. An element of  $F$  is an element of the carrier of  $F$ .

For simplicity we follow a convention:  $x$  will be arbitrary,  $G_1$  will denote a field,  $a$  will denote an element of  $G_1$ ,  $V$  will denote a vector space over  $G_1$ , and  $v, v_1, v_2, w, u$  will denote vectors of  $V$ . Let us consider  $G_1, V, x$ . The predicate  $x \in V$  is defined by:

(Def.1)  $x \in$  the carrier of the carrier of  $V$ .

Next we state two propositions:

- (1)  $x \in V$  if and only if  $x \in$  the carrier of the carrier of  $V$ .
- (2)  $v \in V$ .

We follow a convention:  $F, G, H$  will be finite sequences of elements of the carrier of the carrier of  $V$ ,  $f$  will be a function from  $\mathbb{N}$  into the carrier of the carrier of  $V$ , and  $i, j, k, n$  will be natural numbers. Let us consider  $G_1, V, f, j$ . Then  $f(j)$  is a vector of  $V$ .

Let us consider  $G_1, V, F$ . The functor  $\sum F$  yielding a vector of  $V$  is defined as follows:

(Def.2) there exists  $f$  such that  $\sum F = f(\text{len } F)$  and  $f(0) = \Theta_V$  and for all  $j, v$  such that  $j < \text{len } F$  and  $v = F(j+1)$  holds  $f(j+1) = f(j) + v$ .

We now state a number of propositions:

- (3) If there exists  $f$  such that  $u = f(\text{len } F)$  and  $f(0) = \Theta_V$  and for all  $j, v$  such that  $j < \text{len } F$  and  $v = F(j+1)$  holds  $f(j+1) = f(j) + v$ , then  $u = \sum F$ .

- (4) There exists  $f$  such that  $\sum F = f(\text{len } F)$  and  $f(0) = \Theta_V$  and for all  $j, v$  such that  $j < \text{len } F$  and  $v = F(j+1)$  holds  $f(j+1) = f(j) + v$ .
- (5) If  $k \in \text{Seg } n$  and  $\text{len } F = n$ , then  $F(k)$  is a vector of  $V$ .
- (6) If  $\text{len } F = \text{len } G + 1$  and  $G = F \upharpoonright \text{Seg}(\text{len } G)$  and  $v = F(\text{len } F)$ , then  $\sum F = \sum G + v$ .
- (7)  $\sum(F \frown G) = \sum F + \sum G$ .
- (8) If  $\text{len } F = \text{len } G$  and  $\text{len } F = \text{len } H$  and for every  $k$  such that  $k \in \text{Seg}(\text{len } F)$  holds  $H(k) = \pi_k F + \pi_k G$ , then  $\sum H = \sum F + \sum G$ .
- (9) If  $\text{len } F = \text{len } G$  and for all  $k, v$  such that  $k \in \text{Seg}(\text{len } F)$  and  $v = G(k)$  holds  $F(k) = a \cdot v$ , then  $\sum F = a \cdot \sum G$ .
- (10) If  $\text{len } F = \text{len } G$  and for every  $k$  such that  $k \in \text{Seg}(\text{len } F)$  holds  $G(k) = a \cdot \pi_k F$ , then  $\sum G = a \cdot \sum F$ .
- (11) If  $\text{len } F = \text{len } G$  and for all  $k, v$  such that  $k \in \text{Seg}(\text{len } F)$  and  $v = G(k)$  holds  $F(k) = -v$ , then  $\sum F = -\sum G$ .
- (12) If  $\text{len } F = \text{len } G$  and for every  $k$  such that  $k \in \text{Seg}(\text{len } F)$  holds  $G(k) = -\pi_k F$ , then  $\sum G = -\sum F$ .
- (13) If  $\text{len } F = \text{len } G$  and  $\text{len } F = \text{len } H$  and for every  $k$  such that  $k \in \text{Seg}(\text{len } F)$  holds  $H(k) = \pi_k F - \pi_k G$ , then  $\sum H = \sum F - \sum G$ .
- (14) If  $\text{rng } F = \text{rng } G$  and  $F$  is one-to-one and  $G$  is one-to-one, then  $\sum F = \sum G$ .
- (15) For all  $F, G$  and for every permutation  $f$  of  $\text{dom } F$  such that  $\text{len } F = \text{len } G$  and for every  $i$  such that  $i \in \text{dom } G$  holds  $G(i) = F(f(i))$  holds  $\sum F = \sum G$ .
- (16) For every permutation  $f$  of  $\text{dom } F$  such that  $G = F \cdot f$  holds  $\sum F = \sum G$ .
- (17)  $\sum \varepsilon_{\text{the carrier of the carrier of } V} = \Theta_V$ .
- (18)  $\sum \langle v \rangle = v$ .
- (19)  $\sum \langle v, u \rangle = v + u$ .
- (20)  $\sum \langle v, u, w \rangle = (v + u) + w$ .
- (21)  $a \cdot \sum \varepsilon_{\text{the carrier of the carrier of } V} = \Theta_V$ .
- (22)  $a \cdot \sum \langle v \rangle = a \cdot v$ .
- (23)  $a \cdot \sum \langle v, u \rangle = a \cdot v + a \cdot u$ .
- (24)  $a \cdot \sum \langle v, u, w \rangle = (a \cdot v + a \cdot u) + a \cdot w$ .
- (25)  $-\sum \varepsilon_{\text{the carrier of the carrier of } V} = \Theta_V$ .
- (26)  $-\sum \langle v \rangle = -v$ .
- (27)  $-\sum \langle v, u \rangle = (-v) - u$ .
- (28)  $-\sum \langle v, u, w \rangle = ((-v) - u) - w$ .
- (29)  $\sum \langle v, w \rangle = \sum \langle w, v \rangle$ .
- (30)  $\sum \langle v, w \rangle = \sum \langle v \rangle + \sum \langle w \rangle$ .
- (31)  $\sum \langle \Theta_V, \Theta_V \rangle = \Theta_V$ .
- (32)  $\sum \langle \Theta_V, v \rangle = v$  and  $\sum \langle v, \Theta_V \rangle = v$ .

- (33)  $\sum\langle v, -v \rangle = \Theta_V$  and  $\sum\langle -v, v \rangle = \Theta_V$ .
- (34)  $\sum\langle v, -w \rangle = v - w$  and  $\sum\langle -w, v \rangle = v - w$ .
- (35)  $\sum\langle -v, -w \rangle = -(v + w)$  and  $\sum\langle -w, -v \rangle = -(v + w)$ .
- (36)  $\sum\langle u, v, w \rangle = (\sum\langle u \rangle + \sum\langle v \rangle) + \sum\langle w \rangle$ .
- (37)  $\sum\langle u, v, w \rangle = \sum\langle u, v \rangle + w$ .
- (38)  $\sum\langle u, v, w \rangle = \sum\langle v, w \rangle + u$ .
- (39)  $\sum\langle u, v, w \rangle = \sum\langle u, w \rangle + v$ .
- (40)  $\sum\langle u, v, w \rangle = \sum\langle u, w, v \rangle$ .
- (41)  $\sum\langle u, v, w \rangle = \sum\langle v, u, w \rangle$ .
- (42)  $\sum\langle u, v, w \rangle = \sum\langle v, w, u \rangle$ .
- (43)  $\sum\langle u, v, w \rangle = \sum\langle w, u, v \rangle$ .
- (44)  $\sum\langle u, v, w \rangle = \sum\langle w, v, u \rangle$ .
- (45)  $\sum\langle \Theta_V, \Theta_V, \Theta_V \rangle = \Theta_V$ .
- (46)  $\sum\langle \Theta_V, \Theta_V, v \rangle = v$  and  $\sum\langle \Theta_V, v, \Theta_V \rangle = v$  and  $\sum\langle v, \Theta_V, \Theta_V \rangle = v$ .
- (47)  $\sum\langle \Theta_V, u, v \rangle = u + v$  and  $\sum\langle u, v, \Theta_V \rangle = u + v$  and  $\sum\langle u, \Theta_V, v \rangle = u + v$ .
- (48) If  $\text{len } F = 0$ , then  $\sum F = \Theta_V$ .
- (49) If  $\text{len } F = 1$ , then  $\sum F = F(1)$ .
- (50) If  $\text{len } F = 2$  and  $v_1 = F(1)$  and  $v_2 = F(2)$ , then  $\sum F = v_1 + v_2$ .
- (51) If  $\text{len } F = 3$  and  $v_1 = F(1)$  and  $v_2 = F(2)$  and  $v = F(3)$ , then  $\sum F = (v_1 + v_2) + v$ .
- (52)  $v - v = \Theta_V$ .
- (53)  $-(v + w) = (-v) + (-w)$ .

## References

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [4] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [5] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [6] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.

- [7] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.

*Received July 12, 1990*

---