

Determinant of Some Matrices of Field Elements

Yatsuka Nakamura
Shinshu University
Nagano, Japan

Summary. Here, we present determinants of some square matrices of field elements. First, the determinant of 2×2 matrix is shown. Secondly, the determinants of zero matrix and unit matrix are shown, which are equal to 0 in the field and 1 in the field respectively. Thirdly, the determinant of diagonal matrix is shown, which is a product of all diagonal elements of the matrix. At the end, we prove that the determinant of a matrix is the same as the determinant of its transpose.

MML identifier: MATRIX_7, version: 7.6.01 4.53.937

The articles [19], [26], [2], [27], [5], [4], [8], [24], [18], [17], [14], [6], [23], [7], [25], [20], [21], [3], [12], [28], [10], [15], [16], [11], [13], [1], [9], and [22] provide the notation and terminology for this paper.

In this paper n, i, l are natural numbers.

The following propositions are true:

- (1) For every permutation f of Seg 2 holds $f = \langle 1, 2 \rangle$ or $f = \langle 2, 1 \rangle$.
- (2) For every finite sequence f such that $f = \langle 1, 2 \rangle$ or $f = \langle 2, 1 \rangle$ holds f is a permutation of Seg 2.
- (3) The permutations of 2-element set = $\{\langle 1, 2 \rangle, \langle 2, 1 \rangle\}$.
- (4) For every permutation p of Seg 2 such that p is a transposition holds $p = \langle 2, 1 \rangle$.
- (5) Let D be a non empty set, f be a finite sequence of elements of D , and k_2 be a natural number. If $1 \leq k_2$ and $k_2 < \text{len } f$, then $f = (\text{mid}(f, 1, k_2)) \hat{\ } \text{mid}(f, k_2 + 1, \text{len } f)$.
- (6) For every non empty set D and for every finite sequence f of elements of D such that $2 \leq \text{len } f$ holds $f = (f \upharpoonright (\text{len } f - 1)) \hat{\ } \text{mid}(f, \text{len } f - 1, \text{len } f)$.

- (7) For every non empty set D and for every finite sequence f of elements of D such that $1 \leq \text{len } f$ holds $f = (f \upharpoonright (\text{len } f - 1)) \hat{\ } \text{mid}(f, \text{len } f, \text{len } f)$.
- (8) Let a be an element of A_2 . Given an element q of the permutations of 2-element set such that $q = a$ and q is a transposition. Then $a = \langle 2, 1 \rangle$.
- (9) Let n be a natural number, a, b be elements of A_n , and p_2, p_1 be elements of the permutations of n -element set. If $a = p_2$ and $b = p_1$, then $a \cdot b = p_1 \cdot p_2$.
- (10) Let a, b be elements of A_2 . Suppose that
- (i) there exists an element p of the permutations of 2-element set such that $p = a$ and p is a transposition, and
 - (ii) there exists an element q of the permutations of 2-element set such that $q = b$ and q is a transposition.
- Then $a \cdot b = \langle 1, 2 \rangle$.
- (11) Let l be a finite sequence of elements of A_2 . Suppose that
- (i) $\text{len } l \bmod 2 = 0$, and
 - (ii) for every i such that $i \in \text{dom } l$ there exists an element q of the permutations of 2-element set such that $l(i) = q$ and q is a transposition.
- Then $\prod l = \langle 1, 2 \rangle$.
- (12) For every field K and for every matrix M over K of dimension 2 holds $\text{Det } M = M_{1,1} \cdot M_{2,2} - M_{1,2} \cdot M_{2,1}$.

Let n be a natural number, let K be a field, let M be a matrix over K of dimension n , and let a be an element of K . Then $a \cdot M$ is a matrix over K of dimension n .

The following three propositions are true:

- (13) For every field K and for all natural numbers n, m holds
- $$\text{len} \left(\begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_{n \times m} \right) = n \text{ and } \text{dom} \left(\begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_{n \times m} \right) = \text{Seg } n.$$
- (14) Let K be a field, n be a natural number, p be an element of the permutations of n -element set, and i be a natural number. If $i \in \text{Seg } n$, then $p(i) \in \text{Seg } n$.
- (15) For every field K and for every natural number n such that $n \geq 1$ holds
- $$\text{Det} \left(\begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_{n \times n} \right) = 0_K.$$

Let x, y, a, b be sets. The functor $\text{IFIN}(x, y, a, b)$ is defined by:

$$(\text{Def. 1}) \quad \text{IFIN}(x, y, a, b) = \begin{cases} a, & \text{if } x \in y, \\ b, & \text{otherwise.} \end{cases}$$

We now state the proposition

(16) For every field K and for every natural number n such that $n \geq 1$ holds

$$\text{Det} \left(\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_{K}^{n \times n} \right) = 1_K.$$

Let K be a field, let n be a natural number, and let M be a matrix over K of dimension n . We say that M being diagonal if and only if:

(Def. 2) For all natural numbers i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } n$ and $i \neq j$ holds $M_{i,j} = 0_K$.

One can prove the following propositions:

(17) Let K be a field, n be a natural number, and A be a matrix over K of dimension n . Suppose $n \geq 1$ and A being diagonal. Then $\text{Det } A =$ (the multiplication of K) \otimes (the diagonal of A).

(18) Let n be a natural number and p be an element of the permutations of n -element set. Then p^{-1} is an element of the permutations of n -element set.

Let us consider n and let p be an element of the permutations of n -element set. Then p^{-1} is an element of the permutations of n -element set.

Next we state the proposition

(19) Let n be a natural number, K be a field, and A be a matrix over K of dimension n . Then A^T is a matrix over K of dimension n .

Let n be a natural number, let K be a field, and let A be a matrix over K of dimension n . The functor A^T yields a matrix over K of dimension n and is defined as follows:

(Def. 3) $A^T = (A \text{ qua matrix over } K)^T$.

The following proposition is true

(20) For every group G and for all finite sequences f_1, f_2 of elements of G holds $(\prod(f_1 \wedge f_2))^{-1} = (\prod f_2)^{-1} \cdot (\prod f_1)^{-1}$.

Let G be a group and let f be a finite sequence of elements of G . The functor f^{-1} yields a finite sequence of elements of G and is defined by:

(Def. 4) $\text{len}(f^{-1}) = \text{len } f$ and for every natural number i such that $i \in \text{Seg len } f$ holds $(f^{-1})_i = (f_i)^{-1}$.

One can prove the following propositions:

(21) For every group G holds $(\varepsilon_{(\text{the carrier of } G)})^{-1} = \varepsilon_{(\text{the carrier of } G)}$.

(22) For every group G and for all finite sequences f, g of elements of G holds $(f \wedge g)^{-1} = (f^{-1}) \wedge g^{-1}$.

(23) For every group G and for every element a of G holds $\langle a \rangle^{-1} = \langle a^{-1} \rangle$.

(24) For every group G and for every finite sequence f of elements of G holds $\prod(f \wedge (\text{Rev}(f)))^{-1} = 1_G$.

- (25) For every group G and for every finite sequence f of elements of G holds $\prod(((\text{Rev}(f))^{-1}) \wedge f) = 1_G$.
- (26) For every group G and for every finite sequence f of elements of G holds $(\prod f)^{-1} = \prod((\text{Rev}(f))^{-1})$.
- (27) Let I_1 be an element of the permutations of n -element set and I_2 be an element of A_n . If $I_2 = I_1$ and $n \geq 1$, then $I_1^{-1} = I_2^{-1}$.
- (28) Let n be a natural number and I_3 be an element of the permutations of n -element set. If $n \geq 1$, then I_3 is even iff I_3^{-1} is even.
- (29) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and x be an element of K . If $n \geq 1$, then $(-1)^{\text{sgn}(p)}x = (-1)^{\text{sgn}(p^{-1})}x$.
- (30) Let K be a field and f_1, f_2 be finite sequences of elements of K . Then (the multiplication of K) $\otimes (f_1 \wedge f_2) = ((\text{the multiplication of } K) \otimes (f_1)) \cdot ((\text{the multiplication of } K) \otimes (f_2))$.
- (31) Let K be a field and R_1, R_2 be finite sequences of elements of K . Suppose R_1 and R_2 are fiberwise equipotent. Then (the multiplication of K) $\otimes (R_1) = (\text{the multiplication of } K) \otimes (R_2)$.
- (32) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and f, g be finite sequences of elements of K . If $n \geq 1$ and $\text{len } f = n$ and $g = f \cdot p$, then f and g are fiberwise equipotent.
- (33) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and f, g be finite sequences of elements of K . Suppose $n \geq 1$ and $\text{len } f = n$ and $g = f \cdot p$. Then (the multiplication of K) $\otimes f = (\text{the multiplication of } K) \otimes g$.
- (34) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and f be a finite sequence of elements of K . If $n \geq 1$ and $\text{len } f = n$, then $f \cdot p$ is a finite sequence of elements of K .
- (35) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and A be a matrix over K of dimension n . If $n \geq 1$, then $p^{-1}\text{-Path } A^T = (p\text{-Path } A) \cdot p^{-1}$.
- (36) Let n be a natural number, K be a field, p be an element of the permutations of n -element set, and A be a matrix over K of dimension n . Suppose $n \geq 1$. Then (the product on paths of A^T)(p^{-1}) = (the product on paths of A)(p).
- (37) Let n be a natural number, K be a field, and A be a matrix over K of dimension n . If $n \geq 1$, then $\text{Det } A = \text{Det}(A^T)$.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.

- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [7] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [8] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [9] Czesław Byliński. Some properties of restrictions of finite sequences. *Formalized Mathematics*, 5(2):241–245, 1996.
- [10] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [11] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Formalized Mathematics*, 2(4):475–480, 1991.
- [12] Katarzyna Jankowska. Transpose matrices and groups of permutations. *Formalized Mathematics*, 2(5):711–717, 1991.
- [13] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Formalized Mathematics*, 3(2):275–278, 1992.
- [14] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [15] Yatsuka Nakamura and Roman Matuszewski. Reconstructions of special sequences. *Formalized Mathematics*, 6(2):255–263, 1997.
- [16] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [17] Library Committee of the Association of Mizar Users. Binary operations on numbers. *To appear in Formalized Mathematics*.
- [18] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [19] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [20] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Formalized Mathematics*, 1(1):187–190, 1990.
- [21] Wojciech A. Trybulec. Binary operations on finite sequences. *Formalized Mathematics*, 1(5):979–981, 1990.
- [22] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [23] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [24] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [25] Wojciech A. Trybulec. Lattice of subgroups of a group. Frattini subgroup. *Formalized Mathematics*, 2(1):41–47, 1991.
- [26] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [27] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [28] Katarzyna Zawadzka. The product and the determinant of matrices with entries in a field. *Formalized Mathematics*, 4(1):1–8, 1993.

Received January 4, 2006
