

# Contents

*Formaliz. Math.* 24 (3)

<b>Compactness in Metric Spaces</b> By KAZUHISA NAKASHO <i>et al.</i> .....	167
<b>Double Sequences and Iterated Limits in Regular Space</b> By ROLAND COGHETTO .....	173
<b>Prime Factorization of Sums and Differences of Two Like Powers</b> By RAFAL ZIOBRO .....	187
<b>Riemann-Stieltjes Integral</b> By KEIKO NARITA <i>et al.</i> .....	199
<b>Quasi-uniform Space</b> By ROLAND COGHETTO .....	205
<b>Uniform Space</b> By ROLAND COGHETTO .....	215
<b>Some Algebraic Properties of Polynomial Rings</b> By CHRISTOPH SCHWARZWELLER <i>et al.</i> .....	227



# Compactness in Metric Spaces

Kazuhisa Nakasho  
Shinshu University  
Nagano, Japan

Keiko Narita  
Hirosaki-city  
Aomori, Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** In this article, we mainly formalize in Mizar [2] the equivalence among a few compactness definitions of metric spaces, norm spaces, and the real line. In the first section, we formalized general topological properties of metric spaces. We discussed openness and closedness of subsets in metric spaces in terms of convergence of element sequences. In the second section, we firstly formalize the definition of sequentially compact, and then discuss the equivalence of compactness, countable compactness, sequential compactness, and totally boundedness with completeness in metric spaces.

In the third section, we discuss compactness in norm spaces. We formalize the equivalence of compactness and sequential compactness in norm space. In the fourth section, we formalize topological properties of the real line in terms of convergence of real number sequences. In the last section, we formalize the equivalence of compactness and sequential compactness in the real line. These formalizations are based on [20], [5], [17], [14], and [4].

MSC: 46B50 54E45 03B35

Keywords: metric spaces; normed linear spaces; compactness

MML identifier: TOPMETR4, version: 8.1.05 5.37.1275

## 1. TOPOLOGICAL PROPERTIES OF METRIC SPACES

Now we state the propositions:

- (1) Let us consider a non empty set  $M$ , and a sequence  $x$  of  $M$ . Suppose  $\text{rng } x$  is finite. Then there exists an element  $z$  of  $M$  such that
  - (i)  $x^{-1}(\{z\}) \subseteq \mathbb{N}$ , and
  - (ii)  $x^{-1}(\{z\})$  is infinite.

PROOF: Define  $\mathcal{X}(\text{object}) = x^{-1}(\{\$1\})$ . Set  $K = \{\mathcal{X}(w)$ , where  $w$  is an element of  $M : w \in \text{rng } x\}$ .  $K$  is finite from [18, Sch. 21]. For every set  $Y$  such that  $Y \in K$  holds  $Y$  is finite.  $\text{dom } x \subseteq \bigcup K$  by [6, (3)].  $\square$

- (2) Let us consider a subset  $X$  of  $\mathbb{N}$ . Suppose  $X$  is infinite. Then there exists an increasing sequence  $N$  of  $\mathbb{N}$  such that  $\text{rng } N \subseteq X$ .

PROOF: Reconsider  $B = 2^X$  as a non empty set. Reconsider  $N_0 = \min^* X$  as an element of  $\mathbb{N}$ . Reconsider  $Y_0 = X$  as an element of  $B$ . Define  $\mathcal{P}[\text{object}, \text{object}, \text{set}, \text{object}, \text{set}] \equiv \$5 = \$3 \setminus \{\$2\}$  and  $\$4 = \min^* \$5$ . For every natural number  $n$  and for every element  $x$  of  $\mathbb{N}$  and for every element  $y$  of  $B$ , there exists an element  $x_1$  of  $\mathbb{N}$  and there exists an element  $y_1$  of  $B$  such that  $\mathcal{P}[n, x, y, x_1, y_1]$ . Consider  $N$  being a sequence of  $\mathbb{N}$ ,  $Y$  being a sequence of  $B$  such that  $N(0) = N_0$  and  $Y(0) = Y_0$  and for every natural number  $n$ ,  $\mathcal{P}[n, N(n), Y(n), N(n+1), Y(n+1)]$  from [13, Sch. 3]. Define  $\mathcal{Q}[\text{natural number}] \equiv N(\$1) = \min^*(Y(\$1))$  and  $N(\$1) \in Y(\$1)$  and  $Y(\$1)$  is infinite and  $Y(\$1) \subseteq \mathbb{N}$ . For every natural number  $i$  such that  $\mathcal{Q}[i]$  holds  $\mathcal{Q}[i+1]$  by [8, (31)]. For every natural number  $i$ ,  $\mathcal{Q}[i]$  from [1, Sch. 2].  $\text{rng } N \subseteq X$  by [7, (11)]. For every natural number  $i$ ,  $N(i) < N(i+1)$ .  $\square$

- (3) Let us consider a non empty metric space  $M$ , and a subset  $V$  of  $M_{\text{top}}$ . Suppose  $V$  is open. Then there exists a family  $F$  of subsets of  $M$  such that

- (i)  $F = \{\text{Ball}(x, r)$ , where  $x$  is an element of  $M$ ,  $r$  is a real number :  $r > 0$  and  $\text{Ball}(x, r) \subseteq V\}$ , and
- (ii)  $V = \bigcup F$ .

PROOF: Set  $F = \{\text{Ball}(x, r)$ , where  $x$  is an element of  $M$ ,  $r$  is a real number:  $r > 0$  and  $\text{Ball}(x, r) \subseteq V\}$ . For every object  $z$  such that  $z \in F$  holds  $z \in$  the open set family of  $M$  by [3, (29)]. Reconsider  $Q = \bigcup F$  as a subset of  $M_{\text{top}}$ . For every object  $z$ ,  $z \in V$  iff  $z \in Q$  by [9, (15)], [12, (1), (11)].  $\square$

- (4) Let us consider a non empty metric space  $M$ , a subset  $X$  of  $M_{\text{top}}$ , and an element  $p$  of  $M$ . Then  $p \in \overline{X}$  if and only if for every real number  $r$  such that  $0 < r$  holds  $X$  meets  $\text{Ball}(p, r)$ .
- (5) Let us consider a non empty metric space  $M$ , a subset  $X$  of  $M_{\text{top}}$ , and an object  $x$ . Then  $x \in \overline{X}$  if and only if there exists a sequence  $S$  of  $M$  such that for every natural number  $n$ ,  $S(n) \in X$  and  $S$  is convergent and  $\lim S = x$ .
- (6) Let us consider a non empty metric space  $M$ , and a subset  $X$  of  $M_{\text{top}}$ . Then  $X$  is closed if and only if for every sequence  $S$  of  $M$  such that for every natural number  $n$ ,  $S(n) \in X$  and  $S$  is convergent holds  $\lim S \in X$ . The theorem is a consequence of (5).

- (7) Let us consider non empty metric spaces  $X, Y$ , and a function  $f$  from  $X_{\text{top}}$  into  $Y_{\text{top}}$ . Then  $f$  is continuous if and only if for every sequence  $S$  of  $X$  and for every sequence  $T$  of  $Y$  such that  $S$  is convergent and  $T = f \cdot S$  holds  $T$  is convergent and  $\lim T = f(\lim S)$ .

PROOF: For every subset  $B$  of  $Y_{\text{top}}$  such that  $B$  is closed holds  $f^{-1}(B)$  is closed by [7, (15)], (6).  $\square$

## 2. COMPACTNESS IN METRIC SPACES

Let  $M$  be a non empty metric space and  $X$  be a subset of  $M$ . We say that  $X$  is sequentially compact if and only if

- (Def. 1) for every sequence  $S_1$  of  $M$  such that  $\text{rng } S_1 \subseteq X$  there exists a sequence  $S_2$  of  $M$  such that there exists an increasing sequence  $N$  of  $\mathbb{N}$  such that  $S_2 = S_1 \cdot N$  and  $S_2$  is convergent and  $\lim S_2 \in X$ .

Let us observe that every subset of  $M$  which is empty is also sequentially compact.

We say that  $M$  is sequentially compact if and only if

- (Def. 2)  $\Omega_M$  is sequentially compact.

Now we state the proposition:

- (8) Let us consider a non empty metric space  $M$ , a subset  $X$  of  $M$ , a subset  $Y$  of  $M_{\text{top}}$ , an element  $x$  of  $M$ , and an element  $y$  of  $M_{\text{top}}$ . Suppose  $X = Y$  and  $x = y$ . Then  $y$  is an accumulation point of  $Y$  if and only if for every real number  $r$  such that  $0 < r$  holds  $\text{Ball}(x, r)$  meets  $X \setminus \{x\}$ .

Let us consider a non empty metric space  $M$ . Now we state the propositions:

- (9) If  $M_{\text{top}}$  is countably-compact, then  $M$  is sequentially compact.

PROOF: For every subset  $X$  of  $M$  such that  $X$  is infinite there exists an element  $x$  of  $M$  such that for every real number  $r$  such that  $0 < r$  holds  $\text{Ball}(x, r)$  meets  $X \setminus \{x\}$  by [16, (28)], [11, (16)], (8). For every sequence  $x$  of  $M$  such that  $\text{rng } x \subseteq \Omega_M$  there exists a sequence  $x_1$  of  $M$  such that there exists an increasing sequence  $N$  of  $\mathbb{N}$  such that  $x_1 = x \cdot N$  and  $x_1$  is convergent and  $\lim x_1 \in \Omega_M$  by (1), (2), [7, (4), (38), (15)].  $\square$

- (10) If  $M$  is sequentially compact, then  $M_{\text{top}}$  is countably-compact.

PROOF: For every subset  $X$  of  $M$  such that  $X$  is infinite there exists an element  $x$  of  $M$  such that for every real number  $r$  such that  $0 < r$  holds  $\text{Ball}(x, r)$  meets  $X \setminus \{x\}$  by [15, (3)], [7, (2)], [19, (26)], [7, (112)]. For every subset  $A$  of  $M_{\text{top}}$  such that  $A$  is infinite holds  $\text{Der } A$  is not empty by (8), [11, (16)].  $\square$

- (11)  $M_{\text{top}}$  is compact if and only if  $M$  is sequentially compact. The theorem is a consequence of (9).

- (12)  $M$  is totally bounded and complete if and only if  $M$  is sequentially compact. The theorem is a consequence of (11).

Let us consider a non empty metric space  $M$  and a non empty subset  $S$  of  $M$ . Now we state the propositions:

- (13)  $S$  is sequentially compact if and only if  $M \upharpoonright S$  is sequentially compact.  
 PROOF: For every sequence  $S_1$  of  $M$  such that  $\text{rng } S_1 \subseteq S$  there exists a sequence  $S_2$  of  $M$  such that there exists an increasing sequence  $N$  of  $\mathbb{N}$  such that  $S_2 = S_1 \cdot N$  and  $S_2$  is convergent and  $\lim S_2 \in S$  by [7, (6)].  $\square$
- (14)  $S$  is sequentially compact if and only if  $M \upharpoonright S$  is compact. The theorem is a consequence of (11) and (13).
- (15) Let us consider a non empty metric space  $M$ , a subset  $S$  of  $M$ , and a subset  $T$  of  $M_{\text{top}}$ . If  $T = S$ , then  $T$  is compact iff  $S$  is sequentially compact. The theorem is a consequence of (11) and (13).
- (16) Let us consider a non empty metric space  $M$ , a non empty subset  $S$  of  $M$ , and a non empty subset  $T$  of  $M_{\text{top}}$ . Suppose  $T = S$ . Then  $M_{\text{top}} \upharpoonright T$  is countably-compact if and only if  $S$  is sequentially compact. The theorem is a consequence of (11) and (13).
- (17) Let us consider a non empty metric space  $M$ , and a non empty subset  $S$  of  $M$ . Then  $M \upharpoonright S$  is totally bounded and complete if and only if  $S$  is sequentially compact. The theorem is a consequence of (12) and (13).

### 3. COMPACTNESS IN NORM SPACES

Now we state the propositions:

- (18) Let us consider a real normed space  $N$ , a subset  $S$  of  $N$ , and a subset  $T$  of  $\text{MetricSpaceNorm } N$ . If  $S = T$ , then  $S$  is compact iff  $T$  is sequentially compact.
- (19) Let us consider a real normed space  $N$ , a subset  $S$  of  $N$ , and a subset  $T$  of  $\text{TopSpaceNorm } N$ . If  $S = T$ , then  $S$  is compact iff  $T$  is compact. The theorem is a consequence of (15) and (18).

### 4. TOPOLOGICAL PROPERTIES OF THE REAL LINE

Let us consider a sequence  $S_1$  of the metric space of real numbers, a sequence  $s$  of real numbers, a real number  $g$ , and an element  $g_1$  of the metric space of real numbers. Now we state the propositions:

- (20) Suppose  $S_1 = s$  and  $g = g_1$ . Then for every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number

$m$  such that  $n \leq m$  holds  $|s(m) - g| < p$  if and only if for every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $\rho(S_1(m), g_1) < p$ .

PROOF: For every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|s(m) - g| < p$  by [9, (11)].  $\square$

- (21) If  $S_1 = s$  and  $g = g_1$ , then  $s$  is convergent and  $\lim s = g$  iff  $S_1$  is convergent and  $\lim S_1 = g_1$ . The theorem is a consequence of (20).
- (22) Let us consider a sequence  $S_1$  of the metric space of real numbers, and a sequence  $s$  of real numbers. Suppose  $S_1 = s$  and  $s$  is convergent. Then
- (i)  $S_1$  is convergent, and
  - (ii)  $\lim S_1 = \lim s$ .

The theorem is a consequence of (20).

## 5. COMPACTNESS IN THE REAL LINE

Now we state the propositions:

- (23) Let us consider a subset  $N$  of  $\mathbb{R}$ , and a subset  $M$  of  $\mathbb{R}^1$ . Suppose  $N = M$ . Then for every family  $F$  of subsets of  $\mathbb{R}$  such that  $F$  is a cover of  $N$  and for every subset  $P$  of  $\mathbb{R}$  such that  $P \in F$  holds  $P$  is open there exists a family  $G$  of subsets of  $\mathbb{R}$  such that  $G \subseteq F$  and  $G$  is cover of  $N$  and finite if and only if for every family  $F_1$  of subsets of  $\mathbb{R}^1$  such that  $F_1$  is cover of  $M$  and open there exists a family  $G_1$  of subsets of  $\mathbb{R}^1$  such that  $G_1 \subseteq F_1$  and  $G_1$  is cover of  $M$  and finite.

PROOF: Reconsider  $F_1 = F$  as a family of subsets of  $\mathbb{R}^1$ . For every subset  $P_1$  of  $\mathbb{R}^1$  such that  $P_1 \in F_1$  holds  $P_1$  is open by [10, (39)]. Consider  $G_1$  being a family of subsets of  $\mathbb{R}^1$  such that  $G_1 \subseteq F_1$  and  $G_1$  is cover of  $M$  and finite.  $\square$

- (24) Let us consider a subset  $N$  of  $\mathbb{R}$ . Then  $N$  is compact if and only if for every family  $F$  of subsets of  $\mathbb{R}$  such that  $F$  is a cover of  $N$  and for every subset  $P$  of  $\mathbb{R}$  such that  $P \in F$  holds  $P$  is open there exists a family  $G$  of subsets of  $\mathbb{R}$  such that  $G \subseteq F$  and  $G$  is cover of  $N$  and finite. The theorem is a consequence of (23).

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [3] Leszek Borys. Paracompact and metrizable spaces. *Formalized Mathematics*, 2(4):481–485, 1991.
- [4] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [5] Nicolas Bourbaki, H.G. Eggleston, and S. Madan. *Elements of mathematics: Topological vector spaces*. Springer-Verlag, 1987.
- [6] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [7] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [8] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [9] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces – fundamental concepts. *Formalized Mathematics*, 2(4):605–608, 1991.
- [10] Adam Grabowski. On the subcontinua of a real line. *Formalized Mathematics*, 11(3):313–322, 2003.
- [11] Adam Grabowski. On the boundary and derivative of a set. *Formalized Mathematics*, 13(1):139–146, 2005.
- [12] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Formalized Mathematics*, 1(3):607–610, 1990.
- [13] Artur Korniłowicz. Recursive definitions. Part II. *Formalized Mathematics*, 12(2):167–172, 2004.
- [14] Erwin Kreyszig. *Introductory Functional Analysis with Applications*. Wiley, 1 edition, 1989.
- [15] Gilbert Lee and Piotr Rudnicki. Dickson’s lemma. *Formalized Mathematics*, 10(1):29–37, 2002.
- [16] Karol Pąk. Complete spaces. *Formalized Mathematics*, 16(1):35–43, 2008. doi:10.2478/v10037-008-0006-2.
- [17] Walter Rudin. *Functional Analysis*. New York, McGraw-Hill, 2nd edition, 1991.
- [18] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(3):495–500, 1990.
- [19] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [20] Kosaku Yoshida. *Functional Analysis*. Springer, 1980.

Received June 30, 2016

---



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.



# Double Sequences and Iterated Limits in Regular Space

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** First, we define in Mizar [5], the Cartesian product of two filters bases and the Cartesian product of two filters. After comparing the product of two Fréchet filters on  $\mathbb{N}$  ( $\mathcal{F}_1$ ) with the Fréchet filter on  $\mathbb{N} \times \mathbb{N}$  ( $\mathcal{F}_2$ ), we compare  $\lim_{\mathcal{F}_1}$  and  $\lim_{\mathcal{F}_2}$  for all double sequences in a non empty topological space.

Endou, Okazaki and Shidama formalized in [14] the “convergence in Pringsheim’s sense” for double sequence of real numbers. We show some basic correspondences between the  $p$ -convergence and the filter convergence in a topological space. Then we formalize that the double sequence  $(x_{m,n} = \frac{1}{m+1})_{(m,n)} \in \mathbb{N} \times \mathbb{N}$  converges in “Pringsheim’s sense” but not in Fréchet filter on  $\mathbb{N} \times \mathbb{N}$  sense.

In the next section, we generalize some definitions: “is convergent in the first coordinate”, “is convergent in the second coordinate”, “the  $\lim$  in the first coordinate of”, “the  $\lim$  in the second coordinate of” according to [14], in Hausdorff space.

Finally, we generalize two theorems: (3) and (4) from [14] in the case of double sequences and we formalize the “iterated limit” theorem (“Double limit” [7], p. 81, par. 8.5 “*Double limite*” [6] (TG I,57)), all in regular space. We were inspired by the exercises (2.11.4), (2.17.5) [17] and the corrections B.10 [18].

MSC: 54A20 40A05 40B05 03B35

Keywords: filter; double limits; Pringsheim convergence; iterated limits; regular space

MML identifier: CARDFIL4, version: 8.1.05 5.37.1275

## 1. PRELIMINARIES

From now on  $x$  denotes an object,  $X, Y, Z$  denote sets,  $i, j, k, l, m, n$  denote natural numbers,  $r, s$  denote real numbers,  $n_1$  denotes an element of the ordered  $\mathbb{N}$ , and  $A$  denotes a subset of  $\mathbb{N} \times \mathbb{N}$ .

Now we state the propositions:

- (1) Let us consider a finite subset  $W$  of  $X$ . If  $X \setminus W \subseteq Z$ , then  $X \setminus Z$  is finite.
- (2) If  $Z \subseteq X$  and  $X \setminus Z$  is finite, then there exists a finite subset  $W$  of  $X$  such that  $X \setminus W = Z$ .
- (3) Let us consider sets  $X_1, X_2$ , a family  $S_1$  of subsets of  $X_1$ , and a family  $S_2$  of subsets of  $X_2$ . Then  $\{s, \text{ where } s \text{ is a subset of } X_1 \times X_2 : \text{ there exist sets } s_1, s_2 \text{ such that } s_1 \in S_1 \text{ and } s_2 \in S_2 \text{ and } s = s_1 \times s_2\}$  is a family of subsets of  $X_1 \times X_2$ .
- (4) If  $x \in X \times Y$ , then  $x$  is pair.
- (5) If  $0 < r$ , then there exists  $m$  such that  $m$  is not zero and  $\frac{1}{m} < r$ .
- (6) Let us consider points  $x, y$  of the metric space of real numbers. Then there exist real numbers  $x_1, y_1$  such that
  - (i)  $x = x_1$ , and
  - (ii)  $y = y_1$ , and
  - (iii)  $\rho(x, y) = \rho_{\mathbb{R}}(x, y)$ , and
  - (iv)  $\rho(x, y) = \rho^1(\langle x \rangle, \langle y \rangle)$ , and
  - (v)  $\rho(x, y) = |x_1 - y_1|$ .
- (7) Let us consider points  $x, y$  of  $(\mathcal{E}^1)_{\text{top}}$ . Then there exist points  $x_2, y_2$  of the metric space of real numbers and there exist real numbers  $x_1, y_1$  such that  $x_2 = x_1$  and  $y_2 = y_1$  and  $x = \langle x_1 \rangle$  and  $y = \langle y_1 \rangle$  and  $\rho(x_2, y_2) = \rho_{\mathbb{R}}(x_1, y_1)$  and  $\rho(x_2, y_2) = \rho^1(\langle x_1 \rangle, \langle y_1 \rangle)$  and  $\rho(x_2, y_2) = |x_1 - y_1|$ .
- (8) Let us consider points  $x, y$  of  $\mathcal{E}^1$ , and real numbers  $r, s$ . If  $x = \langle r \rangle$  and  $y = \langle s \rangle$ , then  $\rho(x, y) = |r - s|$ . The theorem is a consequence of (7).

One can check that  $\mathbb{N} \times \mathbb{N}$  is countable and  $\mathbb{N} \times \mathbb{N}$  is denumerable.

Now we state the propositions:

- (9) the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number is infinite.  
 PROOF: Define  $\mathcal{F}(\text{object}) = \langle 0, \$1 \rangle$ . Consider  $f$  being a function such that  $\text{dom } f = \mathbb{N}$  and for every object  $x$  such that  $x \in \mathbb{N}$  holds  $f(x) = \mathcal{F}(x)$  from [9, Sch. 3].  $f$  is one-to-one.  $\text{rng } f =$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number by [9, (3)].  $\square$
- (10) If  $i \leq k$  and  $j \leq l$ , then  $\mathbb{Z}_i \times \mathbb{Z}_j \subseteq \mathbb{Z}_k \times \mathbb{Z}_l$ .
- (11)  $(\mathbb{N} \setminus \mathbb{Z}_m) \times (\mathbb{N} \setminus \mathbb{Z}_n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_m \times \mathbb{Z}_n$ .
- (12) If  $n = n_1$  and  $n \leq m$ , then  $m \in \uparrow n_1$ .
- (13) If  $n = n_1$  and  $m \in \uparrow n_1$ , then  $n \leq m$ .
- (14) If  $n = n_1$ , then  $\uparrow n_1 = \mathbb{N} \setminus \mathbb{Z}_n$ .

PROOF:  $\uparrow n_1 \subseteq \mathbb{N} \setminus \mathbb{Z}_n$  by [12, (50)], (13), [1, (44)].  $\mathbb{N} \setminus \mathbb{Z}_n \subseteq \uparrow n_1$  by [1, (44)], [12, (50)].  $\square$

- (15)  $\pi_1(A) = \{x, \text{ where } x \text{ is an element of } \mathbb{N} : \text{ there exists an element } y \text{ of } \mathbb{N} \text{ such that } \langle x, y \rangle \in A\}$ .
- (16)  $\pi_2(A) = \{y, \text{ where } y \text{ is an element of } \mathbb{N} : \text{ there exists an element } x \text{ of } \mathbb{N} \text{ such that } \langle x, y \rangle \in A\}$ .
- (17) Let us consider a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$ . Then there exists  $m$  and there exists  $n$  such that  $A \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ . The theorem is a consequence of (15) and (16).
- (18) Let us consider a non empty set  $X$ . Then every filter of  $X$  is a proper filter of  $2_{\subseteq}^X$ .
- (19) Let us consider a non empty set  $X$ , and a filter  $\mathcal{F}$  of  $X$ . Then there exists a filter base  $\mathcal{B}$  of  $X$  such that
  - (i)  $\mathcal{B} = \mathcal{F}$ , and
  - (ii)  $[\mathcal{B}] = \mathcal{F}$ .
- (20) Let us consider a non empty topological space  $T$ , and a filter  $\mathcal{F}$  of the carrier of  $T$ . If  $x \in \text{LimFilter}(\mathcal{F})$ , then  $x$  is a cluster point of  $\mathcal{F}, T$ .
- (21) Let us consider an element  $B$  of the base of Frechet filter. Then there exists  $n$  such that  $B = \mathbb{N} \setminus \mathbb{Z}_n$ . The theorem is a consequence of (14).
- (22) Let us consider a subset  $B$  of  $\mathbb{N}$ . Suppose  $B = \mathbb{N} \setminus \mathbb{Z}_n$ . Then  $B$  is an element of the base of Frechet filter. The theorem is a consequence of (14).

## 2. CARTESIAN PRODUCT OF TWO FILTERS

From now on  $X, Y, X_1, X_2$  denote non empty sets,  $\mathcal{A}_1, \mathcal{B}_1$  denote filter bases of  $X_1$ ,  $\mathcal{A}_2, \mathcal{B}_2$  denote filter bases of  $X_2$ ,  $\mathcal{F}_1$  denotes a filter of  $X_1$ ,  $\mathcal{F}_2$  denotes a filter of  $X_2$ ,  $\mathcal{B}_3$  denotes a generalized basis of  $\mathcal{F}_1$ .

Let  $X_1, X_2$  be non empty sets,  $\mathcal{B}_1$  be a filter base of  $X_1$ , and  $\mathcal{B}_2$  be a filter base of  $X_2$ . The functor  $\mathcal{B}_1 \times \mathcal{B}_2$  yielding a filter base of  $X_1 \times X_2$  is defined by the term

(Def. 1) the set of all  $B_1 \times B_2$  where  $B_1$  is an element of  $\mathcal{B}_1$ ,  $B_2$  is an element of  $\mathcal{B}_2$ .

Now we state the propositions:

- (23) Suppose  $\mathcal{F}_1 = [\mathcal{B}_1)$  and  $\mathcal{F}_1 = [\mathcal{A}_1)$  and  $\mathcal{F}_2 = [\mathcal{B}_2)$  and  $\mathcal{F}_2 = [\mathcal{A}_2)$ . Then  $[\mathcal{B}_1 \times \mathcal{B}_2) = [\mathcal{A}_1 \times \mathcal{A}_2)$ .
- (24) If  $\mathcal{B}_3 = \mathcal{B}_1$ , then  $[\mathcal{B}_1] = \mathcal{F}_1$ .

(25) There exists  $\mathcal{B}_1$  such that  $[\mathcal{B}_1] = \mathcal{F}_1$ . The theorem is a consequence of (24).

Let  $X_1, X_2$  be non empty sets,  $\mathcal{F}_1$  be a filter of  $X_1$ , and  $\mathcal{F}_2$  be a filter of  $X_2$ . The functor  $\langle \mathcal{F}_1, \mathcal{F}_2 \rangle$  yielding a filter of  $X_1 \times X_2$  is defined by

(Def. 2) there exists a filter base  $\mathcal{B}_1$  of  $X_1$  and there exists a filter base  $\mathcal{B}_2$  of  $X_2$  such that  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$  and  $it = [\mathcal{B}_1 \times \mathcal{B}_2]$ .

Let  $\mathcal{B}_1$  be a generalized basis of  $\mathcal{F}_1$  and  $\mathcal{B}_2$  be a generalized basis of  $\mathcal{F}_2$ . The functor  $\mathcal{B}_1 \times \mathcal{B}_2$  yielding a generalized basis of  $\langle \mathcal{F}_1, \mathcal{F}_2 \rangle$  is defined by

(Def. 3) there exists a filter base  $\mathcal{B}_3$  of  $X_1$  and there exists a filter base  $\mathcal{B}_4$  of  $X_2$  such that  $\mathcal{B}_1 = \mathcal{B}_3$  and  $\mathcal{B}_2 = \mathcal{B}_4$  and  $it = \mathcal{B}_3 \times \mathcal{B}_4$ .

Let  $n$  be a natural number. The functor  $\uparrow^2(n)$  yielding a subset of  $\mathbb{N} \times \mathbb{N}$  is defined by

(Def. 4) for every element  $x$  of  $\mathbb{N} \times \mathbb{N}$ ,  $x \in it$  iff there exist natural numbers  $n_1, n_2$  such that  $n_1 = (x)_1$  and  $n_2 = (x)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .

Now we state the proposition:

(26)  $\langle n, n \rangle \in \uparrow^2(n)$ .

Let us consider  $n$ . One can check that  $\uparrow^2(n)$  is non empty.

Now we state the propositions:

(27) If  $\langle i, j \rangle \in \uparrow^2(n)$ , then  $\langle i + k, j \rangle, \langle i, j + l \rangle \in \uparrow^2(n)$ .

(28)  $\uparrow^2(n)$  is an infinite subset of  $\mathbb{N} \times \mathbb{N}$ . The theorem is a consequence of (17).

(29) If  $n_1 = n$ , then  $\uparrow^2(n) = \uparrow n_1 \times \uparrow n_1$ . The theorem is a consequence of (12) and (13).

(30) If  $m = n - 1$ , then  $\uparrow^2(n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \text{Seg } m \times \text{Seg } m$ .

PROOF: Reconsider  $y = x$  as an element of  $\mathbb{N} \times \mathbb{N}$ . Consider  $n_1, n_2$  being natural numbers such that  $n_1 = (y)_1$  and  $n_2 = (y)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .  $x \notin \text{Seg } m \times \text{Seg } m$  by [3, (1)].  $\square$

(31)  $\uparrow^2(n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_n \times \mathbb{Z}_n$ .

PROOF: Reconsider  $y = x$  as an element of  $\mathbb{N} \times \mathbb{N}$ . Consider  $n_1, n_2$  being natural numbers such that  $n_1 = (y)_1$  and  $n_2 = (y)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .  $x \notin \mathbb{Z}_n \times \mathbb{Z}_n$  by [16, (10)].  $\square$

(32)  $\uparrow^2(n) = (\mathbb{N} \setminus \mathbb{Z}_n) \times (\mathbb{N} \setminus \mathbb{Z}_n)$ . The theorem is a consequence of (14) and (29).

(33) There exists  $n$  such that  $\uparrow^2(n) \subseteq (\mathbb{N} \setminus \mathbb{Z}_i) \times (\mathbb{N} \setminus \mathbb{Z}_j)$ . The theorem is a consequence of (4).

(34) If  $n = \max(i, j)$ , then  $\uparrow^2(n) \subseteq (\uparrow^2(i)) \cap (\uparrow^2(j))$ .

Let  $n$  be a natural number. The functor  $\downarrow^2(n)$  yielding a subset of  $\mathbb{N} \times \mathbb{N}$  is defined by

(Def. 5) for every element  $x$  of  $\mathbb{N} \times \mathbb{N}$ ,  $x \in it$  iff there exist natural numbers  $n_1, n_2$  such that  $n_1 = (x)_1$  and  $n_2 = (x)_2$  and  $n_1 < n$  and  $n_2 < n$ .

Now we state the propositions:

(35)  $\downarrow^2(n) = \mathbb{Z}_n \times \mathbb{Z}_n$ .

PROOF:  $\downarrow^2(n) \subseteq \mathbb{Z}_n \times \mathbb{Z}_n$  by [1, (44)]. Consider  $y_2, y_1$  being objects such that  $y_2 \in \mathbb{Z}_n$  and  $y_1 \in \mathbb{Z}_n$  and  $x = \langle y_2, y_1 \rangle$ .  $\square$

(36) Let us consider a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$ . Then there exists  $n$  such that  $A \subseteq \downarrow^2(n)$ .

PROOF: Consider  $m, n$  such that  $A \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ . Reconsider  $m_1 = \max(m, n)$  as a natural number.  $A \subseteq \downarrow^2(m_1)$  by [1, (39)], [11, (96)], (35).  $\square$

(37)  $\downarrow^2(n)$  is a finite subset of  $\mathbb{N} \times \mathbb{N}$ . The theorem is a consequence of (35).

### 3. COMPARISON BETWEEN CARTESIAN PRODUCT OF FRECHET FILTER ON $\mathbb{N}$ AND THE FRECHET FILTER OF $\mathbb{N} \times \mathbb{N}$

Let us consider an element  $x$  of (the base of Frechet filter)  $\times$  (the base of Frechet filter). Now we state the propositions:

(38) There exists  $i$  and there exists  $j$  such that  $x = (\mathbb{N} \setminus \mathbb{Z}_i) \times (\mathbb{N} \setminus \mathbb{Z}_j)$ . The theorem is a consequence of (21).

(39) There exists  $n$  such that  $\uparrow^2(n) \subseteq x$ . The theorem is a consequence of (38) and (33).

(40) (The base of Frechet filter)  $\times$  (the base of Frechet filter) is a filter base of  $\mathbb{N} \times \mathbb{N}$ .

(41) There exists a generalized basis  $\mathcal{B}$  of  $\text{FrechetFilter}(\mathbb{N})$  such that

(i)  $\mathcal{B} =$  the base of Frechet filter, and

(ii)  $\mathcal{B} \times \mathcal{B}$  is a generalized basis of  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

The functor  $\uparrow_{\mathbb{N}}^2$  yielding a filter base of  $\mathbb{N} \times \mathbb{N}$  is defined by the term

(Def. 6) the set of all  $\uparrow^2(n)$  where  $n$  is a natural number.

Now we state the propositions:

(42)  $\uparrow_{\mathbb{N}}^2$  and (the base of Frechet filter)  $\times$  (the base of Frechet filter) are equivalent generators. The theorem is a consequence of (22), (32), and (39).

(43)  $[(\text{the base of Frechet filter}) \times (\text{the base of Frechet filter})] = \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ . The theorem is a consequence of (41).

(44)  $[\uparrow_{\mathbb{N}}^2] = \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

(45)  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  is finer than  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$ .  
 The theorem is a consequence of (17), (11), (22), and (43).

(46) (i)  $\mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number  $\in \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ , and

(ii)  $\mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number  $\notin \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$ .

PROOF: Set  $X = \mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number.  $\uparrow^2(1) \subseteq X$  by (32), [1, (44)].  $X \notin \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  by [12, (51)], [15, (5)], (9).  $\square$

(47)  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N}) \neq \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

#### 4. TOPOLOGICAL SPACE AND DOUBLE SEQUENCE

In the sequel  $T$  denotes a non empty topological space,  $s$  denotes a function from  $\mathbb{N} \times \mathbb{N}$  into the carrier of  $T$ ,  $M$  denotes a subset of the carrier of  $T$ , and  $\mathcal{F}_1, \mathcal{F}_2$  denote filters of the carrier of  $T$ . Now we state the propositions:

(48) If  $\mathcal{F}_2$  is finer than  $\mathcal{F}_1$ , then  $\text{LimFilter}(\mathcal{F}_1) \subseteq \text{LimFilter}(\mathcal{F}_2)$ .

(49) Let us consider a function  $f$  from  $X$  into  $Y$ , and filters  $\mathcal{F}_1, \mathcal{F}_2$  of  $X$ . Suppose  $\mathcal{F}_2$  is finer than  $\mathcal{F}_1$ . Then the image of filter  $\mathcal{F}_2$  under  $f$  is finer than the image of filter  $\mathcal{F}_1$  under  $f$ .

(50)  $s^{-1}(M) \in \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  if and only if there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(M) = \mathbb{N} \times \mathbb{N} \setminus A$ .

(51)  $s^{-1}(M) \in \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  if and only if there exists  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(M)$ . The theorem is a consequence of (43), (39), and (42).

(52) The image of filter  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  under  $s = \{M$ , where  $M$  is a subset of the carrier of  $T$  : there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(M) = \mathbb{N} \times \mathbb{N} \setminus A\}$ . The theorem is a consequence of (50).

(53) The image of filter  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  under  $s = \{M$ , where  $M$  is a subset of the carrier of  $T$  : there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(M)\}$ . The theorem is a consequence of (51).

Let us consider a point  $x$  of  $T$ . Now we state the propositions:

(54)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every neighbourhood  $A$  of  $x$ , there exists a finite subset  $B$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(A) = \mathbb{N} \times \mathbb{N} \setminus B$ . The theorem is a consequence of (52).

(55)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every neighbourhood  $A$  of  $x$ ,  $\mathbb{N} \times \mathbb{N} \setminus s^{-1}(A)$  is finite. The theorem is a consequence of (54), (1), and (2).

- (56)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every neighbourhood  $A$  of  $x$ , there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(A)$ . The theorem is a consequence of (53).

Let us consider a point  $x$  of  $T$  and a generalized basis  $\mathcal{B}$  of BooleanFilter ToFilter(the neighborhood system of  $x$ ). Now we state the propositions:

- (57)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(B)$ . The theorem is a consequence of (56).
- (58)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(B) = \mathbb{N} \times \mathbb{N} \setminus A$ . The theorem is a consequence of (54), (1), and (55).
- (59)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that  $s^\circ(\uparrow^2(n)) \subseteq B$ . The theorem is a consequence of (57).

- (60)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^\circ(\mathbb{N} \times \mathbb{N} \setminus A) \subseteq B$ .

PROOF: For every neighbourhood  $A$  of  $x$ ,  $\mathbb{N} \times \mathbb{N} \setminus s^{-1}(A)$  is finite by [4, (2)], [19, (143)], [9, (76)].  $\square$

- (61)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists  $n$  and there exists  $m$  such that  $s^\circ(\mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_n \times \mathbb{Z}_m) \subseteq B$ . The theorem is a consequence of (60) and (17).

- (62)  $x \in s^\circ(\uparrow^2(n))$  if and only if there exists  $i$  and there exists  $j$  such that  $n \leq i$  and  $n \leq j$  and  $x = s(i, j)$ .

- (63)  $x \in s^\circ(\mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_i \times \mathbb{Z}_j)$  if and only if there exist natural numbers  $n, m$  such that  $(i \leq n$  or  $j \leq m)$  and  $x = s(n, m)$ .

PROOF: Consider  $n, m$  being natural numbers such that  $i \leq n$  or  $j \leq m$  and  $x = s(n, m)$ .  $\langle n, m \rangle \notin \mathbb{Z}_i \times \mathbb{Z}_j$  by [1, (44)].  $\square$

Let us consider a point  $x$  of  $T$  and a generalized basis  $\mathcal{B}$  of BooleanFilter ToFilter(the neighborhood system of  $x$ ). Now we state the propositions:

- (64)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in B$ . The theorem is a consequence of (62) and (59).

- (65)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists  $i$  and there exists  $j$  such that for every  $m$  and  $n$  such that  $i \leq m$  or  $j \leq n$  holds  $s(m, n) \in B$ . The theorem is a consequence of (61).

- (66)  $\lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s \subseteq \lim_{[\uparrow^2_{\mathbb{N}}]} s$ . The theorem is a consequence of (42), (43), (45), (48), and (49).

## 5. METRIC SPACE AND DOUBLE SEQUENCE

Now we state the propositions:

- (67) Let us consider a non empty metric space  $M$ , a point  $p$  of  $M$ , a point  $x$  of  $M_{\text{top}}$ , and a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $M_{\text{top}}$ . Suppose  $x = p$ . Then  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$ .

PROOF:  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  iff for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$  by [13, (6)], (64).  $\square$

- (68) Let us consider a non empty metric space  $M$ , a point  $p$  of  $M$ , a point  $x$  of  $M_{\text{top}}$ , a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $M_{\text{top}}$ , and a function  $s_2$  from  $\mathbb{N} \times \mathbb{N}$  into  $M$ . Suppose  $x = p$  and  $s = s_2$ . Then  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s_2(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$ .

## 6. ONE-DIMENSIONAL EUCLIDEAN METRIC SPACE AND DOUBLE SEQUENCE

In the sequel  $R$  denotes a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ .

Now we state the proposition:

- (69) Let us consider a point  $x$  of  $(\mathcal{E}^1)_{\text{top}}$ , a point  $y$  of  $\mathcal{E}^1$ , a generalized basis  $\mathcal{B}$  of  $\text{BooleanFilterToFilter}(\text{the neighborhood system of } x)$ , and an element  $b$  of  $\mathcal{B}$ . Suppose  $x = y$  and  $\mathcal{B} = \text{Balls } x$ . Then there exists a natural number  $n$  such that  $b = \{q, \text{ where } q \text{ is an element of } \mathcal{E}^1 : \rho(y, q) < \frac{1}{n}\}$ .

Let  $s$  be a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ . The functor  $\# s$  yielding a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}^1$  is defined by the term

(Def. 7)  $s$ .

Now we state the propositions:

- (70) Let us consider a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $(\mathcal{E}^1)_{\text{top}}$ , and a point  $y$  of  $\mathcal{E}^1$ . Then  $s^\circ(\uparrow^2(n)) \subseteq \{q, \text{ where } q \text{ is an element of } \mathcal{E}^1 : \rho(y, q) < \frac{1}{m}\}$  if and only if for every object  $x$  such that  $x \in s^\circ(\uparrow^2(n))$  there exist real numbers  $r_1, r_2$  such that  $x = \langle r_1 \rangle$  and  $y = \langle r_2 \rangle$  and  $|r_2 - r_1| < \frac{1}{m}$ . The theorem is a consequence of (8).

- (71)  $r \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every



natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|R(n_1, n_2) - r| < \frac{1}{m}$ .

PROOF: Reconsider  $p = r$  as a point of the metric space of real numbers. for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $R(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of the metric space of real numbers} : \rho(p, q) < \frac{1}{m}\}$  iff for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|R(n_1, n_2) - r| < \frac{1}{m}$  by (6), [8, (60)].  $\square$

## 7. BASIC RELATIONS CONVERGENCE IN PRINGSHEIM'S SENSE AND FILTER CONVERGENCE

Now we state the propositions:

- (72) Suppose  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R \neq \emptyset$ . Then there exists a real number  $x$  such that  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R = \{x\}$ .
- (73) If  $R$  is P-convergent, then  $\text{P-lim } R \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ . The theorem is a consequence of (71).
- (74)  $R$  is P-convergent if and only if  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R \neq \emptyset$ . The theorem is a consequence of (71) and (5).
- (75) Suppose  $R$  is P-convergent. Then  $\{\text{P-lim } R\} = \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ . The theorem is a consequence of (73) and (72).
- (76) Suppose  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$  is not empty. Then
- (i)  $R$  is P-convergent, and
  - (ii)  $\{\text{P-lim } R\} = \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ .

## 8. EXAMPLE: DOUBLE SEQUENCE CONVERGES IN PRINGSHEIM'S SENSE BUT NOT IN FRECHET FILTER OF $\mathbb{N} \times \mathbb{N}$ SENSE

The functor  $\text{DbSeq-ex1}$  yielding a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$  is defined by (Def. 8) for every natural numbers  $m, n$ ,  $it(m, n) = \frac{1}{m+1}$ .

Now we state the propositions:

- (77) Let us consider a non zero natural number  $m$ . Then there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|(\text{DbSeq-ex1})(n_1, n_2) - 0| < \frac{1}{m}$ .
- (78)  $0 \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# \text{DbSeq-ex1}$ .

- (79)  $\lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} \# \text{DblSeq-ex1} = \emptyset$ . The theorem is a consequence of (66), (42), (43), (72), (78), and (65).
- (80)  $\lim_{(\text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}))} \# \text{DblSeq-ex1} \neq \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} \# \text{DblSeq-ex1}$ .

9. CORRESPONDENCE WITH SOME DEFINITIONS FROM [14]

Let  $X_1, X_2$  be non empty sets,  $\mathcal{F}_1$  be a filter of  $X_1$ ,  $Y$  be a Hausdorff, non empty topological space, and  $f$  be a function from  $X_1 \times X_2$  into  $Y$ . Assume for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . The functor  $\lim_1(f, \mathcal{F}_1)$  yielding a function from  $X_2$  into  $Y$  is defined by

(Def. 9) for every element  $x$  of  $X_2$ ,  $\{it(x)\} = \lim_{\mathcal{F}_1} \text{curry}'(f, x)$ .

Let  $\mathcal{F}_2$  be a filter of  $X_2$ . Assume for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . The functor  $\lim_2(f, \mathcal{F}_2)$  yielding a function from  $X_1$  into  $Y$  is defined by

(Def. 10) for every element  $x$  of  $X_1$ ,  $\{it(x)\} = \lim_{\mathcal{F}_2} \text{curry}(f, x)$ .

Now we state the propositions:

- (81) Every function from  $X$  into  $\mathbb{R}$  is a function from  $X$  into  $\mathbb{R}^1$ .
- (82) Every sequence of  $\mathbb{R}$  is a function from  $\mathbb{N}$  into  $\mathbb{R}^1$ .

From now on  $f$  denotes a function from  $\Omega_{\text{the ordered } \mathbb{N}}$  into  $\mathbb{R}^1$  and  $s_1$  denotes a function from  $\mathbb{N}$  into  $\mathbb{R}$ .

Now we state the propositions:

- (83) Suppose  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ . Then
  - (i)  $s_1$  is convergent, and
  - (ii) there exists a real number  $z$  such that  $z \in \text{LimF}(f)$  and for every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|s_1(m) - z| < p$ .

PROOF: Consider  $x$  being an object such that  $x \in \text{LimF}(f)$ . Reconsider  $y = x$  as a point of (the metric space of real numbers)<sub>top</sub>. Reconsider  $z = y$  as a real number. Consider  $y_1$  being a point of the metric space of real numbers such that  $y_1 = y$  and  $\text{Balls } y = \{\text{Ball}(y_1, \frac{1}{n}), \text{ where } n \text{ is a natural number : } n \neq 0\}$ . For every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|s_1(m) - z| < p$  by (5), [12, (84), (50)], [2, (18)].  $\square$

- (84) If  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ , then  $\text{LimF}(f) = \{\lim s_1\}$ .

PROOF: Consider  $x$  being an object such that  $x \in \text{LimF}(f)$ . Consider  $u$  being an object such that  $\text{LimF}(f) = \{u\}$ .  $\text{LimF}(f) = \{\lim s_1\}$  by (83), [11, (3)].  $\square$

- (85) Let us consider a function  $f$  from  $\Omega_\alpha$  into  $T$ , and a sequence  $s$  of  $T$ . If  $f = s$ , then  $\text{LimF}(f) = \text{LimF}(s)$ , where  $\alpha$  is the ordered  $\mathbb{N}$ .
- (86) Let us consider a function  $f$  from  $\Omega_\alpha$  into  $T$ , and a function  $g$  from  $\mathbb{N}$  into  $T$ . If  $f = g$ , then  $\text{LimF}(f) = \text{LimF}(g)$ , where  $\alpha$  is the ordered  $\mathbb{N}$ .
- (87) Let us consider a function  $f$  from  $\mathbb{N}$  into  $\mathbb{R}^1$ . Suppose  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ . Then  $\text{LimF}(f) = \{\lim s_1\}$ . The theorem is a consequence of (84).
- (88) for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(\# R, x) \neq \emptyset$  if and only if  $R$  is convergent in the first coordinate. The theorem is a consequence of (5).
- (89) for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(\# R, x) \neq \emptyset$  if and only if  $R$  is convergent in the second coordinate. The theorem is a consequence of (5).

Let us consider an element  $t$  of  $\mathbb{N}$ , a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}^1$ , and a function  $s_1$  from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ . Now we state the propositions:

- (90) Suppose  $f = s_1$  and for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, t) = \{\lim \text{curry}(s_1, t)\}$ . The theorem is a consequence of (87).
- (91) Suppose  $f = s_1$  and for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, t) = \{\lim \text{curry}'(s_1, t)\}$ . The theorem is a consequence of (87).
- (92) Let us consider a Hausdorff, non empty topological space  $Y$ , and a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $Y$ . Suppose for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, x) \neq \emptyset$  and  $f = R$  and  $Y = \mathbb{R}^1$ . Then  $\lim_1(f, \text{FrechetFilter}(\mathbb{N})) =$  the lim in the first coordinate of  $R$ . The theorem is a consequence of (91).
- (93) Let us consider a non empty, Hausdorff topological space  $Y$ , and a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $Y$ . Suppose for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, x) \neq \emptyset$  and  $f = R$  and  $Y = \mathbb{R}^1$ . Then  $\lim_2(f, \text{FrechetFilter}(\mathbb{N})) =$  the lim in the second coordinate of  $R$ . The theorem is a consequence of (90).

## 10. REGULAR SPACE, DOUBLE LIMIT AND ITERATED LIMIT

From now on  $Y$  denotes a non empty topological space,  $x$  denotes a point of  $Y$ , and  $f$  denotes a function from  $X_1 \times X_2$  into  $Y$ .

Now we state the proposition:

- (94) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Let us consider a subset  $V$  of  $Y$ . Suppose  $V$  is open and  $x \in V$ . Then there exists an ele-

ment  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq V$ .

Let us consider a neighbourhood  $U$  of  $x$ . Now we state the propositions:

- (95) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ .
- (96) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $y$  of  $B_1$ ,  $f^\circ(\{y\} \times B_2) \subseteq \text{Int } U$ . The theorem is a consequence of (95).
- (97) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $z$  of  $X_1$  for every element  $y$  of  $Y$  such that  $z \in B_1$  and  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds  $y \in \overline{\text{Int } U}$ .

PROOF: Consider  $B_1$  being an element of  $\mathcal{B}_1$ ,  $B_2$  being an element of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ . For every element  $y$  of  $B_1$ ,  $f^\circ(\{y\} \times B_2) \subseteq \text{Int } U$  by [11, (95)], [19, (125)]. For every element  $z$  of  $B_1$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$  is a proper filter of  $2_{\subseteq}^{\Omega_Y}$  and  $\text{Int } U \in$  the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$  and  $y$  is a cluster point of the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$ ,  $Y$  by (18), [19, (132)], [10, (95)], (20). For every element  $z$  of  $B_1$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds  $y \in \overline{\text{Int } U}$  by [4, (25)].  $\square$

- (98) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $z$  of  $X_2$  for every element  $y$  of  $Y$  such that  $z \in B_2$  and  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds  $y \in \overline{\text{Int } U}$ .

PROOF: Consider  $B_1$  being an element of  $\mathcal{B}_1$ ,  $B_2$  being an element of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ . For every element  $y$  of  $B_2$ ,  $f^\circ(B_1 \times \{y\}) \subseteq \text{Int } U$  by [11, (95)], [19, (125)]. For every element  $z$  of  $B_2$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$  is a proper filter of  $2_{\subseteq}^{\Omega_Y}$  and  $\text{Int } U \in$  the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$  and  $y$  is a cluster point of the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$ ,  $Y$  by (18), [19, (132)], [10, (95)], (20). For every element  $z$  of  $B_2$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds  $y \in \overline{\text{Int } U}$  by [4, (25)].  $\square$

Let us consider a Hausdorff, regular, non empty topological space  $Y$  and a function  $f$  from  $X_1 \times X_2$  into  $Y$ . Now we state the propositions:

- (99) Suppose for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle}$

$f \subseteq \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (19) and (98).

- (100) Suppose for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \subseteq \lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2)$ . The theorem is a consequence of (19) and (97).

Let us consider non empty sets  $X_1$ ,  $X_2$ , a filter  $\mathcal{F}_1$  of  $X_1$ , a filter  $\mathcal{F}_2$  of  $X_2$ , a Hausdorff, regular, non empty topological space  $Y$ , and a function  $f$  from  $X_1 \times X_2$  into  $Y$ . Now we state the propositions:

- (101) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f = \lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2)$ . The theorem is a consequence of (100).
- (102) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f = \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (99).
- (103) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$  and for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2) = \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (102) and (101).

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Formalized Mathematics*, 6(1):93–107, 1997.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Grzegorz Bancerek, Noboru Endou, and Yuji Sakai. On the characterizations of compactness. *Formalized Mathematics*, 9(4):733–738, 2001.
- [5] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [6] Nicolas Bourbaki. *Topologie générale: Chapitres 1 à 4*. Eléments de mathématique. Springer Science & Business Media, 2007.
- [7] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [8] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Roland Coghetto. Convergent filter bases. *Formalized Mathematics*, 23(3):189–203, 2015. doi:10.1515/forma-2015-0016.
- [13] Roland Coghetto. Summable family in a commutative group. *Formalized Mathematics*, 23(4):279–288, 2015. doi:10.1515/forma-2015-0022.

- [14] Noboru Endou, Hiroyuki Okazaki, and Yasunari Shidama. Double sequences and limits. *Formalized Mathematics*, 21(3):163–170, 2013. doi:10.2478/forma-2013-0018.
- [15] Andrzej Owsiejczuk. Combinatorial Grassmannians. *Formalized Mathematics*, 15(2):27–33, 2007. doi:10.2478/v10037-007-0004-9.
- [16] Karol Pałk. Stirling numbers of the second kind. *Formalized Mathematics*, 13(2):337–345, 2005.
- [17] Claude Wagschal. *Topologie et analyse fonctionnelle*. Hermann, 1995.
- [18] Claude Wagschal. *Topologie: Exercices et problèmes corrigés*. Hermann, 1995.
- [19] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

*Received June 30, 2016*

---



The English version of this volume of *Formalized Mathematics* was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.

# Prime Factorization of Sums and Differences of Two Like Powers

Rafał Ziobro  
Department of Carbohydrate Technology  
University of Agriculture  
Krakow, Poland

**Summary.** Representation of a non zero integer as a signed product of primes is unique similarly to its representations in various types of positional notations [4], [3]. The study focuses on counting the prime factors of integers in the form of sums or differences of two equal powers (thus being represented by 1 and a series of zeroes in respective digital bases).

Although the introduced theorems are not particularly important, they provide a couple of shortcuts useful for integer factorization, which could serve in further development of Mizar projects [2]. This could be regarded as one of the important benefits of proof formalization [9].

MSC: 11A51 03B35

Keywords: integers; factorization; primes

MML identifier: NEWTON03, version: 8.1.05 5.37.1275

From now on  $a, b, c, d, x, j, k, l, m, n, o$  denote natural numbers,  $p, q, t, z, u, v$  denote integers, and  $a_1, b_1, c_1, d_1$  denote complexes.

Now we state the propositions:

- (1)  $a_1^{n+k} + b_1^{n+k} = a_1^n \cdot (a_1^k + b_1^k) + b_1^k \cdot (b_1^n - a_1^n)$ .
- (2)  $a_1^{n+k} - b_1^{n+k} = a_1^n \cdot (a_1^k - b_1^k) + b_1^k \cdot (a_1^n - b_1^n)$ .
- (3)  $a_1^{m+2} + b_1^{m+2} = (a_1 + b_1) \cdot (a_1^{m+1} + b_1^{m+1}) - a_1 \cdot b_1 \cdot (a_1^m + b_1^m)$ .

Let  $a$  be a natural number. Let us note that  $a$  is trivial if and only if the condition (Def. 1) is satisfied.

(Def. 1)  $a \leq 1$ .

Let  $a$  be a complex. Let us note that the functor  $a^2$  yields a set and is defined by the term

(Def. 2)  $a^2$ .

Let  $a, b$  be integers. The functors:  $\gcd(a, b)$  and  $\text{lcm}(a, b)$  yielding natural numbers are defined by terms

(Def. 3)  $\gcd(|a|, |b|)$ ,

(Def. 4)  $\text{lcm}(|a|, |b|)$ ,

respectively. Let  $a, b$  be positive real numbers. Note that  $\max(a, b)$  is positive and  $\min(a, b)$  is positive.

Let  $a$  be a non zero integer and  $b$  be an integer. One can check that  $\gcd(a, b)$  is non zero.

Let  $a$  be a non zero complex and  $n$  be a natural number. Let us observe that  $a^n$  is non zero.

Let  $a$  be a non trivial natural number and  $n$  be a non zero natural number. Note that  $a^n$  is non trivial.

Let  $a$  be an integer. One can check that  $|a|$  is natural.

Let  $a$  be an even integer. Note that  $|a|$  is even.

Let  $a$  be a natural number. Let us note that  $\text{lcm}(a, a)$  reduces to  $a$  and  $\gcd(a, a)$  reduces to  $a$ .

Let  $a$  be a non zero integer and  $b$  be an integer. Note that  $\gcd(a, b)$  is positive.

Let  $a, b$  be integers. One can check that  $\gcd(a, \gcd(a, b))$  reduces to  $\gcd(a, b)$  and  $\text{lcm}(a, \text{lcm}(a, b))$  reduces to  $\text{lcm}(a, b)$ .

Let  $a$  be an integer. Observe that  $\gcd(a, 1)$  reduces to 1 and  $\gcd(a + 1, a)$  reduces to 1.

Now we state the proposition:

(4) Let us consider integers  $t, z$ . Then  $\gcd(t^n, z^n) = (\gcd(t, z))^n$ .

Let  $a$  be an integer and  $n$  be a natural number.

One can verify that  $\gcd((a + 1)^n, a^n)$  reduces to 1.

Let us consider  $a_1$  and  $b_1$ . One can verify that  $a_1^0 - b_1^0$  reduces to 0.

Let  $a$  be a non negative real number and  $n$  be a natural number. One can verify that  $a^n$  is non negative and there exists an odd natural number which is non trivial and there exists an even natural number which is non trivial.

Let  $a$  be a positive real number and  $n$  be a natural number. One can verify that  $a^n$  is positive.

Let  $a$  be an integer. One can verify that  $a \cdot a$  is square and  $\frac{a}{a}$  is square and there exists an element of  $\mathbb{N}$  which is non square and every element of  $\mathbb{N}$  which is prime is also non square and there exists a prime natural number which is even and there exists a prime natural number which is odd and every integer which is prime is also non square.

Let  $a$  be a square element of  $\mathbb{N}$ . Observe that  $\sqrt{a}$  is natural.



Let  $a$  be an integer. Let us note that  $a^2$  is square and  $a \cdot a$  is square and there exists an integer which is non square and every natural number which is zero is also trivial and there exists a natural number which is square and there exists an element of  $\mathbb{N}$  which is non zero and there exists a square element of  $\mathbb{N}$  which is non trivial and every natural number which is trivial is also square and every integer which is non square is also non zero.

Now we state the propositions:

(5) Let us consider integers  $a, b, c, d$ . If  $a \mid b$  and  $c \mid d$ , then  $a \cdot c \mid b \cdot d$ .

(6) Let us consider integers  $a, b$ . Then  $a \mid b$  if and only if  $\text{lcm}(a, b) = |b|$ .

PROOF: If  $a \mid b$ , then  $\text{lcm}(a, b) = |b|$  by [8, (16)], [7, (44)].  $\square$

Let  $a$  be an integer. Observe that  $\text{lcm}(a, 0)$  reduces to 0.

Let  $a$  be a natural number. Note that  $\text{lcm}(a, 1)$  reduces to  $a$ .

Let us consider  $a$  and  $b$ . Let us observe that  $\text{lcm}(a \cdot b, a)$  reduces to  $a \cdot b$  and  $\text{lcm}(\text{gcd}(a, b), b)$  reduces to  $b$  and  $\text{gcd}(a, \text{lcm}(a, b))$  reduces to  $a$ .

Let us consider integers  $a, b$ . Now we state the propositions:

(7)  $|a \cdot b| = (\text{gcd}(a, b)) \cdot \text{lcm}(a, b)$ .

(8)  $\text{lcm}(a^n, b^n) = \text{lcm}(a, b)^n$ . The theorem is a consequence of (4) and (7).

Let  $a$  be a square element of  $\mathbb{N}$  and  $b$  be a square element of  $\mathbb{N}$ . One can check that  $\text{gcd}(a, b)$  is square and  $\text{lcm}(a, b)$  is square.

Let  $a, b$  be square integers. One can verify that  $\text{gcd}(a, b)$  is square and  $\text{lcm}(a, b)$  is square.

Now we state the proposition:

(9) Let us consider an integer  $t$ . Then  $t$  is odd if and only if  $\text{gcd}(t, 2) = 1$ .

PROOF: If  $t$  is odd, then  $\text{gcd}(t, 2) = 1$  by [13, (1)], [14, (5)].  $\square$

Let  $t$  be an integer. One can check that  $t$  is odd if and only if the condition (Def. 5) is satisfied.

(Def. 5)  $\text{gcd}(t, 2) = 1$ .

Let  $a$  be an odd integer. Let us observe that  $|a|$  is odd and  $-a$  is odd.

Let  $a, b$  be even integers. Note that  $\text{gcd}(a, b)$  is even.

Let  $a$  be an integer and  $b$  be an odd integer. Note that  $\text{gcd}(a, b)$  is odd.

Let  $a$  be a natural number. One can check that  $|-a|$  reduces to  $a$ .

Let  $t, z$  be even integers. One can check that  $t + z$  is even and  $t - z$  is even and  $t \cdot z$  is even.

Let  $t, z$  be odd integers. Note that  $t + z$  is even and  $t - z$  is even and  $t \cdot z$  is odd.

Let  $t$  be an odd integer and  $z$  be an even integer. Let us observe that  $t + z$  is odd and  $t - z$  is odd and  $t \cdot z$  is even.

Now we state the proposition:

(10) Let us consider a non zero, square integer  $a$ , and an integer  $b$ . If  $a \cdot b$  is square, then  $b$  is square.

Let  $a$  be a square element of  $\mathbb{N}$  and  $n$  be a natural number. Let us observe that  $a^n$  is square.

Let  $a$  be a square integer. Note that  $a^n$  is square.

Let  $a$  be a non zero, square integer and  $b$  be a non square integer. Let us note that  $a \cdot b$  is non square.

Let  $a$  be an element of  $\mathbb{N}$  and  $b$  be an even natural number. Note that  $a^b$  is square.

Let  $a$  be a non square element of  $\mathbb{N}$  and  $b$  be an odd natural number. Note that  $a^b$  is non square.

Let  $a$  be a non zero, square integer. Note that  $a + 1$  is non square.

Let  $a$  be a non zero, square element of  $\mathbb{N}$ . Let us observe that  $a + 1$  is non square.

Let  $a$  be a non zero, square object and  $b$  be a non square element of  $\mathbb{N}$ . Let us observe that  $a \cdot b$  is non square.

Let  $a$  be a non zero, square integer and  $n, m$  be natural numbers. Let us observe that  $a^n + a^m$  is non square.

Let  $a$  be a non zero, square element of  $\mathbb{N}$ . Let us note that  $a^n + a^m$  is non square.

Let  $a$  be a non zero, square integer and  $p$  be a prime natural number. Note that  $p \cdot a$  is non square.

Let  $a$  be a non trivial element of  $\mathbb{N}$ . One can verify that  $a - 1$  is non zero.

Let  $q$  be a square integer. Let us observe that  $|q|$  is square.

Let  $x$  be a non zero integer. Let us observe that  $|x|$  is non zero.

Let  $a$  be a non trivial, square element of  $\mathbb{N}$ . Let us observe that  $a - 1$  is non square.

Let  $a$  be a non trivial element of  $\mathbb{N}$ . Let us note that  $a \cdot (a - 1)$  is non square.

Let  $a, b$  be integers and  $n, m$  be natural numbers. One can verify that  $(a^n + b^n) \cdot (a^m - b^m) + (a^m + b^m) \cdot (a^n - b^n)$  is even and  $(a^n + b^n) \cdot (a^m + b^m) + (a^m - b^m) \cdot (a^n - b^n)$  is even.

Let  $a$  be an even integer. Let us note that  $\frac{a}{2}$  is integer.

Let  $a, b$  be non zero natural numbers. Note that  $a + b$  is non trivial.

Let  $b$  be a non zero natural number and  $a, c$  be non trivial natural numbers. Let us observe that  $c$ -count( $c^{a$ -count( $b$ )) reduces to  $a$ -count( $b$ ).

Let  $a, b$  be non zero integers. Let us note that  $\frac{a}{\gcd(a,b)}$  is integer and  $\frac{\text{lcm}(a,b)}{b}$  is integer and  $\frac{\text{lcm}(a,b)}{\gcd(a,b)}$  is integer.

Let  $a$  be an even integer. One can verify that  $\gcd(a, 2)$  reduces to 2.

Let us observe that there exists an even natural number which is non zero.

Let  $a$  be an even integer and  $n$  be a non zero natural number. Let us observe that  $a \cdot n$  is even and  $a^n$  is even.

Let  $a$  be an integer and  $n$  be a zero natural number. One can check that  $a \cdot n$  is even and  $a^n$  is odd.

Let  $a$  be an element of  $\mathbb{N}$ . Note that  $|a|$  reduces to  $a$ .

One can check that every integer which is non negative is also natural.

Let  $a$  be a non negative real number and  $n$  be a non zero natural number. Let us note that  $\sqrt[n]{a^n}$  reduces to  $a$  and  $(\sqrt[n]{a})^n$  reduces to  $a$ .

Now we state the propositions:

(11) If  $a \nmid b$ , then  $a \cdot c \nmid b$ .

(12) Let us consider non negative real numbers  $a$ ,  $b$ , and a positive natural number  $n$ . Then  $a^n = b^n$  if and only if  $a = b$ .

Let  $a$  be a real number and  $n$  be an even natural number. One can verify that  $a^n$  is non negative.

Let  $a$  be a negative real number and  $n$  be an odd natural number. One can verify that  $a^n$  is negative.

Now we state the propositions:

(13) Let us consider real numbers  $a$ ,  $b$ , and an odd natural number  $n$ . Then  $a^n = b^n$  if and only if  $a = b$ . The theorem is a consequence of (12).

(14) If  $a$  and  $b$  are relatively prime, then for every non zero natural number  $n$ ,  $a \cdot b = c^n$  iff  $\sqrt[n]{a}$ ,  $\sqrt[n]{b} \in \mathbb{N}$  and  $c = \sqrt[n]{a} \cdot \sqrt[n]{b}$ .

PROOF: If  $a \cdot b = c^n$ , then  $\sqrt[n]{a}$ ,  $\sqrt[n]{b} \in \mathbb{N}$  and  $c = \sqrt[n]{a} \cdot \sqrt[n]{b}$  by [14, (30)], [11, (11)], [1, (14)].  $\square$

(15) Let us consider a non zero natural number  $n$ , an integer  $a$ , and an integer  $b$ . Then  $b^n \mid a^n$  if and only if  $b \mid a$ .

PROOF: If  $b^n \mid a^n$ , then  $b \mid a$  by [10, (1)], [14, (3)], (4), [5, (3)].  $\square$

(16) Let us consider an integer  $a$ , and natural numbers  $m$ ,  $n$ . If  $m \geq n$ , then  $a^n \mid a^m$ .

(17) Let us consider integers  $a$ ,  $b$ . If  $a \mid b$  and  $b^m \mid c$ , then  $a^m \mid c$ . The theorem is a consequence of (4).

(18) Let us consider integers  $a$ ,  $p$ . If  $p^{2 \cdot n + k} \mid a^2$ , then  $p^n \mid a$ . The theorem is a consequence of (16), (4), and (12).

(19) Let us consider odd, square elements  $a$ ,  $b$  of  $\mathbb{N}$ . Then  $8 \mid a - b$ .

Let us consider odd natural numbers  $a$ ,  $b$ . Now we state the propositions:

(20) If  $4 \mid a - b$ , then  $4 \nmid a^n + b^n$ .

(21) If  $4 \mid a^n + b^n$ , then  $4 \nmid a^{2 \cdot n} + b^{2 \cdot n}$ .

(22) If  $4 \mid a^n - b^n$ , then  $4 \nmid a^{2 \cdot n} + b^{2 \cdot n}$ .

- (23) Let us consider odd natural numbers  $a, b$ . If  $2^m \mid a^n - b^n$ , then  $2^{m+1} \mid a^{2^n} - b^{2^n}$ .
- (24)  $a_1^3 - b_1^3 = (a_1 - b_1) \cdot (a_1^2 + b_1^2 + a_1 \cdot b_1)$ . The theorem is a consequence of (2).
- (25) Let us consider an odd natural number  $n$ . Then  $3 \mid a^n + b^n$  if and only if  $3 \mid a + b$ .  
 PROOF: Consider  $k$  such that  $n = 2 \cdot k + 1$ . If  $3 \mid a^n + b^n$ , then  $3 \mid a + b$  by [14, (173)], [5, (4)], [8, (1), (10)].  $\square$
- (26) Let us consider an integer  $c$ . If  $c \mid a - b$ , then  $c \mid a^n - b^n$ .
- (27) Let us consider an odd natural number  $n$ . Then  $3 \mid a^n - b^n$  if and only if  $3 \mid a - b$ .  
 PROOF: Consider  $k$  such that  $n = 2 \cdot k + 1$ . If  $3 \mid a^n - b^n$ , then  $3 \mid a - b$  by [14, (173)], [8, (10)], [5, (4)], [8, (1)].  $\square$
- (28) Let us consider a natural number  $n$ . Then  $a^n \equiv (a - b)^n \pmod{b}$ .
- (29) Let us consider a non trivial natural number  $a$ . Then there exists a prime natural number  $n$  such that  $n \mid a$ .
- (30) Let us consider a prime natural number  $p$ . If  $p \mid (p + (k + 1)) \cdot (p - (k + 1))$ , then  $k + 1 \geq p$ .
- (31) Let us consider a prime natural number  $p$ , and a non zero natural number  $k$ . If  $k < p$ , then  $p \nmid p^2 - k^2$ . The theorem is a consequence of (30).
- (32) Let us consider integers  $a, b$ , and an odd, prime natural number  $p$ . If  $p \nmid b$ , then if  $p \mid a - b$ , then  $p \nmid a + b$ .
- (33) Let us consider a non zero, square element  $a$  of  $\mathbb{N}$ , and a prime natural number  $p$ . If  $p \mid a$ , then  $a + p$  is not square.
- (34) Let us consider a non zero, square element  $a$  of  $\mathbb{N}$ , and a prime natural number  $p$ . If  $a + p$  is square, then  $p = 2 \cdot \sqrt{a} + 1$ .
- (35) Let us consider integers  $a, b, c$ . Suppose  $a$  and  $b$  are relatively prime. Then  $\gcd(c, a \cdot b) = (\gcd(c, a)) \cdot (\gcd(c, b))$ .
- (36) Let us consider a prime natural number  $p$ . If  $a \mid p^n$ , then there exists  $k$  such that  $a = p^k$ .

Let us consider non zero natural numbers  $a, b$  and a prime natural number  $p$ . Now we state the propositions:

- (37) If  $a + b = p$ , then  $a$  and  $b$  are relatively prime.
- (38) If  $a^n + b^n = p^n$ , then  $a$  and  $b$  are relatively prime.
- (39) Let us consider non zero natural numbers  $a, b$ . If  $c \geq a + b$ , then  $c^{k+1} \cdot (a + b) > a^{k+2} + b^{k+2}$ .

- (40) Let us consider natural numbers  $a, c$ , and a non zero natural number  $b$ .  
If  $a \cdot b < c < a \cdot (b + 1)$ , then  $a \nmid c$  and  $c \nmid a$ .
- (41) Let us consider real numbers  $a, b$ . Then  $a + b = \min(a, b) + \max(a, b)$ .
- (42) Let us consider non negative real numbers  $a, b$ . Then
- (i)  $\max(a^n, b^n) = (\max(a, b))^n$ , and
  - (ii)  $\min(a^n, b^n) = (\min(a, b))^n$ .
- (43) Let us consider a prime natural number  $p$ . Suppose  $a \cdot b = p^n$ . Then there exist natural numbers  $k, l$  such that
- (i)  $a = p^k$ , and
  - (ii)  $b = p^l$ , and
  - (iii)  $k + l = n$ .
- (44) Let us consider non trivial natural numbers  $a, b$ . If  $a$  and  $b$  are relatively prime, then  $a \nmid b$  and  $b \nmid a$ .
- (45) Let us consider a non trivial natural number  $a$ , and a prime natural number  $p$ . If  $p > a$ , then  $p \nmid a$  and  $a \nmid p$ . The theorem is a consequence of (44).
- (46) Let us consider a prime natural number  $p$ . Then
- (i)  $\gcd(a, p) = 1$ , or
  - (ii)  $\gcd(a, p) = p$ .
- (47) Let us consider a non trivial natural number  $a$ , and a prime natural number  $p$ . If  $a \mid p^n$ , then  $p \mid a$ . The theorem is a consequence of (46).
- (48) Let us consider odd natural numbers  $a, b$ , and an even natural number  $m$ . Then  $2\text{-count}(a^m + b^m) = 1$ .
- (49) Let us consider a non zero natural number  $a$ . Then there exists an odd natural number  $k$  such that  $a = 2^{2\text{-count}(a)} \cdot k$ .
- (50) Let us consider a non zero natural number  $b$ . Suppose  $a > b$ . Then there exists a prime natural number  $p$  such that  $p\text{-count}(a) > p\text{-count}(b)$ .  
PROOF: If for every prime natural number  $p$ ,  $p\text{-count}(a) \leq p\text{-count}(b)$ , then  $a \leq b$  by [12, (20)], [1, (14)].  $\square$
- (51) Let us consider natural numbers  $a, b, c$ . Suppose  $a \neq 1$  and  $b \neq 0$  and  $c \neq 0$  and  $b > a\text{-count}(c)$ . Then  $a^b \nmid c$ . The theorem is a consequence of (11).

Let us consider a non zero integer  $b$  and an integer  $a$ . Now we state the propositions:

- (52) If  $|a| \neq 1$ , then  $a^{|a|\text{-count}(|b|)} \mid b$  and  $a^{(|a|\text{-count}(|b|))+1} \nmid b$ .
- (53) If  $|a| \neq 1$ , then if  $a^n \mid b$  and  $a^{n+1} \nmid b$ , then  $n = |a|\text{-count}(|b|)$ .

(54) Let us consider a non zero natural number  $b$ , and a non trivial natural number  $a$ . Then  $a \mid b$  if and only if  $a\text{-count}(\gcd(a, b)) = 1$ .

PROOF: If  $a \mid b$ , then  $a\text{-count}(\gcd(a, b)) = 1$  by [14, (3)], [6, (22)].  $\square$

(55) Let us consider non zero natural numbers  $b, n$ , and a non trivial natural number  $a$ . Then  $a\text{-count}(\gcd(a, b)) = 1$  if and only if  $a^n\text{-count}((\gcd(a, b))^n) = 1$ . The theorem is a consequence of (15), (54), and (4).

(56) Let us consider a non zero natural number  $b$ , and a non trivial natural number  $a$ . Then  $a\text{-count}(\gcd(a, b)) = 0$  if and only if  $a\text{-count}(\gcd(a, b)) \neq 1$ . The theorem is a consequence of (54).

Let  $a, b$  be integers. The functor  $a\text{-count}(b)$  yielding a natural number is defined by the term

(Def. 6)  $|a|\text{-count}(|b|)$ .

Let  $a$  be an integer. Assume  $|a| \neq 1$ . Let  $b$  be a non zero integer. One can check that the functor  $a\text{-count}(b)$  is defined by

(Def. 7)  $a^{it} \mid b$  and  $a^{it+1} \nmid b$ .

Now we state the propositions:

(57) Let us consider a prime natural number  $p$ , and non zero integers  $a, b$ . Then  $p\text{-count}(a \cdot b) = (p\text{-count}(a)) + (p\text{-count}(b))$ .

(58) Let us consider a non trivial natural number  $a$ , and a non zero natural number  $b$ . Then  $a^{a\text{-count}(b)} \leq b$ .

(59) Let us consider a non trivial natural number  $a$ , and a non zero integer  $b$ . Then  $a^n \mid b$  if and only if  $n \leq a\text{-count}(b)$ .

PROOF: If  $a^n \mid b$ , then  $n \leq a\text{-count}(b)$  by [8, (9)], [7, (89)], [1, (13)]. If  $a^n \nmid b$ , then  $a\text{-count}(b) < n$  by [8, (9)], [7, (89)].  $\square$

(60) Let us consider a non trivial natural number  $a$ , a non zero integer  $b$ , and a non zero natural number  $n$ . Then  $n \cdot (a\text{-count}(b)) \leq a\text{-count}(b^n) < n \cdot ((a\text{-count}(b)) + 1)$ . The theorem is a consequence of (4) and (59).

(61) Let us consider a non trivial natural number  $a$ , and non zero natural numbers  $b, n$ . If  $b < a$ , then  $a\text{-count}(b^n) < n$ . The theorem is a consequence of (60).

(62) Let us consider a non trivial natural number  $a$ , and a non zero natural number  $b$ . If  $b < a^n$ , then  $a\text{-count}(b) < n$ . The theorem is a consequence of (59).

(63) Let us consider non zero natural numbers  $a, b$ , and a non trivial natural number  $n$ . Then  $a + b\text{-count}(a^n + b^n) < n$ . The theorem is a consequence of (62).

(64) Let us consider non zero natural numbers  $a, b$ . Then  $\gcd(a, b) = 1$  if and only if for every non trivial natural number  $c$ ,  $(c\text{-count}(a)) \cdot (c\text{-count}(b)) = 0$ .

PROOF: If  $\gcd(a, b) = 1$ , then for every non trivial natural number  $c$ ,  $(c\text{-count}(a)) \cdot (c\text{-count}(b)) = 0$  by [6, (27)]. If for every prime natural number  $c$ ,  $(c\text{-count}(a)) \cdot (c\text{-count}(b)) = 0$ , then  $\gcd(a, b) = 1$  by [6, (27)].  
□

Let us consider a non zero, even natural number  $m$  and odd natural numbers  $a, b$ . Now we state the propositions:

(65) If  $a \neq b$ , then  $2\text{-count}(a^{2 \cdot m} - b^{2 \cdot m}) \geq (2\text{-count}(a^m - b^m)) + 1$ . The theorem is a consequence of (12), (23), and (59).

(66) If  $a \neq b$ , then  $2\text{-count}(a^{2 \cdot m} - b^{2 \cdot m}) = (2\text{-count}(a^m - b^m)) + 1$ . The theorem is a consequence of (12), (57), and (48).

Let us consider a prime natural number  $p$  and integers  $a, b$ . Now we state the propositions:

(67) If  $|a| \neq |b|$ , then  $p\text{-count}(a^2 - b^2) = (p\text{-count}(a - b)) + (p\text{-count}(a + b))$ .

(68) If  $|a| \neq |b|$ , then  $p\text{-count}(a^3 - b^3) = (p\text{-count}(a - b)) + (p\text{-count}(a^2 + a \cdot b + b^2))$ . The theorem is a consequence of (24).

(69) Let us consider non zero natural numbers  $a, b$ . Then  $\frac{a}{\gcd(a, b)} = \frac{\text{lcm}(a, b)}{b}$ .

Let us consider a non zero natural number  $b$ . Now we state the propositions:

(70)  $\text{lcm}(a, a \cdot n + b) = ((\frac{a \cdot n}{b}) + 1) \cdot \text{lcm}(a, b)$ . The theorem is a consequence of (69).

(71)  $\text{lcm}(a, (n \cdot a + 1) \cdot b) = (n \cdot a + 1) \cdot \text{lcm}(a, b)$ . The theorem is a consequence of (70).

(72) Let us consider a non trivial natural number  $a$ , and non zero natural numbers  $n, b$ . Then  $a\text{-count}(b) \geq n \cdot (a^n\text{-count}(b))$ . The theorem is a consequence of (51).

Let us consider odd integers  $a, b$ . Now we state the propositions:

(73)  $4 \mid a - b$  if and only if  $4 \nmid a + b$ .

(74)  $2\text{-count}(a^2 + b^2) = 1$ . The theorem is a consequence of (5) and (73).

(75) Let us consider a prime natural number  $p$ , and natural numbers  $a, b$ . Suppose  $a \neq b$ . Then  $p\text{-count}(a + b) \geq p\text{-count}(\gcd(a, b))$ .

(76) Let us consider a non zero integer  $a$ , a non trivial natural number  $b$ , and an integer  $c$ . If  $a = b^{b\text{-count}(a)} \cdot c$ , then  $b \nmid c$ .

Let  $a$  be a non zero integer and  $b$  be a non trivial natural number. Let us note that  $\frac{a}{b^{b\text{-count}(a)}}$  is integer and  $\frac{a}{2^{2\text{-count}(a)}}$  is integer and  $\frac{a}{2^{2\text{-count}(a)}}$  is odd.

Now we state the proposition:

(77) Let us consider a non zero integer  $a$ , and a non trivial natural number  $b$ . Then  $b\text{-count}(a) = 0$  if and only if  $b \nmid a$ .

Let  $a$  be an odd integer. Observe that  $2\text{-count}(a)$  is zero.

Observe that  $\frac{a}{2^{2\text{-count}(a)}}$  reduces to  $a$ .

Now we state the propositions:

- (78) Let us consider a prime natural number  $a$ , a non zero integer  $b$ , and a natural number  $c$ . Then  $a\text{-count}(b^c) = c \cdot (a\text{-count}(b))$ .
- (79) Let us consider non zero natural numbers  $a, b$ , and an odd natural number  $n$ . Then  $\frac{a^{n+2}+b^{n+2}}{a+b} = a^{n+1} + b^{n+1} - a \cdot b \cdot (\frac{a^n+b^n}{a+b})$ . The theorem is a consequence of (3).
- (80) Let us consider odd integers  $a, b$ , and a natural number  $n$ . Then  $2\text{-count}(a^{2 \cdot n+1} - b^{2 \cdot n+1}) = 2\text{-count}(a - b)$ . The theorem is a consequence of (13), (2), and (57).
- (81) Let us consider odd integers  $a, b$ , and an odd natural number  $m$ . Then  $2\text{-count}(a^m + b^m) = 2\text{-count}(a + b)$ . The theorem is a consequence of (80).
- (82) Let us consider odd natural numbers  $a, b$ . Suppose  $a \neq b$ . Then  $1 = \min(2\text{-count}(a - b), 2\text{-count}(a + b))$ .

Let us consider a non trivial natural number  $a$  and non zero integers  $b, c$ . Now we state the propositions:

- (83) If  $a\text{-count}(b) > a\text{-count}(c)$ , then  $a^{a\text{-count}(c)} \mid b$  and  $a^{a\text{-count}(b)} \nmid c$ .
- (84) If  $a^{a\text{-count}(b)} \mid c$  and  $a^{a\text{-count}(c)} \mid b$ , then  $a\text{-count}(b) = a\text{-count}(c)$ . The theorem is a consequence of (83).
- (85) Let us consider integers  $a, b$ , and natural numbers  $m, n$ . If  $a^n \mid b$  and  $a^m \nmid b$ , then  $m > n$ . The theorem is a consequence of (16).

Let us consider a non trivial natural number  $a$  and non zero integers  $b, c$ . Now we state the propositions:

- (86) If  $a\text{-count}(b) = a\text{-count}(c)$  and  $a^n \mid b$ , then  $a^n \mid c$ . The theorem is a consequence of (85).
- (87)  $a\text{-count}(b) = a\text{-count}(c)$  if and only if for every natural number  $n$ ,  $a^n \mid b$  iff  $a^n \mid c$ .

PROOF: If  $a\text{-count}(b) \neq a\text{-count}(c)$ , then there exists a natural number  $n$  such that  $a^n \mid b$  and  $a^n \nmid c$  or  $a^n \mid c$  and  $a^n \nmid b$  by (83), [1, (13)], [7, (89)], [8, (9)].  $\square$

- (88) Let us consider odd integers  $a, b$ . Suppose  $|a| \neq |b|$ . Then
- (i)  $2\text{-count}((a - b)^2) \neq 2\text{-count}((a + b)^2)$ , and
  - (ii)  $2\text{-count}((a - b)^2) \neq (2\text{-count}(a^2)) - b^2$ .

The theorem is a consequence of (78), (73), and (87).

- (89) Let us consider a non trivial natural number  $b$ , and a non zero integer  $a$ . Then  $b\text{-count}(a) \neq 0$  if and only if  $b \mid a$ .

PROOF:  $b\text{-count}(|a|) \neq 0$  iff  $b \mid |a|$  by [6, (27)].  $\square$



- (90) Let us consider a non trivial natural number  $b$ , and a non zero natural number  $a$ . Then  $b$ -count( $a$ ) = 0 if and only if  $a \bmod b \neq 0$ . The theorem is a consequence of (89).
- (91) Let us consider a prime natural number  $p$ , and a non trivial natural number  $a$ . Then  $a$ -count( $p$ )  $\leq 1$ .
- (92) Let us consider non trivial natural numbers  $a$ ,  $b$ , and a non zero natural number  $c$ . Then  $a^{(a\text{-count}(b)) \cdot (b\text{-count}(c))} \leq c$ . The theorem is a consequence of (58).
- (93) Let us consider a prime natural number  $p$ , a non trivial natural number  $a$ , and a non zero natural number  $b$ . Then  $a$ -count( $p^b$ )  $\leq b$ . The theorem is a consequence of (89) and (59).
- (94) Let us consider a prime natural number  $p$ , and a non trivial natural number  $a$ . Then  $(p$ -count( $a$ ))  $\cdot$  ( $a$ -count( $p^n$ ))  $\leq n$ . The theorem is a consequence of (92).
- (95) Let us consider non trivial natural numbers  $a$ ,  $b$ , and a non zero natural number  $c$ . Then  $(a$ -count( $b$ ))  $\cdot$  ( $b$ -count( $c$ ))  $\leq a$ -count( $c$ ). The theorem is a consequence of (17).
- (96) Let us consider a non zero natural number  $a$ , and an odd natural number  $b$ . Then  $2$ -count( $a \cdot b$ ) =  $2$ -count( $a$ ).

Let us consider a non trivial natural number  $a$ . Now we state the propositions:

- (97)  $a^{n+1} + a^n < a^{n+2}$ .
- (98)  $(a + 1)^n + (a + 1)^n < (a + 1)^{n+1}$ .
- (99) Let us consider a non trivial, odd natural number  $a$ . Then  $a^n + a^n < a^{n+1}$ . The theorem is a consequence of (98).
- (100) Let us consider a non trivial natural number  $p$ . If  $a \nmid b$ , then  $(p^a)^c \neq p^b$ .
- (101) Let us consider non zero integers  $a$ ,  $b$ , and a non zero natural number  $n$ . Suppose there exists a prime natural number  $p$  such that  $n \nmid p$ -count( $a$ ). Then  $a \neq b^n$ .
- (102) Let us consider non zero integers  $a$ ,  $b$ , and a non zero natural number  $n$ . Suppose  $a = b^n$ . Let us consider a prime natural number  $p$ . Then  $n \mid p$ -count( $a$ ).
- (103) Let us consider positive real numbers  $a$ ,  $b$ , and a non trivial natural number  $n$ . Then  $(a + b)^n > a^n + b^n$ . The theorem is a consequence of (42) and (41).
- (104) Let us consider non zero integers  $a$ ,  $b$ , and an odd, prime natural number  $p$ . Suppose  $|a| \neq |b|$  and  $p \nmid b$ . Then  $p$ -count( $a^2 - b^2$ ) = max( $p$ -count( $a - b$ ),  $p$ -count( $a + b$ )). The theorem is a consequence of (32), (77), and (57).

- (105) Let us consider a non trivial natural number  $a$ , and a non zero integer  $b$ . Then  $a$ -count( $a^n \cdot b$ ) =  $n + (a$ -count( $b$ )).

ACKNOWLEDGEMENT: Ad Maiorem Dei Gloriam

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [3] Paul Erdős and János Surányi. *Topics in the Theory of Numbers*, chapter Divisibility, the Fundamental Theorem of Number Theory, pages 1–37. Springer New York, 2003. doi:10.1007/978-1-4613-0015-1\_1.
- [4] Jacek Gancarzewicz. *Arytmetyka*. Wydawnictwo UJ, Kraków, 2000. In Polish.
- [5] Andrzej Kondracki. The Chinese Remainder Theorem. *Formalized Mathematics*, 6(4):573–577, 1997.
- [6] Artur Korniłowicz and Piotr Rudnicki. Fundamental Theorem of Arithmetic. *Formalized Mathematics*, 12(2):179–186, 2004.
- [7] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [8] Rafał Kwiatek and Grzegorz Zwara. The divisibility of integers and integer relatively primes. *Formalized Mathematics*, 1(5):829–832, 1990.
- [9] Adam Naumowicz. An example of formalizing recent mathematical results in Mizar. *Journal of Applied Logic*, 4(4):396–413, 2006. doi:10.1016/j.jal.2005.10.003. Towards Computer Aided Mathematics.
- [10] Akira Nishino and Yasunari Shidama. The Maclaurin expansions. *Formalized Mathematics*, 13(3):421–425, 2005.
- [11] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(2):213–216, 1991.
- [12] Marco Riccardi. Pocklington’s theorem and Bertrand’s postulate. *Formalized Mathematics*, 14(2):47–52, 2006. doi:10.2478/v10037-006-0007-y.
- [13] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [14] Rafał Ziobro. Fermat’s Little Theorem via divisibility of Newton’s binomial. *Formalized Mathematics*, 23(3):215–229, 2015. doi:10.1515/forma-2015-0018.

Received June 30, 2016



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.

# Riemann-Stieltjes Integral<sup>1</sup>

Keiko Narita  
Hirosaki-city  
Aomori, Japan

Kazuhisa Nakasho  
Akita Prefectural University  
Akita, Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** In this article, the definitions and basic properties of Riemann-Stieltjes integral are formalized in Mizar [1]. In the first section, we showed the preliminary definition. We proved also some properties of finite sequences of real numbers. In Sec. 2, we defined variation. Using the definition, we also defined bounded variation and total variation, and proved theorems about related properties.

In Sec. 3, we defined Riemann-Stieltjes integral. Referring to the way of the article [5], we described the definitions. In the last section, we proved theorems about linearity of Riemann-Stieltjes integral. Because there are two types of linearity in Riemann-Stieltjes integral, we proved linearity in two ways. We showed the proof of theorems based on the description of the article [5]. These formalizations are based on [6], [4], [2], and [3].

MSC: 26A42 26A45 03B35

Keywords: Riemann-Stieltjes integral; bounded variation; linearity

MML identifier: INTEGR22, version: 8.1.05 5.37.1275

## 1. PROPERTIES OF REAL FINITE SEQUENCES

Let  $A$  be a subset of  $\mathbb{R}$  and  $\varrho$  be a real-valued function. The functor  $\text{vol}(A, \varrho)$  yielding a real number is defined by the term

---

<sup>1</sup>This work was supported by JSPS KAKENHI 22300285.

(Def. 1)  $\begin{cases} 0, & \text{if } A \text{ is empty,} \\ \varrho(\sup A) - \varrho(\inf A), & \text{otherwise.} \end{cases}$

Now we state the propositions:

- (1) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a partition  $D$  of  $A$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , a non empty, closed interval subset  $B$  of  $\mathbb{R}$ , and a finite sequence  $v$  of elements of  $\mathbb{R}$ . Suppose  $B \subseteq A$  and  $\text{len } D = \text{len } v$  and for every natural number  $i$  such that  $i \in \text{dom } v$  holds  $v(i) = \text{vol}(B \cap \text{divset}(D, i), \varrho)$ . Then  $\sum v = \text{vol}(B, \varrho)$ .
- (2) Let us consider natural numbers  $n, m$ , a function  $a$  from  $\text{Seg } n \times \text{Seg } m$  into  $\mathbb{R}$ , and finite sequences  $p, q$  of elements of  $\mathbb{R}$ . Suppose  $\text{dom } p = \text{Seg } n$  and for every natural number  $i$  such that  $i \in \text{dom } p$  there exists a finite sequence  $r$  of elements of  $\mathbb{R}$  such that  $\text{dom } r = \text{Seg } m$  and  $p(i) = \sum r$  and for every natural number  $j$  such that  $j \in \text{dom } r$  holds  $r(j) = a(i, j)$  and  $\text{dom } q = \text{Seg } m$  and for every natural number  $j$  such that  $j \in \text{dom } q$  there exists a finite sequence  $s$  of elements of  $\mathbb{R}$  such that  $\text{dom } s = \text{Seg } n$  and  $q(j) = \sum s$  and for every natural number  $i$  such that  $i \in \text{dom } s$  holds  $s(i) = a(i, j)$ . Then  $\sum p = \sum q$ .

## 2. THE DEFINITIONS OF BOUNDED VARIATION

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$ ,  $\varrho$  be a real-valued function, and  $t$  be a partition of  $A$ . A var-volume of  $\varrho$  and  $t$  is a finite sequence of elements of  $\mathbb{R}$  and is defined by

(Def. 2)  $\text{len } it = \text{len } t$  and for every natural number  $k$  such that  $k \in \text{dom } t$  holds  $it(k) = |\text{vol}(\text{divset}(t, k), \varrho)|$ .

Now we state the propositions:

- (3) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , a partition  $t$  of  $A$ , a var-volume  $F$  of  $\varrho$  and  $t$ , and a natural number  $k$ . If  $k \in \text{dom } F$ , then  $0 \leq F(k)$ .
- (4) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , a partition  $t$  of  $A$ , and a var-volume  $F$  of  $\varrho$  and  $t$ . Then  $0 \leq \sum F$ . The theorem is a consequence of (3).

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$  and  $\varrho$  be a function from  $A$  into  $\mathbb{R}$ . We say that  $\varrho$  is bounded-variation if and only if

(Def. 3) there exists a real number  $d$  such that  $0 < d$  and for every partition  $t$  of  $A$  and for every var-volume  $F$  of  $\varrho$  and  $t$ ,  $\sum F \leq d$ .

Assume  $\varrho$  is bounded-variation. The functor  $\text{TotalVD}(\varrho)$  yielding a real number is defined by

(Def. 4) there exists a non empty subset  $V$  of  $\mathbb{R}$  such that  $V$  is upper bounded and  $V = \{r, \text{ where } r \text{ is a real number} : \text{ there exists a partition } t \text{ of } A \text{ and there exists a var-volume } F \text{ of } \varrho \text{ and } t \text{ such that } r = \sum F\}$  and  $it = \sup V$ .

Now we state the propositions:

- (5) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and a partition  $T$  of  $A$ . Suppose  $\varrho$  is bounded-variation. Let us consider a var-volume  $F$  of  $\varrho$  and  $T$ . Then  $\sum F \leq \text{TotalVD}(\varrho)$ .
- (6) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , and a function  $\varrho$  from  $A$  into  $\mathbb{R}$ . If  $\varrho$  is bounded-variation, then  $0 \leq \text{TotalVD}(\varrho)$ . The theorem is a consequence of (4).

### 3. THE DEFINITIONS OF RIEMANN-STIELTJES INTEGRAL

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$ ,  $\varrho$  be a function from  $A$  into  $\mathbb{R}$ , and  $u$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . Assume  $\varrho$  is bounded-variation and  $\text{dom } u = A$ . Let  $t$  be a partition of  $A$ .

A middle volume of  $\varrho$ ,  $u$  and  $t$  is a finite sequence of elements of  $\mathbb{R}$  and is defined by

(Def. 5)  $\text{len } it = \text{len } t$  and for every natural number  $k$  such that  $k \in \text{dom } t$  there exists a real number  $r$  such that  $r \in \text{rng}(u \upharpoonright \text{divset}(t, k))$  and  $it(k) = r \cdot \text{vol}(\text{divset}(t, k), \varrho)$ .

Let  $T$  be a division sequence of  $A$ . A middle volume sequence of  $\varrho$ ,  $u$  and  $T$  is a sequence of  $\mathbb{R}^*$  and is defined by

(Def. 6) for every element  $k$  of  $\mathbb{N}$ ,  $it(k)$  is a middle volume of  $\varrho$ ,  $u$  and  $T(k)$ .

Let  $S$  be a middle volume sequence of  $\varrho$ ,  $u$  and  $T$  and  $k$  be a natural number. One can check that the functor  $S(k)$  yields a middle volume of  $\varrho$ ,  $u$  and  $T(k)$ . From now on  $A$  denotes a non empty, closed interval subset of  $\mathbb{R}$ ,  $\varrho$  denotes a function from  $A$  into  $\mathbb{R}$ ,  $u$  denotes a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ ,  $T$  denotes a division sequence of  $A$ ,  $S$  denotes a middle volume sequence of  $\varrho$ ,  $u$  and  $T$ , and  $k$  denotes a natural number.

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$ ,  $\varrho$  be a function from  $A$  into  $\mathbb{R}$ ,  $u$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ ,  $T$  be a division sequence of  $A$ , and  $S$  be a middle volume sequence of  $\varrho$ ,  $u$  and  $T$ . The functor  $\text{middle-sum}(S)$  yielding a sequence of real numbers is defined by

(Def. 7) for every natural number  $i$ ,  $it(i) = \sum(S(i))$ .

We say that  $u$  is Riemann-Stieltjes integrable with  $\varrho$  if and only if

(Def. 8) there exists a real number  $I$  such that for every division sequence  $T$  of  $A$  for every middle volume sequence  $S$  of  $\varrho$ ,  $u$  and  $T$  such that  $\delta_T$  is convergent and  $\lim \delta_T = 0$  holds middle-sum( $S$ ) is convergent and  $\lim$  middle-sum( $S$ ) =  $I$ .

Assume  $\varrho$  is bounded-variation and  $\text{dom } u = A$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho$ . The functor  $\int_{\varrho} u(x)dx$  yielding a real number is defined by

(Def. 9) for every division sequence  $T$  of  $A$  and for every middle volume sequence  $S$  of  $\varrho$ ,  $u$  and  $T$  such that  $\delta_T$  is convergent and  $\lim \delta_T = 0$  holds middle-sum( $S$ ) is convergent and  $\lim$  middle-sum( $S$ ) =  $it$ .

#### 4. LINEARITY OF RIEMANN-STIELTJES INTEGRAL

Now we state the propositions:

(7) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a real number  $r$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and partial functions  $u, w$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\text{dom } u = A$  and  $\text{dom } w = A$  and  $w = r \cdot u$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho$ . Then

(i)  $w$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} w(x)dx = r \cdot \int_{\varrho} u(x)dx.$$

(8) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and partial functions  $u, w$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\text{dom } u = A$  and  $\text{dom } w = A$  and  $w = -u$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho$ . Then

(i)  $w$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} w(x)dx = - \int_{\varrho} u(x)dx.$$

The theorem is a consequence of (7).

Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and partial functions  $u, v, w$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Now we state the propositions:

(9) Suppose  $\varrho$  is bounded-variation and  $\text{dom } u = A$  and  $\text{dom } v = A$  and  $\text{dom } w = A$  and  $w = u + v$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho$  and  $v$  is Riemann-Stieltjes integrable with  $\varrho$ . Then

(i)  $w$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} w(x)dx = \int_{\varrho} u(x)dx + \int_{\varrho} v(x)dx.$$

- (10) Suppose  $\varrho$  is bounded-variation and  $\text{dom } u = A$  and  $\text{dom } v = A$  and  $\text{dom } w = A$  and  $w = u - v$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho$  and  $v$  is Riemann-Stieltjes integrable with  $\varrho$ . Then

(i)  $w$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} w(x)dx = \int_{\varrho} u(x)dx - \int_{\varrho} v(x)dx.$$

The theorem is a consequence of (8) and (9).

- (11) Let us consider non empty, closed interval subsets  $A, B$  of  $\mathbb{R}$ , a real number  $r$ , and functions  $\varrho, \varrho_1$  from  $A$  into  $\mathbb{R}$ . Suppose  $B \subseteq A$  and  $\varrho = r \cdot \varrho_1$ . Then  $\text{vol}(B, \varrho) = r \cdot \text{vol}(B, \varrho_1)$ .

- (12) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a real number  $r$ , functions  $\varrho, \varrho_1$  from  $A$  into  $\mathbb{R}$ , and a partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\varrho_1$  is bounded-variation and  $\text{dom } u = A$  and  $\varrho = r \cdot \varrho_1$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_1$ . Then

(i)  $u$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} u(x)dx = r \cdot \int_{\varrho_1} u(x)dx.$$

The theorem is a consequence of (11).

- (13) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , functions  $\varrho, \varrho_1$  from  $A$  into  $\mathbb{R}$ , and a partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\varrho_1$  is bounded-variation and  $\text{dom } u = A$  and  $\varrho = -\varrho_1$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_1$ . Then

(i)  $u$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} u(x)dx = - \int_{\varrho_1} u(x)dx.$$

The theorem is a consequence of (12).

- (14) Let us consider non empty, closed interval subsets  $A, B$  of  $\mathbb{R}$ , and functions  $\varrho, \varrho_1, \varrho_2$  from  $A$  into  $\mathbb{R}$ . Suppose  $B \subseteq A$  and  $\varrho = \varrho_1 + \varrho_2$ . Then  $\text{vol}(B, \varrho) = \text{vol}(B, \varrho_1) + \text{vol}(B, \varrho_2)$ .

- (15) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , functions  $\varrho, \varrho_1, \varrho_2$  from  $A$  into  $\mathbb{R}$ , and a partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\varrho_1$  is bounded-variation and  $\varrho_2$  is bounded-variation and  $\text{dom } u = A$  and  $\varrho = \varrho_1 + \varrho_2$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_1$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_2$ . Then

(i)  $u$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} u(x)dx = \int_{\varrho_1} u(x)dx + \int_{\varrho_2} u(x)dx.$$

The theorem is a consequence of (14).

(16) Let us consider non empty, closed interval subsets  $A, B$  of  $\mathbb{R}$ , and functions  $\varrho, \varrho_1, \varrho_2$  from  $A$  into  $\mathbb{R}$ . Suppose  $B \subseteq A$  and  $\varrho = \varrho_1 - \varrho_2$ . Then  $\text{vol}(B, \varrho) = \text{vol}(B, \varrho_1) - \text{vol}(B, \varrho_2)$ . The theorem is a consequence of (14).

(17) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , functions  $\varrho, \varrho_1, \varrho_2$  from  $A$  into  $\mathbb{R}$ , and a partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose  $\varrho$  is bounded-variation and  $\varrho_1$  is bounded-variation and  $\varrho_2$  is bounded-variation and  $\text{dom } u = A$  and  $\varrho = \varrho_1 - \varrho_2$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_1$  and  $u$  is Riemann-Stieltjes integrable with  $\varrho_2$ . Then

(i)  $u$  is Riemann-Stieltjes integrable with  $\varrho$ , and

$$(ii) \int_{\varrho} u(x)dx = \int_{\varrho_1} u(x)dx - \int_{\varrho_2} u(x)dx.$$

The theorem is a consequence of (16).

#### REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] S.L. Gupta and Nisha Rani. *Fundamental Real Analysis*. Vikas Pub., 1986.
- [3] Einar Hille. *Methods in classical and functional analysis*. Addison-Wesley Publishing Co., Halsted Press, 1974.
- [4] H. Kestelman. *Modern theories of integration*. Dover Publications, 2nd edition, 1960.
- [5] Keiichi Miyajima, Takahiro Kato, and Yasunari Shidama. Riemann integral of functions from  $\mathbb{R}$  into real normed space. *Formalized Mathematics*, 19(1):17–22, 2011. doi:10.2478/v10037-011-0003-8.
- [6] Daniel W. Stroock. *A Concise Introduction to the Theory of Integration*. Springer Science & Business Media, 1999.

Received June 30, 2016



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.



# Quasi-uniform Space

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** In this article, using mostly Pervin [9], Kunzi [6], [8], [7], Williams [11] and Bourbaki [3] works, we formalize in Mizar [2] the notions of quasi-uniform space, semi-uniform space and locally uniform space.

We define the topology induced by a quasi-uniform space. Finally we formalize from the sets of the form  $((X \setminus \Omega) \times X) \cup (X \times \Omega)$ , the Csaszar-Pervin quasi-uniform space induced by a topological space.

MSC: 54E15 03B35

Keywords: quasi-uniform space; quasi-uniformity; Pervin space; Csaszar-Pervin quasi-uniformity

MML identifier: UNIFORM2, version: 8.1.05 5.37.1275

## 1. PRELIMINARIES

From now on  $X$  denotes a set,  $A$  denotes a subset of  $X$ , and  $R, S$  denote binary relations on  $X$ .

Now we state the propositions:

- (1)  $(X \setminus A) \times X \cup X \times A \subseteq X \times X$ .
- (2)  $(X \setminus A) \times X \cup X \times A = A \times A \cup (X \setminus A) \times X$ .

PROOF:  $(X \setminus A) \times X \cup X \times A \subseteq A \times A \cup (X \setminus A) \times X$  by (1), [4, (87)].  $\square$

- (3)  $R \cdot S = \{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } X : \text{ there exists an element } z \text{ of } X \text{ such that } \langle x, z \rangle \in R \text{ and } \langle z, y \rangle \in S\}$ .

PROOF:  $R \cdot S \subseteq \{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } X : \text{ there exists an element } z \text{ of } X \text{ such that } \langle x, z \rangle \in R \text{ and } \langle z, y \rangle \in S\}$  by [4, (87)].  $\{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } X : \text{ there exists an element } z \text{ of } X \text{ such that } \langle x, z \rangle \in R \text{ and } \langle z, y \rangle \in S\} \subseteq R \cdot S$ .  $\square$

Let  $X$  be a set and  $\mathcal{B}$  be a family of subsets of  $X$ . One can check that  $[\mathcal{B}]$  is non empty.

Let  $\mathcal{B}$  be a family of subsets of  $X \times X$ . Note that every element of  $\mathcal{B}$  is relation-like.

Let  $B$  be an element of  $\mathcal{B}$ . We introduce the notation  $B[\sim]$  as a synonym of  $B^\sim$ .

Let us observe that the functor  $B[\sim]$  yields a subset of  $X \times X$ . Let  $B_1, B_2$  be elements of  $\mathcal{B}$ . We introduce the notation  $B_1 \otimes B_2$  as a synonym of  $B_1 \cdot B_2$ .

One can verify that the functor  $B_1 \otimes B_2$  yields a subset of  $X \times X$ . Now we state the propositions:

- (4) Let us consider a set  $X$ , and a family  $G$  of subsets of  $X$ . If  $G$  is upper, then  $\text{FinMeetCl}(G)$  is upper.
- (5) If  $R$  is symmetric in  $X$ , then  $R^\sim$  is symmetric in  $X$ .

## 2. UNIFORM SPACE STRUCTURE

We consider uniform space structures which extend 1-sorted structures and are systems

$$\langle \text{a carrier, entourages} \rangle$$

where the carrier is a set, the entourages constitute a family of subsets of (the carrier)  $\times$  (the carrier).

Let  $U$  be a uniform space structure. We say that  $U$  is void if and only if

(Def. 1) the entourages of  $U$  is empty.

Let  $X$  be a set. The functor  $\text{UniformSpace}(X)$  yielding a strict uniform space structure is defined by the term

(Def. 2)  $\langle X, \emptyset_{2^{X \times X}} \rangle$ .

The functors: the trivial uniform space and the non empty trivial uniform space yielding strict uniform space structures are defined by terms

(Def. 3)  $\langle \emptyset, 2_*^{\emptyset \times \emptyset} \rangle$ ,

(Def. 4) there exists a family  $S_1$  of subsets of  $\{\emptyset\} \times \{\emptyset\}$  such that  $S_1 = \{\{\emptyset\} \times \{\emptyset\}\}$  and the non empty trivial uniform space =  $\langle \{\emptyset\}, S_1 \rangle$ ,

respectively. Let  $X$  be an empty set. One can verify that  $\text{UniformSpace}(X)$  is empty.

Let  $X$  be a non empty set. One can check that  $\text{UniformSpace}(X)$  is non empty.

Let  $X$  be a set. Note that  $\text{UniformSpace}(X)$  is void and the trivial uniform space is empty and non void and the non empty trivial uniform space is non empty and non void and there exists a uniform space structure which is empty,

strict, and void and there exists a uniform space structure which is empty, strict, and non void and there exists a uniform space structure which is non empty, strict, and void and there exists a uniform space structure which is non empty, strict, and non void.

Let  $X$  be a set and  $S_1$  be a family of subsets of  $X \times X$ . The functor  $S_1[\sim]$  yielding a family of subsets of  $X \times X$  is defined by the term

(Def. 5) the set of all  $S[\sim]$  where  $S$  is an element of  $S_1$ .

Let  $U$  be a uniform space structure. The functor  $U[\sim]$  yielding a uniform space structure is defined by the term

(Def. 6)  $\langle$ the carrier of  $U$ , (the entourages of  $U$ )  $[\sim]$  $\rangle$ .

Let  $U$  be a non empty uniform space structure. One can verify that  $U[\sim]$  is non empty.

### 3. AXIOMS

Let  $U$  be a uniform space structure. We say that  $U$  is upper if and only if

(Def. 7) the entourages of  $U$  is upper.

We say that  $U$  is  $\cap$ -closed if and only if

(Def. 8) the entourages of  $U$  is  $\cap$ -closed.

We say that  $U$  satisfies axiom U1 if and only if

(Def. 9) for every element  $S$  of the entourages of  $U$ ,  $\text{id}_\alpha \subseteq S$ , where  $\alpha$  is the carrier of  $U$ .

We say that  $U$  satisfies axiom U2 if and only if

(Def. 10) for every element  $S$  of the entourages of  $U$ ,  $S[\sim] \in$  the entourages of  $U$ .

We say that  $U$  satisfies axiom U3 if and only if

(Def. 11) for every element  $S$  of the entourages of  $U$ , there exists an element  $W$  of the entourages of  $U$  such that  $W \otimes W \subseteq S$ .

Let us consider a non void uniform space structure  $U$ . Now we state the propositions:

(6)  $U$  satisfies axiom U1 if and only if for every element  $S$  of the entourages of  $U$ , there exists a binary relation  $R$  on the carrier of  $U$  such that  $R = S$  and  $R$  is reflexive in the carrier of  $U$ .

(7)  $U$  satisfies axiom U1 if and only if for every element  $S$  of the entourages of  $U$ , there exists a total, reflexive binary relation  $R$  on the carrier of  $U$  such that  $R = S$ . The theorem is a consequence of (6).

Note that every uniform space structure which is void does not satisfy also axiom U2.

Now we state the proposition:

- (8) Let us consider a uniform space structure  $U$ . Suppose  $U$  satisfies axiom U2. Let us consider an element  $S$  of the entourages of  $U$ , and elements  $x, y$  of  $U$ . Suppose  $\langle x, y \rangle \in S$ . Then  $\langle y, x \rangle \in \bigcup(\text{the entourages of } U)$ .

Let us consider a non void uniform space structure  $U$ . Now we state the propositions:

- (9) Suppose for every element  $S$  of the entourages of  $U$ , there exists a binary relation  $R$  on the carrier of  $U$  such that  $S = R$  and  $R$  is symmetric in the carrier of  $U$ . Then  $U$  satisfies axiom U2. The theorem is a consequence of (5).
- (10) Suppose for every element  $S$  of the entourages of  $U$ , there exists a binary relation  $R$  on the carrier of  $U$  such that  $S = R$  and  $R$  is symmetric. Then  $U$  satisfies axiom U2. The theorem is a consequence of (9).
- (11) If for every element  $S$  of the entourages of  $U$ , there exists a tolerance  $R$  of the carrier of  $U$  such that  $S = R$ , then  $U$  satisfies axiom U1 and axiom U2. The theorem is a consequence of (7) and (10).

Let  $X$  be an empty set. Observe that  $\text{UniformSpace}(X)$  is upper and  $\cap$ -closed and satisfies axiom U1 and axiom U3 and does not satisfy axiom U2 and  $\text{UniformSpace}(\{\emptyset\})$  is upper and  $\cap$ -closed and does not satisfy axiom U2 and the trivial uniform space is upper and  $\cap$ -closed and satisfies axiom U1, axiom U2, and axiom U3 and the non empty trivial uniform space is upper and  $\cap$ -closed and satisfies axiom U1, axiom U2, and axiom U3.

There exists a uniform space structure which is strict, empty, non void, upper, and  $\cap$ -closed and satisfies axiom U1, axiom U2, and axiom U3 and every strict uniform space structure which is empty satisfies also axiom U1 and there exists a uniform space structure which is strict, non empty, non void, upper, and  $\cap$ -closed and satisfies axiom U1, axiom U2, and axiom U3.

Let  $S_4$  be a non empty uniform space structure satisfying axiom U1,  $x$  be an element of  $S_4$ , and  $V$  be an element of the entourages of  $S_4$ . The functor  $\text{Nbh}(V, x)$  yielding a non empty subset of  $S_4$  is defined by the term

(Def. 12)  $\{y, \text{ where } y \text{ is an element of } S_4 : \langle x, y \rangle \in V\}$ .

Now we state the proposition:

- (12) Let us consider a non empty uniform space structure  $U$  satisfying axiom U1, an element  $x$  of the carrier of  $U$ , and an element  $V$  of the entourages of  $U$ . Then  $x \in \text{Nbh}(V, x)$ .

Let  $U$  be a  $\cap$ -closed uniform space structure and  $V_1, V_2$  be elements of the entourages of  $U$ . One can check that the functor  $V_1 \cap V_2$  yields an element of the entourages of  $U$ . Now we state the proposition:

- (13) Let us consider a non empty,  $\cap$ -closed uniform space structure  $U$  satisfying axiom U1, an element  $x$  of  $U$ , and elements  $V, W$  of the entourages

of  $U$ . Then  $\text{Nbh}(V, x) \cap \text{Nbh}(W, x) = \text{Nbh}(V \cap W, x)$ .

Let  $U$  be a non empty uniform space structure satisfying axiom U1. Let us observe that the entourages of  $U$  has non empty elements and the entourages of  $U$  is non empty.

Let  $x$  be a point of  $U$ . The functor Neighborhood  $x$  yielding a family of subsets of  $U$  is defined by the term

(Def. 13) the set of all  $\text{Nbh}(V, x)$  where  $V$  is an element of the entourages of  $U$ .

Let us note that Neighborhood  $x$  is non empty.

Now we state the proposition:

(14) Let us consider a non empty uniform space structure  $S_4$  satisfying axiom U1, an element  $x$  of the carrier of  $S_4$ , and an element  $V$  of the entourages of  $S_4$ . Then

- (i)  $\text{Nbh}(V, x) = V^\circ\{x\}$ , and
- (ii)  $\text{Nbh}(V, x) = \text{rng}(V \upharpoonright \{x\})$ , and
- (iii)  $\text{Nbh}(V, x) = V^\circ x$ , and
- (iv)  $\text{Nbh}(V, x) = [x]_V$ , and
- (v)  $\text{Nbh}(V, x) = \text{neighbourhood}(x, V)$ .

PROOF:  $\text{Nbh}(V, x) = V^\circ\{x\}$  by [4, (87)].  $\square$

Let  $U$  be a non empty uniform space structure satisfying axiom U1. The functor Neighborhood  $U$  yielding a function from the carrier of  $U$  into  $2^{2^{\text{(the carrier of } U\text{)}}}$  is defined by

(Def. 14) for every element  $x$  of  $U$ ,  $it(x) = \text{Neighborhood } x$ .

We say that  $U$  is topological if and only if

(Def. 15)  $\langle \text{the carrier of } U, \text{Neighborhood } U \rangle$  is a topology from neighbourhoods.

#### 4. QUASI-UNIFORM SPACE

A quasi-uniform space is an upper,  $\cap$ -closed uniform space structure satisfying axiom U1 and axiom U3. In the sequel  $Q$  denotes a quasi-uniform space.

Now we state the propositions:

- (15) If the entourages of  $Q$  is empty, then the entourages of  $Q[\sim] = \{\emptyset\}$ .
- (16) Suppose the entourages of  $Q[\sim] = \{\emptyset\}$  and the entourages of  $Q[\sim]$  is upper. Then the carrier of  $Q$  is empty.

Let  $Q$  be a non void quasi-uniform space. One can check that  $Q[\sim]$  is upper and  $\cap$ -closed and satisfies axiom U1 and axiom U3.

Let  $X$  be a set and  $\mathcal{B}$  be a family of subsets of  $X \times X$ . We say that  $\mathcal{B}$  satisfies axiom UP1 if and only if

(Def. 16) for every element  $B$  of  $\mathcal{B}$ ,  $\text{id}_X \subseteq B$ .

We say that  $\mathcal{B}$  satisfies axiom UP3 if and only if

(Def. 17) for every element  $B_1$  of  $\mathcal{B}$ , there exists an element  $B_2$  of  $\mathcal{B}$  such that  $B_2 \otimes B_2 \subseteq B_1$ .

Now we state the propositions:

(17) Let us consider a non empty set  $X$ , and an empty family  $\mathcal{B}$  of subsets of  $X \times X$ . Then  $\mathcal{B}$  does not satisfy axiom UP1.

(18) Let us consider a set  $X$ , and a family  $\mathcal{B}$  of subsets of  $X \times X$ . Suppose  $\mathcal{B}$  is quasi-basis and satisfies axiom UP1 and axiom UP3. Then  $\langle X, [\mathcal{B}] \rangle$  is a quasi-uniform space.

## 5. SEMI-UNIFORM SPACE

A semi-uniform space is an upper,  $\cap$ -closed uniform space structure satisfying axiom U1 and axiom U2. From now on  $S_4$  denotes a semi-uniform space.

Let us observe that every semi-uniform space is non void.

Now we state the proposition:

(19) If  $S_4$  is empty, then  $\emptyset \in$  the entourages of  $S_4$ .

Let  $S_4$  be an empty semi-uniform space. One can verify that the entourages of  $S_4$  has the empty element.

## 6. LOCALLY UNIFORM SPACE

Let  $S_4$  be a non empty semi-uniform space. We say that  $S_4$  satisfies axiom UL if and only if

(Def. 18) for every element  $S$  of the entourages of  $S_4$  and for every element  $x$  of  $S_4$ , there exists an element  $W$  of the entourages of  $S_4$  such that  $\{y, \text{ where } y \text{ is an element of } S_4 : \langle x, y \rangle \in W \otimes W\} \subseteq \text{Nbh}(S, x)$ .

One can verify that every non empty semi-uniform space which satisfies axiom U3 satisfies also axiom UL and there exists a non empty semi-uniform space which satisfies axiom UL.

A locally uniform space is a non empty semi-uniform space satisfying axiom UL. Now we state the propositions:

(20) Let us consider a non empty, upper uniform space structure  $U$  satisfying axiom U1, and an element  $x$  of the carrier of  $U$ . Then Neighborhood  $x$  is upper.

- (21) Let us consider a non empty uniform space structure  $U$  satisfying axiom U1, an element  $x$  of  $U$ , and an element  $V$  of the entourages of  $U$ . Then  $x \in \text{Nbh}(V, x)$ .
- (22) Let us consider a non empty,  $\cap$ -closed uniform space structure  $U$  satisfying axiom U1, and an element  $x$  of  $U$ . Then Neighborhood  $x$  is  $\cap$ -closed. The theorem is a consequence of (13).
- (23) Let us consider a non empty, upper,  $\cap$ -closed uniform space structure  $U$  satisfying axiom U1, and an element  $x$  of  $U$ . Then Neighborhood  $x$  is a filter of the carrier of  $U$ . The theorem is a consequence of (22) and (20).

Let us observe that every locally uniform space is topological.

### 7. TOPOLOGICAL SPACE INDUCED BY A UNIFORM SPACE STRUCTURE

Let  $U$  be a topological, non empty uniform space structure satisfying axiom U1. The FMT induced by  $U$  yielding a non empty, strict topology from neighbourhoods is defined by the term

(Def. 19)  $\langle \text{the carrier of } U, \text{Neighborhood } U \rangle$ .

The topological space induced by  $U$  yielding a topological space is defined by the term

(Def. 20)  $\text{FMT2TopSpace}(\text{the FMT induced by } U)$ .

### 8. THE QUASI-UNIFORM PERVIN SPACE INDUCED BY A TOPOLOGICAL SPACE

Let  $X$  be a set and  $A$  be a subset of  $X$ . The functor  $\text{qBlock}(A)$  yielding a subset of  $X \times X$  is defined by the term

(Def. 21)  $(X \setminus A) \times X \cup X \times A$ .

Now we state the proposition:

- (24) (i)  $\text{id}_X \subseteq \text{qBlock}(A)$ , and
- (ii)  $\text{qBlock}(A) \cdot \text{qBlock}(A) \subseteq \text{qBlock}(A)$ .

PROOF:  $\text{id}_X \subseteq \text{qBlock}(A)$  by [4, (96)].  $\square$

Let  $T$  be a topological space. The functor  $\text{qBlocks}(T)$  yielding a family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 22) the set of all  $\text{qBlock}(O)$  where  $O$  is an element of the topology of  $T$ .

Let  $T$  be a non empty topological space. One can check that  $\text{qBlocks}(T)$  is non empty.

Let  $T$  be a topological space. The functor  $\text{FMCqBlocks}(T)$  yielding a family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 23)  $\text{FinMeetCl}(\text{qBlocks}(T))$ .

Let  $X$  be a set. One can check that every non empty family of subsets of  $X \times X$  which is  $\cap$ -closed is also quasi-basis.

In the sequel  $T$  denotes a topological space.

Let us consider  $T$ . One can check that  $\text{FMCqBlocks}(T)$  is  $\cap$ -closed and  $\text{FMCqBlocks}(T)$  is quasi-basis and  $\text{FMCqBlocks}(T)$  satisfies axiom UP1 and  $\text{FMCqBlocks}(T)$  satisfies axiom UP3.

Let  $T$  be a topological space. The Pervin quasi-uniformity of  $T$  yielding a strict quasi-uniform space is defined by the term

(Def. 24)  $\langle \text{the carrier of } T, [\text{FMCqBlocks}(T)] \rangle$ .

Now we state the propositions:

(25) Let us consider a non empty topological space  $T$ , and an element  $V$  of the entourages of the Pervin quasi-uniformity of  $T$ . Then there exists an element  $b$  of  $\text{FinMeetCl}(\text{qBlocks}(T))$  such that  $b \subseteq V$ .

(26) Let us consider a non empty topological space  $T$ , and a subset  $V$  of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$ . Suppose there exists an element  $b$  of  $\text{FinMeetCl}(\text{qBlocks}(T))$  such that  $b \subseteq V$ . Then  $V$  is an element of the entourages of the Pervin quasi-uniformity of  $T$ .

(27)  $\text{qBlocks}(T) \subseteq$  the entourages of the Pervin quasi-uniformity of  $T$ .

Let us consider a non void quasi-uniform space  $Q$ . Now we state the propositions:

(28)  $(\text{The carrier of } Q) \times (\text{the carrier of } Q) \in$  the entourages of  $Q$ .

(29) Suppose the carrier of  $T =$  the carrier of  $Q$  and  $\text{qBlocks}(T) \subseteq$  the entourages of  $Q$ . Then the entourages of the Pervin quasi-uniformity of  $T \subseteq$  the entourages of  $Q$ .

PROOF: The entourages of the Pervin quasi-uniformity of  $T \subseteq$  the entourages of  $Q$  by (28), [1, (1)].  $\square$

Let  $T$  be a non empty topological space. One can check that the Pervin quasi-uniformity of  $T$  is non empty and the Pervin quasi-uniformity of  $T$  is topological.

Now we state the propositions:

(30) Let us consider a non empty topological space  $T$ , an element  $x$  of  $\text{qBlocks}(T)$ , and an element  $y$  of the Pervin quasi-uniformity of  $T$ . Then  $\{z, \text{ where } z \text{ is an element of the Pervin quasi-uniformity of } T : \langle y, z \rangle \in x\} \in$  the topology of  $T$ .

(31) Let us consider a non empty topological space  $T$ , an element  $x$  of the carrier of the Pervin quasi-uniformity of  $T$ , and an element  $b$  of  $\text{FinMeetCl}(\text{qBlocks}(T))$ . Then  $\{y, \text{ where } y \text{ is an element of } T : \langle x, y \rangle \in b\} \in$  the to-



pology of  $T$ . The theorem is a consequence of (30).

- (32) Let us consider a non empty, strict quasi-uniform space  $U$ , a non empty, strict formal topological space  $F$ , and an element  $x$  of  $F$ . Suppose  $F = \langle \text{the carrier of } U, \text{Neighborhood } U \rangle$ . Then there exists an element  $y$  of  $U$  such that
- (i)  $x = y$ , and
  - (ii)  $U_F(x) = \text{Neighborhood } y$ .
- (33) Let us consider a non empty topological space  $T$ . Then the open set family of the FMT induced by the Pervin quasi-uniformity of  $T =$  the topology of  $T$ .
- PROOF: The open set family of the FMT induced by the Pervin quasi-uniformity of  $T \subseteq$  the topology of  $T$  by (32), [5, (18)], (31), [12, (25)]. The topology of  $T \subseteq$  the open set family of the FMT induced by the Pervin quasi-uniformity of  $T$  by (32), [10, (4)], [5, (18)], [4, (87)].  $\square$
- (34) Let us consider a non empty, strict topological space  $T$ . Then the topological space induced by the Pervin quasi-uniformity of  $T = T$ . The theorem is a consequence of (33).

ACKNOWLEDGEMENT: The author wants to express his gratitude to the anonymous referee for his/her work for the introduction of new notations and to make the presentation more readable.

#### REFERENCES

- [1] William W. Armstrong, Yatsuka Nakamura, and Piotr Rudnicki. Armstrong's axioms. *Formalized Mathematics*, 11(1):39–51, 2003.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [3] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [4] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [5] Roland Coghetto. Convergent filter bases. *Formalized Mathematics*, 23(3):189–203, 2015. doi:10.1515/forma-2015-0016.
- [6] Hans-Peter A. Künzi. Quasi-uniform spaces - eleven years later. In *Topology Proceedings*, volume 18, pages 143–171, 1993.
- [7] Hans-Peter A. Künzi. An introduction to quasi-uniform spaces. *Beyond Topology*, 486: 239–304, 2009.
- [8] Hans-Peter A. Künzi and Carolina Ryser. The Bourbaki quasi-uniformity. In *Topology Proceedings*, volume 20, pages 161–183, 1995.
- [9] William J. Pervin. Quasi-uniformization of topological spaces. *Mathematische Annalen*, 147(4):316–317, 1962.

- [10] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Formalized Mathematics*, 5(2):233–236, 1996.
- [11] James Williams. Locally uniform spaces. *Transactions of the American Mathematical Society*, 168:435–469, 1972.
- [12] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Formalized Mathematics*, 1(1):231–237, 1990.

*Received June 30, 2016*

---



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.

# Uniform Space

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** In this article, we formalize in Mizar [1] the notion of uniform space introduced by André Weil using the concepts of entourages [2].

We present some results between uniform space and pseudo metric space. We introduce the concepts of left-uniformity and right-uniformity of a topological group.

Next, we define the concept of the partition topology. Following the Vlach's works [11, 10], we define the semi-uniform space induced by a tolerance and the uniform space induced by an equivalence relation.

Finally, using mostly Gehrke, Grigorieff and Pin [4] works, a Pervin uniform space defined from the sets of the form  $((X \setminus A) \times (X \setminus A)) \cup (A \times A)$  is presented.

MSC: 54E15 03B35

Keywords: uniform space; uniformity; pseudo-metric space; topological group; partition topology; Pervin uniform space

MML identifier: UNIFORM3, version: 8.1.05 5.37.1275

## 1. PRELIMINARIES

From now on  $X$  denotes a set,  $D$  denotes a partition of  $X$ ,  $T$  denotes a non empty topological group, and  $A$  denotes a subset of  $X$ .

Now we state the propositions:

- (1)  $A \times A \cup (X \setminus A) \times (X \setminus A) \subseteq (X \setminus A) \times X \cup X \times A$ .
- (2)  $\{1, 2, 3\} \setminus \{1\} = \{2, 3\}$ .
- (3) Suppose  $X = \{1, 2, 3\}$  and  $A = \{1\}$ . Then
  - (i)  $\langle 2, 1 \rangle \in (X \setminus A) \times X \cup X \times A$ , and
  - (ii)  $\langle 2, 1 \rangle \notin A \times A \cup (X \setminus A) \times (X \setminus A)$ .

The theorem is a consequence of (2).

- (4) Let us consider a subset  $A$  of  $X$ . Then  $(A \times A \cup (X \setminus A) \times (X \setminus A))^\smile = A \times A \cup (X \setminus A) \times (X \setminus A)$ .
- (5) Let us consider subsets  $P_1, P_2$  of  $D$ . If  $\bigcup P_1 = \bigcup P_2$ , then  $P_1 = P_2$ .
- (6) Let us consider a subset  $P$  of  $D$ . Then  $\bigcup(D \setminus P) = \bigcup D \setminus \bigcup P$ .
- (7) Let us consider an upper family  $S_1$  of subsets of  $X$ , and an element  $S$  of  $S_1$ . Then  $\bigcap S_1 \subseteq S$ .
- (8) Let us consider an additive group  $G$ , and subsets  $A, B, C, D$  of  $G$ . If  $A \subseteq B$  and  $C \subseteq D$ , then  $A + C \subseteq B + D$ .

Let us consider an element  $e$  of  $T$  and a neighbourhood  $V$  of  $\mathbf{1}_T$ . Now we state the propositions:

- (9)  $\{e\} \cdot V$  is a neighbourhood of  $e$ .
- (10)  $V \cdot \{e\}$  is a neighbourhood of  $e$ .
- (11) Let us consider a neighbourhood  $V$  of  $\mathbf{1}_T$ . Then  $V^{-1}$  is a neighbourhood of  $\mathbf{1}_T$ .

## 2. UNIFORM SPACE

A uniform space is an upper,  $\cap$ -closed uniform space structure satisfying axiom U1, axiom U2, and axiom U3. From now on  $Q$  denotes a uniform space.

Now we state the propositions:

- (12)  $Q$  is a quasi-uniform space.
- (13)  $Q$  is a semi-uniform space.

Let  $X$  be a set and  $\mathcal{B}$  be a family of subsets of  $X \times X$ . We say that  $\mathcal{B}$  satisfies axiom UP2 if and only if

- (Def. 1) for every element  $B_1$  of  $\mathcal{B}$ , there exists an element  $B_2$  of  $\mathcal{B}$  such that  $B_2 \subseteq B_1^\smile$ .

Now we state the proposition:

- (14) Let us consider an empty set  $X$ . Then every empty family of subsets of  $X \times X$  is quasi-basis and satisfies axiom UP1, axiom UP2, and axiom UP3.

One can verify that there exists a uniform space which is strict.

Now we state the proposition:

- (15) Let us consider a set  $X$ , and a family  $S_1$  of subsets of  $X \times X$ . Suppose  $X = \{\emptyset\}$  and  $S_1 = \{X \times X\}$ . Then  $\langle X, S_1 \rangle$  is a uniform space.

Let us observe that there exists a strict uniform space which is non empty.

Now we state the proposition:

- (16) Let us consider a set  $X$ , and a family  $\mathcal{B}$  of subsets of  $X \times X$ . Suppose  $\mathcal{B}$  is quasi-basis and satisfies axiom UP1, axiom UP2, and axiom UP3. Then there exists a strict uniform space  $Q$  such that
- (i) the carrier of  $Q = X$ , and
  - (ii) the entourages  $Q = [\mathcal{B}]$ .

### 3. OPEN SET AND UNIFORM SPACE

Now we state the propositions:

- (17) Let us consider a non empty uniform space  $Q$ . Then
- (i) the carrier of the topological space induced by  $Q =$  the carrier of  $Q$ , and
  - (ii) the topology of the topological space induced by  $Q =$  the open set family of the FMTinduced by  $Q$ .
- (18) Let us consider a non empty uniform space  $Q$ , and a subset  $S$  of the FMTinduced by  $Q$ . Then  $S$  is open if and only if for every element  $x$  of  $Q$  such that  $x \in S$  holds  $S \in$  Neighborhood  $x$ .
- (19) Let us consider a non empty uniform space  $Q$ . Then the open set family of the FMTinduced by  $Q =$  the set of all  $O$  where  $O$  is an open subset of the FMTinduced by  $Q$ .

Let us consider a non empty uniform space  $Q$  and a subset  $S$  of the FMTinduced by  $Q$ . Now we state the propositions:

- (20)  $S$  is open if and only if  $S \in$  the open set family of the FMTinduced by  $Q$ .
- (21)  $S \in$  the open set family of the FMTinduced by  $Q$  if and only if for every element  $x$  of  $Q$  such that  $x \in S$  holds  $S \in$  Neighborhood  $x$ .

### 4. PSEUDO METRIC SPACE AND UNIFORM SPACE

Let  $M$  be a non empty metric structure and  $r$  be a positive real number. The functor  $\text{ent}(M, r)$  yielding a subset of (the carrier of  $M$ )  $\times$  (the carrier of  $M$ ) is defined by the term

(Def. 2)  $\{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } M : \rho(x, y) \leq r\}$ .

Let  $M$  be a non empty, reflexive metric structure. Let us observe that  $\text{ent}(M, r)$  is non empty.

Let  $M$  be a non empty metric structure. The functor  $\text{ENT}(M)$  yielding a non empty family of subsets of (the carrier of  $M$ )  $\times$  (the carrier of  $M$ ) is defined by the term

(Def. 3) the set of all  $\text{ent}(M, r)$  where  $r$  is a positive real number.

The uniformity induced by  $M$  yielding a uniform space structure is defined by the term

(Def. 4)  $\langle \text{the carrier of } M, [\text{ENT}(M)] \rangle$ .

Let  $M$  be a pseudo metric space. The uniformity induced by  $M$  yielding a non empty, strict uniform space is defined by the term

(Def. 5)  $\langle \text{the carrier of } M, [\text{ENT}(M)] \rangle$ .

Let us consider a pseudo metric space  $M$ . Now we state the propositions:

(22) The open set family of the FMT induced by the uniformity induced by  $M =$  the open set family of  $M$ .

PROOF: Set  $X =$  the open set family of the FMT induced by the uniformity induced by  $M$ . Set  $Y =$  the open set family of  $M$ .  $X \subseteq Y$  by (18), (20), [5, (11)]. Reconsider  $t_1 = t$  as a subset of  $M$ . For every element  $x$  of the uniformity induced by  $M$  such that  $x \in t_1$  holds  $t_1 \in \text{Neighborhood } x$  by [5, (11)].  $\square$

(23) The topological space induced by the uniformity induced by  $M = M_{\text{top}}$ . The theorem is a consequence of (22).

## 5. UNIFORM SPACE AND TOPOLOGICAL GROUP

Let  $G$  be a topological group and  $Q$  be a neighbourhood of  $\mathbf{1}_G$ . The functor  $\text{leftU}(Q)$  yielding a subset of  $(\text{the carrier of } G) \times (\text{the carrier of } G)$  is defined by the term

(Def. 6)  $\{ \langle x, y \rangle, \text{ where } x \text{ is an element of } G, y \text{ is an element of } G : x^{-1} \cdot y \in Q \}$ .

Let  $T$  be a non empty topological group. The functor  $\text{SleftU}(T)$  yielding a non empty family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 7) the set of all  $\text{leftU}(Q)$  where  $Q$  is a neighbourhood of  $\mathbf{1}_T$ .

The left-uniformity  $T$  yielding a non empty uniform space is defined by the term

(Def. 8)  $\langle \text{the carrier of } T, [\text{SleftU}(T)] \rangle$ .

Let  $G$  be a topological group and  $Q$  be a neighbourhood of  $\mathbf{1}_G$ . The functor  $\text{rightU}(Q)$  yielding a subset of  $(\text{the carrier of } G) \times (\text{the carrier of } G)$  is defined by the term

(Def. 9)  $\{ \langle x, y \rangle, \text{ where } x \text{ is an element of } G, y \text{ is an element of } G : y \cdot x^{-1} \in Q \}$ .

Let  $T$  be a non empty topological group. The functor  $\text{SrightU}(T)$  yielding a non empty family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 10) the set of all  $\text{rightU}(Q)$  where  $Q$  is a neighbourhood of  $\mathbf{1}_T$ .

The right-uniformity  $T$  yielding a non empty uniform space is defined by the term

(Def. 11)  $\langle \text{the carrier of } T, [\text{SrightU}(T)] \rangle$ .

Now we state the propositions:

(24) Let us consider a non empty, commutative topological group  $T$ , and a neighbourhood  $Q$  of  $\mathbf{1}_T$ . Then  $\text{leftU}(Q) = \text{rightU}(Q)$ .

(25) Let us consider a non empty, commutative topological group  $T$ . Then the left-uniformity  $T =$  the right-uniformity  $T$ . The theorem is a consequence of (24).

Let  $G$  be a semi additive topological group and  $Q$  be a neighbourhood of  $0_G$ . The functor  $\text{leftU}(Q)$  yielding a subset of  $(\text{the carrier of } G) \times (\text{the carrier of } G)$  is defined by the term

(Def. 12)  $\{ \langle x, y \rangle, \text{ where } x \text{ is an element of } G, y \text{ is an element of } G : -x + y \in Q \}$ .

Let  $T$  be a non empty semi additive topological group. The functor  $\text{SleftU}(T)$  yielding a non empty family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 13) the set of all  $\text{leftU}(Q)$  where  $Q$  is a neighbourhood of  $0_T$ .

Let  $T$  be a non empty topological additive group. The left-uniformity  $T$  yielding a non empty uniform space is defined by the term

(Def. 14)  $\langle \text{the carrier of } T, [\text{SleftU}(T)] \rangle$ .

Let  $G$  be a semi additive topological group and  $Q$  be a neighbourhood of  $0_G$ . The functor  $\text{rightU}(Q)$  yielding a subset of  $(\text{the carrier of } G) \times (\text{the carrier of } G)$  is defined by the term

(Def. 15)  $\{ \langle x, y \rangle, \text{ where } x \text{ is an element of } G, y \text{ is an element of } G : y + -x \in Q \}$ .

Let  $T$  be a non empty semi additive topological group. The functor  $\text{SrightU}(T)$  yielding a non empty family of subsets of  $(\text{the carrier of } T) \times (\text{the carrier of } T)$  is defined by the term

(Def. 16) the set of all  $\text{rightU}(Q)$  where  $Q$  is a neighbourhood of  $0_T$ .

Let  $T$  be a non empty topological additive group. The right-uniformity  $T$  yielding a non empty uniform space is defined by the term

(Def. 17)  $\langle \text{the carrier of } T, [\text{SrightU}(T)] \rangle$ .

Now we state the propositions:

(26) Let us consider an Abelian semi additive topological group  $T$ , and a neighbourhood  $Q$  of  $0_T$ . Then  $\text{leftU}(Q) = \text{rightU}(Q)$ .

(27) Let us consider a non empty topological additive group  $T$ . Suppose  $T$

is Abelian. Then the left-uniformity  $T =$  the right-uniformity  $T$ . The theorem is a consequence of (26).

- (28) The topology of the topological space induced by the left-uniformity  $T =$  the topology of  $T$ .

PROOF: Set  $X =$  the topology of  $\text{FMT2TopSpace}$ (the FMT induced by the left-uniformity  $T$ ). Set  $Y =$  the topology of  $T$ .  $X \subseteq Y$  by (9), [6, (7)].  $Y \subseteq X$  by [9, (3)], [6, (6)], [8, (6)].  $\square$

- (29) The topology of the topological space induced by the right-uniformity  $T =$  the topology of  $T$ .

PROOF: Set  $X =$  the topology of  $\text{FMT2TopSpace}$ (the FMT induced by the right-uniformity  $T$ ). Set  $Y =$  the topology of  $T$ .  $X \subseteq Y$  by (10), [6, (7)].  $Y \subseteq X$  by [9, (3)], [6, (6)], [8, (6)].  $\square$

## 6. FUNCTION UNIFORMLY CONTINUOUS

Let  $Q_1, Q_2$  be uniform space structures and  $f$  be a function from  $Q_1$  into  $Q_2$ . We say that  $f$  is uniformly continuous if and only if

- (Def. 18) for every element  $V$  of the entourages  $Q_2$ , there exists an element  $Q$  of the entourages  $Q_1$  such that for every objects  $x, y$  such that  $\langle x, y \rangle \in Q$  holds  $\langle f(x), f(y) \rangle \in V$ .

Let  $Q_1, Q_2$  be non empty uniform space structures satisfying axiom U1. One can check that there exists a function from  $Q_1$  into  $Q_2$  which is uniformly continuous.

## 7. PARTITION TOPOLOGY

Now we state the propositions:

- (30) the set of all  $\cup P$  where  $P$  is a subset of  $D = \text{UniCl}(D)$ .  
 (31)  $X \in \text{UniCl}(D)$ . The theorem is a consequence of (30).  
 (32) If  $D = \emptyset$ , then  $X$  is empty and  $\text{UniCl}(D) = \{\emptyset\}$ .

Let  $X$  be a set and  $D$  be a partition of  $X$ . Let us note that  $\text{UniCl}(D)$  is  $\cap$ -closed and  $\text{UniCl}(D)$  is union-closed and every family of subsets of  $X$  which is union-closed is also  $\cup$ -closed.

Let  $D$  be a partition of  $X$ . Let us note that  $\text{UniCl}(D)$  is closed for complement operator and  $\text{UniCl}(D)$  is  $\cup$ -closed and  $\setminus$ -closed.

Now we state the proposition:

- (33)  $\text{UniCl}(D)$  is a ring of sets. The theorem is a consequence of (30).



Let us consider  $X$  and  $D$ . One can verify that  $\text{UniCl}(D)$  has the empty element.

Let  $X$  be a set and  $D$  be a partition of  $X$ . Let us observe that  $\text{UniCl}(D)$  is non empty.

Now we state the proposition:

(34)  $\text{UniCl}(D)$  is a field of subsets of  $X$ .

Let  $X$  be a set and  $D$  be a partition of  $X$ . Observe that  $\text{UniCl}(D)$  is  $\sigma$ -additive and  $\text{UniCl}(D)$  is  $\sigma$ -multiplicative.

Now we state the proposition:

(35)  $\text{UniCl}(D)$  is a  $\sigma$ -field of subsets of  $X$ .

Let  $X$  be a set and  $D$  be a partition of  $X$ . Observe that  $\text{UniCl}(D)$  is closed for countable unions and closed for countable meets.

Now we state the proposition:

(36) Let us consider a non empty set  $\Omega$ , and a partition  $D$  of  $\Omega$ . Then  $\text{UniCl}(D)$  is a Dynkin system of  $\Omega$ .

Let  $X$  be a set and  $D$  be a partition of  $X$ . The partition topology  $D$  yielding a topological space is defined by the term

(Def. 19)  $\langle X, \text{UniCl}(D) \rangle$ .

Now we state the propositions:

(37) Every open subset of the partition topology  $D$  is closed.

(38) Every closed subset of the partition topology  $D$  is open.

(39) Let us consider a subset  $S$  of the partition topology  $D$ . Then  $S$  is open if and only if  $S$  is closed.

Let  $X$  be a non empty set and  $D$  be a partition of  $X$ . Observe that the partition topology  $D$  is non empty.

Let us consider a non empty set  $X$  and a partition  $D$  of  $X$ . Now we state the propositions:

(40)  $\text{LC}(\text{the partition topology } D) = \text{UniCl}(D)$ . The theorem is a consequence of (38) and (31).

(41)  $\text{OpenClosedSet}(\text{the partition topology } D) = \text{the topology of the partition topology } D$ . The theorem is a consequence of (37).

## 8. UNIFORM SPACE AND PARTITION TOPOLOGY

In the sequel  $R$  denotes a binary relation on  $X$ .

Let  $X$  be a set and  $R$  be a binary relation on  $X$ . The functor  $\rho(R)$  yielding a non empty family of subsets of  $X \times X$  is defined by the term

(Def. 20)  $\{S, \text{ where } S \text{ is a subset of } X \times X : R \subseteq S\}$ .

Now we state the propositions:

$$(42) \quad [\rho(R)] = \rho(R).$$

$$(43) \quad [\{R\}] = \rho(R).$$

$$(44) \quad \rho(R) \text{ is upper and } \cap\text{-closed.}$$

Let us consider  $X$  and  $R$ . Observe that  $\rho(R)$  is quasi-basis.

Now we state the propositions:

(45) Let us consider a total, reflexive binary relation  $R$  on  $X$ . Then  $\rho(R)$  satisfies axiom UP1.

(46) Let us consider a symmetric binary relation  $R$  on  $X$ . Then  $\rho(R)$  satisfies axiom UP2.

(47) Let us consider a total, transitive binary relation  $R$  on  $X$ . Then  $\rho(R)$  satisfies axiom UP3.

Let  $X$  be a set and  $R$  be a binary relation on  $X$ . The uniformity induced by  $R$  yielding an upper,  $\cap$ -closed, strict uniform space structure is defined by the term

(Def. 21)  $\langle X, \rho(R) \rangle$ .

Now we state the propositions:

(48) Let us consider a set  $X$ , and a total, reflexive binary relation  $R$  on  $X$ . Then the uniformity induced by  $R$  satisfies axiom U1. The theorem is a consequence of (45).

(49) Let us consider a set  $X$ , and a symmetric binary relation  $R$  on  $X$ . Then the uniformity induced by  $R$  satisfies axiom U2. The theorem is a consequence of (46).

(50) Let us consider a set  $X$ , and a total, transitive binary relation  $R$  on  $X$ . Then the uniformity induced by  $R$  satisfies axiom U3. The theorem is a consequence of (47).

Let  $X$  be a set and  $R$  be a tolerance of  $X$ . Note that the uniformity induced by  $R$  yields a strict semi-uniform space. Now we state the proposition:

(51) Let us consider a set  $X$ , and an equivalence relation  $R$  of  $X$ . Then the uniformity induced by  $R$  is a uniform space.

Let  $X$  be a set and  $R$  be an equivalence relation of  $X$ . Observe that the uniformity induced by  $R$  yields a strict uniform space. Let  $X$  be a non empty set and  $R$  be a tolerance of  $X$ . Let us note that the uniformity induced by  $R$  is non empty and every non empty uniform space is topological.

Let  $Q$  be a non empty uniform space. The functor  ${}^{\textcircled{Q}}$  yielding a topological, non empty uniform space structure satisfying axiom U1 is defined by the term

(Def. 22)  $Q$ .

Now we state the proposition:

- (52) Let us consider a non empty set  $X$ , and an equivalence relation  $R$  of  $X$ . Then the topological space induced by  $\textcircled{R}$ (the uniformity induced by  $R$ ) = the partition topology Classes  $R$ . The theorem is a consequence of (30) and (18).

9. UNIFORMITY INDUCED BY A TOLERANCE OR BY AN EQUIVALENCE

Now we state the proposition:

- (53) Let us consider an upper uniform space structure  $Q$ . Suppose  $\bigcap$ (the entourages  $Q$ )  $\in$  the entourages  $Q$ . Then there exists a binary relation  $R$  on the carrier of  $Q$  such that
- (i)  $\bigcap$ (the entourages  $Q$ ) =  $R$ , and
  - (ii) the entourages  $Q = \rho(R)$ .

PROOF: Reconsider  $R = \bigcap$ (the entourages  $Q$ ) as a binary relation on the carrier of  $Q$ .  $\rho(R) \subseteq$  the entourages  $Q$ . The entourages  $Q \subseteq \rho(R)$  by [7, (3)].  $\square$

Let  $Q$  be a uniform space structure. The functor  $\text{Uniformity2InternalRel}(Q)$  yielding a binary relation on the carrier of  $Q$  is defined by the term

(Def. 23)  $\bigcap$ (the entourages  $Q$ ).

The functor  $\text{UniformSpaceStr2RelStr}(Q)$  yielding a relational structure is defined by the term

(Def. 24)  $\langle$ the carrier of  $Q$ ,  $\text{Uniformity2InternalRel}(Q)\rangle$ .

Let  $R_1$  be a relational structure. The functor  $\text{InternalRel2Uniformity}(R_1)$  yielding a family of subsets of (the carrier of  $R_1$ )  $\times$  (the carrier of  $R_1$ ) is defined by the term

(Def. 25)  $\{R$ , where  $R$  is a binary relation on the carrier of  $R_1$  : the internal relation of  $R_1 \subseteq R\}$ .

The functor  $\text{RelStr2UniformSpaceStr}(R_1)$  yielding a strict uniform space structure is defined by the term

(Def. 26)  $\langle$ the carrier of  $R_1$ ,  $\text{InternalRel2Uniformity}(R_1)\rangle$ .

The functor  $\text{InternalRel2Element}(R_1)$  yielding an element of the entourages  $\text{RelStr2UniformSpaceStr}(R_1)$  is defined by the term

(Def. 27) the internal relation of  $R_1$ .

Now we state the propositions:

- (54) Let us consider a binary relation  $R$  on  $X$ . Then  $\bigcap \rho(R) = R$ .

- (55) Let us consider a strict relational structure  $R_1$ . Then  $\text{UniformSpaceStr2-RelStr}(\text{RelStr2UniformSpaceStr}(R_1)) = R_1$ . The theorem is a consequence of (54).
- (56) Let us consider a uniform space structure  $Q$ . Then
- (i) the carrier of  $\text{RelStr2UniformSpaceStr}(\text{UniformSpaceStr2RelStr}(Q))$  = the carrier of  $Q$ , and
  - (ii) the entourages  $\text{RelStr2UniformSpaceStr}(\text{UniformSpaceStr2RelStr}(Q)) = \rho(\bigcap(\text{the entourages } Q))$ .
- (57) Let us consider a family  $S_1$  of subsets of  $X \times X$ , and a binary relation  $R$  on  $X$ . If  $S_1 = \rho(R)$ , then  $S_1 \subseteq \rho(\bigcap S_1)$ .
- (58) Let us consider an upper family  $S_1$  of subsets of  $X \times X$ . If  $\bigcap S_1 \in S_1$ , then  $\rho(\bigcap S_1) \subseteq S_1$ .
- (59) Let us consider an upper family  $S_1$  of subsets of  $X \times X$ , and a binary relation  $R$  on  $X$ . Suppose  $R \in S_1$  and  $S_1 = \rho(R)$  and  $\bigcap S_1 \in S_1$ . Then  $\rho(\bigcap S_1) = S_1$ .
- (60) Let us consider an upper uniform space structure  $Q$ . Suppose there exists a binary relation  $R$  on the carrier of  $Q$  such that the entourages  $Q = \rho(R)$  and  $\bigcap(\text{the entourages } Q) \in \text{the entourages } Q$ . Then the entourages  $Q = \rho(\bigcap(\text{the entourages } Q))$ . The theorem is a consequence of (57) and (58).
- (61) Let us consider an upper uniform space structure  $Q$ , and a binary relation  $R$  on the carrier of  $Q$ . Suppose the entourages  $Q = \rho(R)$  and  $\bigcap(\text{the entourages } Q) \in \text{the entourages } Q$ . Then the entourages  $Q = \rho(\bigcap(\text{the entourages } Q))$ .

Let us consider a tolerance  $R$  of  $X$ . Now we state the propositions:

- (62) (i) the uniformity induced by  $R$  is a semi-uniform space, and
- (ii) the entourages the uniformity induced by  $R = \rho(R)$ , and
  - (iii)  $\bigcap(\text{the entourages the uniformity induced by } R) = R$ .
- (63)  $\text{RelStr2UniformSpaceStr}(\text{UniformSpaceStr2RelStr}(\text{the uniformity induced by } R)) = \text{the uniformity induced by } R$ . The theorem is a consequence of (54).
- (64) Let us consider an equivalence relation  $R$  of  $X$ . Then  $\text{RelStr2UniformSpaceStr}(\text{UniformSpaceStr2RelStr}(\text{the uniformity induced by } R)) = \text{the uniformity induced by } R$ . The theorem is a consequence of (54).

10. UNIFORM PERVIN SPACE

Let  $X$  be a set,  $S_1$  be a family of subsets of  $X$ , and  $A$  be an element of  $S_1$ . The functor  $\text{Block}(A)$  yielding a subset of  $X \times X$  is defined by the term

(Def. 28)  $(X \setminus A) \times (X \setminus A) \cup A \times A$ .

From now on  $S_1$  denotes a family of subsets of  $X$  and  $A$  denotes an element of  $S_1$ .

Now we state the propositions:

(65) If  $A = \emptyset$ , then  $\text{Block}(A) = X \times X$ .

(66) Suppose  $X$  is not empty. Then  $\text{Block}(A) = \{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } X : x \in A \text{ iff } y \in A\}$ .

PROOF: Set  $S = \{\langle x, y \rangle, \text{ where } x, y \text{ are elements of } X : x \in A \text{ iff } y \in A\}$ .  $\text{Block}(A) \subseteq S$  by [3, (87)].  $S \subseteq \text{Block}(A)$  by [3, (87)].  $\square$

(67) (i)  $\text{id}_X \subseteq \text{Block}(A)$ , and

(ii)  $\text{Block}(A) \cdot \text{Block}(A) \subseteq \text{Block}(A)$ .

Let  $X$  be a set and  $S_1$  be a family of subsets of  $X$ . The functor  $\text{Blocks}(S_1)$  yielding a family of subsets of  $X \times X$  is defined by the term

(Def. 29) the set of all  $\text{Block}(A)$  where  $A$  is an element of  $S_1$ .

Let us observe that  $\text{Blocks}(S_1)$  is non empty.

The functor  $\text{FMCBLOCKS}(S_1)$  yielding a family of subsets of  $X \times X$  is defined by the term

(Def. 30)  $\text{FinMeetCl}(\text{Blocks}(S_1))$ .

Now we state the propositions:

(68)  $\text{FMCBLOCKS}(S_1)$  is  $\cap$ -closed.

(69)  $\text{FMCBLOCKS}(S_1)$  is quasi-basis. The theorem is a consequence of (68).

(70)  $\text{FMCBLOCKS}(S_1)$  satisfies axiom UP1.

(71) Let us consider an element  $A$  of  $S_1$ , and a binary relation  $R$  on  $X$ . If  $R = \text{Block}(A)$ , then  $R^\sim = \text{Block}(A)$ . The theorem is a consequence of (65) and (4).

(72) Let us consider a binary relation  $R$  on  $X$ . Suppose  $R$  is an element of  $\text{Blocks}(S_1)$ . Then  $R^\sim$  is an element of  $\text{Blocks}(S_1)$ . The theorem is a consequence of (71).

Let us consider a non empty family  $Y$  of subsets of  $X \times X$ . Now we state the propositions:

(73) If  $Y \subseteq \text{Blocks}(S_1)$ , then  $Y[\sim] = Y$ . The theorem is a consequence of (71).

- (74) If  $Y \subseteq \text{Blocks}(S_1)$ , then  $(\bigcap Y)^\smile = \bigcap Y [\sim]$ . The theorem is a consequence of (73) and (71).
- (75) If  $Y \subseteq \text{Blocks}(S_1)$ , then  $\bigcap Y = (\bigcap Y)^\smile$ . The theorem is a consequence of (73) and (74).
- (76)  $\text{FMCBlocks}(S_1)$  satisfies axiom UP2. The theorem is a consequence of (73) and (75).
- (77)  $\text{FMCBlocks}(S_1)$  satisfies axiom UP3. The theorem is a consequence of (67).

Let  $X$  be a set and  $S_1$  be a family of subsets of  $X$ . The Pervin uniform space of  $S_1$  yielding a strict uniform space is defined by the term

(Def. 31)  $\langle X, [\text{FMCBlocks}(S_1)] \rangle$ .

ACKNOWLEDGEMENT: The author wants to express his gratitude to the anonymous referee for his/her work, to make the presentation more readable.

#### REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [3] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [4] Mai Gehrke, Serge Grigorieff, and Jean-Éric Pin. A topological approach to recognition. In *Automata, Languages and Programming*, pages 151–162. Springer, 2010.
- [5] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Formalized Mathematics*, 1(3):607–610, 1990.
- [6] Beata Padlewska. Locally connected spaces. *Formalized Mathematics*, 2(1):93–96, 1991.
- [7] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [8] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [9] Wojciech A. Trybulec. Subgroup and cosets of subgroups. *Formalized Mathematics*, 1(5):855–864, 1990.
- [10] Milan Vlach. Algebraic and topological aspects of rough set theory. In *Fourth International Workshop on Computational Intelligence & Applications, IEEE SMC Hiroshima Chapter, Hiroshima University, Japan, December 10&11, 2008*.
- [11] Milan Vlach. Topologies of approximation spaces of rough set theory. In *Integral/Probabilistic Uncertainty and Non-Classical Logics*, pages 176–186. Springer, 2008.

Received June 30, 2016



The English version of this volume of *Formalized Mathematics* was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.

# Some Algebraic Properties of Polynomial Rings

Christoph Schwarzweller  
Institute of Computer Science  
University of Gdańsk  
Poland

Artur Korniłowicz  
Institute of Informatics  
University of Białystok  
Poland

Agnieszka Rowińska-Schwarzweller  
Sopot, Poland

**Summary.** In this article we extend the algebraic theory of polynomial rings, formalized in Mizar [1], based on [2], [3]. After introducing constant and monic polynomials we present the canonical embedding of  $R$  into  $R[X]$  and deal with both unit and irreducible elements. We also define polynomial GCDs and show that for fields  $F$  and irreducible polynomials  $p$  the field  $F[X]/\langle p \rangle$  is isomorphic to the field of polynomials with degree smaller than the one of  $p$ .

MSC: 12E05 11T55 03B35

Keywords: polynomial; polynomial ring; polynomial GCD

MML identifier: RING\_4, version: 8.1.05 5.37.1275

## 1. PRELIMINARIES

Let  $R$  be a non empty double loop structure and  $a$  be an element of  $R$ . Observe that the functor  $\{a\}$  yields a subset of  $R$ . Observe that every ring which is almost left invertible and commutative is also almost right invertible and every ring which is almost right invertible and commutative is also almost left invertible and every ring which is almost left cancelable and commutative is also almost right cancelable and every ring which is almost right cancelable and commutative is also almost left cancelable.

Let  $L$  be a non empty zero structure and  $X$  be a set. We say that  $X$  is  $L$ -polynomial membered if and only if

(Def. 1) for every object  $p$  such that  $p \in X$  holds  $p$  is a polynomial over  $L$ .

Let  $X$  be a 1-sorted structure. We say that  $X$  is  $L$ -polynomial membered if and only if

(Def. 2) the carrier of  $X$  is  $L$ -polynomial membered.

Let us note that there exists a set which is non empty and  $L$ -polynomial membered and there exists a 1-sorted structure which is non empty and  $L$ -polynomial membered.

Let  $X$  be a non empty,  $L$ -polynomial membered 1-sorted structure. One can check that the carrier of  $X$  is  $L$ -polynomial membered.

Let  $L$  be an add-associative, right zeroed, right complementable, distributive, non empty double loop structure. Let us observe that  $\text{Polynom-Ring}(L)$  is  $L$ -polynomial membered.

Let  $L$  be a non empty zero structure and  $X$  be a non empty,  $L$ -polynomial membered set.

Observe that an element of  $X$  is a polynomial over  $L$ . Let  $R$  be a ring. One can verify that there exists an element of the carrier of  $\text{Polynom-Ring}(R)$  which is zero and there exists an element of  $\text{Polynom-Ring}(R)$  which is zero and there exists a polynomial over  $R$  which is zero.

Let  $R$  be a non degenerated ring. Let us note that there exists an element of the carrier of  $\text{Polynom-Ring}(R)$  which is non zero and there exists an element of  $\text{Polynom-Ring}(R)$  which is non zero.

Let  $L$  be an add-associative, right zeroed, right complementable, distributive, non empty double loop structure and  $p, q$  be polynomials over  $L$ . We say that  $p \mid q$  if and only if

(Def. 3) there exist elements  $a, b$  of  $\text{Polynom-Ring}(L)$  such that  $a = p$  and  $b = q$  and  $a \mid b$ .

Now we state the proposition:

- (1) Let us consider an add-associative, right zeroed, right complementable, distributive, non empty double loop structure  $L$ , and polynomials  $p, q$  over  $L$ . Then  $p \mid q$  if and only if there exists a polynomial  $r$  over  $L$  such that  $p * r = q$ .

Let us consider a field  $F$  and polynomials  $p, q$  over  $F$ . Now we state the propositions:

- (2) If  $\deg p < \deg q$ , then  $p \bmod q = p$ .
- (3)  $p \bmod q = \mathbf{0}$  if and only if  $q \mid p$ . The theorem is a consequence of (1).
- (4)  $p = (p \text{ div } q) * q + (p \bmod q)$ .

Let us consider a field  $F$ , polynomials  $p, r$  over  $F$ , and a non zero polynomial  $q$  over  $F$ . Now we state the propositions:



- (5) (i)  $p + r \operatorname{div} q = (p \operatorname{div} q) + (r \operatorname{div} q)$ , and  
 (ii)  $p + r \operatorname{mod} q = (p \operatorname{mod} q) + (r \operatorname{mod} q)$ .

The theorem is a consequence of (4).

- (6)  $p * r \operatorname{mod} q = (p \operatorname{mod} q) * (r \operatorname{mod} q) \operatorname{mod} q$ . The theorem is a consequence of (4), (5), (3), and (1).

Now we state the propositions:

- (7) Let us consider a field  $F$ , polynomials  $r, q, u$  over  $F$ , and a non zero polynomial  $p$  over  $F$ . Then  $(r * q \operatorname{mod} p) * u \operatorname{mod} p = (r * q) * u \operatorname{mod} p$ . The theorem is a consequence of (5), (3), and (1).
- (8) Let us consider an add-associative, right zeroed, right complementable, left distributive, non empty double loop structure  $L$ , and a sequence  $p$  of  $L$ . Then  $0_L \cdot p = \mathbf{0} \cdot L$ .
- (9) Let us consider a left unital, non empty double loop structure  $L$ , and a sequence  $p$  of  $L$ . Then  $1_L \cdot p = p$ .
- (10) Let us consider an add-associative, right zeroed, right complementable, right unital, distributive, associative, commutative, non empty double loop structure  $L$ , sequences  $p, q$  of  $L$ , and an element  $a$  of  $L$ . Then  $a \cdot (p * q) = p * (a \cdot q)$ .
- (11) Let us consider an associative, non empty multiplicative magma  $L$ , a sequence  $p$  of  $L$ , and elements  $a, b$  of  $L$ . Then  $(a \cdot b) \cdot p = a \cdot (b \cdot p)$ .
- (12) Let us consider an add-associative, right zeroed, right complementable, left distributive, left unital, non empty double loop structure  $L$ , and a sequence  $p$  of  $L$ . Then  $\mathbf{1} \cdot L * p = p$ .

Let  $L$  be an add-associative, right zeroed, right complementable, well unital, distributive, non empty double loop structure. Let us observe that  $\operatorname{Polynom-Ring}(L)$  is well unital.

## 2. CONSTANT POLYNOMIALS

Let  $R$  be an add-associative, right zeroed, right complementable, distributive, non empty double loop structure and  $x$  be an element of the carrier of  $\operatorname{Polynom-Ring}(R)$ . We say that  $x$  is constant if and only if

(Def. 4)  $\deg x \leq 0$ .

Let  $R$  be a non degenerated ring. Observe that there exists an element of  $\operatorname{Polynom-Ring}(R)$  which is non zero and constant and there exists an element of the carrier of  $\operatorname{Polynom-Ring}(R)$  which is non zero and constant.

Let  $R$  be an integral domain. Let us observe that there exists an element of  $\text{Polynom-Ring}(R)$  which is non constant and there exists an element of the carrier of  $\text{Polynom-Ring}(R)$  which is non constant.

Let  $L$  be a non empty zero structure and  $a$  be an element of  $L$ . The functor  $a\downarrow L$  yielding a sequence of  $L$  is defined by the term

(Def. 5)  $\mathbf{0}.L + \cdot (0, a)$ .

Note that  $a\downarrow L$  is finite-Support and  $a\downarrow L$  is constant.

Let  $a$  be a non zero element of  $L$ . Let us note that  $a\downarrow L$  is non zero and there exists a polynomial over  $L$  which is non zero and constant.

Now we state the propositions:

(13) Let us consider a non empty zero structure  $L$ . Then  $0_L\downarrow L = \mathbf{0}.L$ .

(14) Let us consider a non empty multiplicative loop with zero structure  $L$ .  
Then  $1_L\downarrow L = \mathbf{1}.L$ .

Let  $L$  be a non empty zero structure. Observe that  $0_L\downarrow L$  is zero.

Let  $L$  be a non degenerated multiplicative loop with zero structure. Let us note that  $1_L\downarrow L$  is non zero.

Now we state the propositions:

(15) Let us consider an add-associative, right zeroed, right complementable, distributive, non empty double loop structure  $L$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(L)$ . Then  $p$  is non zero and constant if and only if  $\deg p = 0$ .

(16) Let us consider an add-associative, right zeroed, right complementable, right distributive, right unital, non empty double loop structure  $L$ , and an element  $a$  of  $L$ . Then  $a\downarrow L = a \cdot \mathbf{1}.L$ .

Let us consider a ring  $R$  and elements  $a, b$  of  $R$ . Now we state the propositions:

(17)  $a\downarrow R + b\downarrow R = (a + b)\downarrow R$ .

(18)  $(a\downarrow R) * (b\downarrow R) = a \cdot b\downarrow R$ .

(19)  $a\downarrow R = b\downarrow R$  if and only if  $a = b$ .

(20) Let us consider a ring  $R$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(R)$ . Then  $p$  is constant if and only if there exists an element  $a$  of  $R$  such that  $p = a\downarrow R$ .

(21) Let us consider a ring  $R$ , and an element  $a$  of  $R$ . Then  $\deg(a\downarrow R) = 0$  if and only if  $a \neq 0_R$ . The theorem is a consequence of (19).

## 3. MONIC POLYNOMIALS

Let  $L$  be a non empty double loop structure and  $p$  be a polynomial over  $L$ . We introduce the notation  $p$  is monic as a synonym of  $p$  is normalized.

Let  $L$  be an add-associative, right zeroed, right complementable, distributive, non degenerated double loop structure. Let us observe that  $\mathbf{1}.L$  is monic and  $\mathbf{0}.L$  is non monic and there exists a polynomial over  $L$  which is monic and there exists a polynomial over  $L$  which is non monic and there exists an element of the carrier of  $\text{Polynom-Ring}(L)$  which is monic and there exists an element of the carrier of  $\text{Polynom-Ring}(L)$  which is non monic.

Let  $L$  be a well unital, non degenerated double loop structure and  $x$  be an element of  $L$ . One can verify that  $\text{rpoly}(1, x)$  is monic.

Let  $L$  be a field and  $p$  be an element of the carrier of  $\text{Polynom-Ring}(L)$ . Let us observe that the functor  $\text{NormPolynomial } p$  yields an element of the carrier of  $\text{Polynom-Ring}(L)$ . Let  $F$  be a field and  $p$  be a non zero polynomial over  $F$ . Observe that  $\text{NormPolynomial } p$  is monic.

Let  $L$  be a field and  $p$  be a non zero element of the carrier of  $\text{Polynom-Ring}(L)$ . Observe that  $\text{NormPolynomial } p$  is monic.

Now we state the proposition:

(22) Let us consider a field  $F$ . Then  $\text{NormPolynomial } \mathbf{0}.F = \mathbf{0}.F$ .

Let us consider a field  $F$  and a non zero element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Now we state the propositions:

(23)  $\text{NormPolynomial } p = (\text{LC } p)^{-1} \cdot p$ .

(24)  $p$  is monic if and only if  $\text{NormPolynomial } p = p$ . The theorem is a consequence of (23) and (9).

Let us consider a field  $F$  and elements  $p, q$  of the carrier of  $\text{Polynom-Ring}(F)$ . Now we state the propositions:

(25)  $q \mid p$  if and only if  $\text{NormPolynomial } q \mid p$ . The theorem is a consequence of (22), (1), (9), (11), (10), and (23).

(26)  $q \mid p$  if and only if  $q \mid \text{NormPolynomial } p$ . The theorem is a consequence of (22), (1), (23), (10), (9), and (11).

Let us consider a field  $F$  and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Now we state the propositions:

(27)  $\text{NormPolynomial } p$  is associated to  $p$ . The theorem is a consequence of (1), (26), and (25).

(28)  $\text{NormPolynomial } p$  is irreducible if and only if  $p$  is irreducible. The theorem is a consequence of (27).

Now we state the propositions:

- (29) Let us consider an integral domain  $R$ , and elements  $p, q$  of the carrier of  $\text{Polynom-Ring}(R)$ . If  $p$  is associated to  $q$ , then  $\deg p = \deg q$ .
- (30) Let us consider an integral domain  $R$ , and monic elements  $p, q$  of the carrier of  $\text{Polynom-Ring}(R)$ . Then  $p$  is associated to  $q$  if and only if  $p = q$ . The theorem is a consequence of (29), (20), (16), (10), and (12).

#### 4. THE CANONICAL HOMOMORPHISM FROM $R$ INTO $R[X]$

Let  $R$  be a ring. The canonical homomorphism of  $R$  into quotient field yielding a function from  $R$  into  $\text{Polynom-Ring}(R)$  is defined by

(Def. 6) for every element  $x$  of  $R$ ,  $it(x) = x \upharpoonright R$ .

Note that the canonical homomorphism of  $R$  into quotient field is additive, multiplicative, and unity-preserving and the canonical homomorphism of  $R$  into quotient field is monomorphic and  $\text{Polynom-Ring}(R)$  is  $R$ -homomorphic and  $R$ -monomorphic.

Now we state the proposition:

- (31) Let us consider a ring  $R$ . Then  $\text{char}(\text{Polynom-Ring}(R)) = \text{char}(R)$ .

Let  $R$  be a non degenerated ring. Let us note that  $\text{Polynom-Ring}(R)$  is infinite and every ring with characteristic 0 is infinite.

Now we state the proposition:

- (32) Let us consider a ring  $R$ . If  $\text{char}(R) = 0$ , then  $R$  is infinite.

Let  $n$  be a non trivial natural number.

One can verify that  $\text{Polynom-Ring}(\mathbb{Z}/n)$  is infinite. Now we state the proposition:

- (33) Let us consider a non trivial natural number  $n$ .

Then  $\text{char}(\text{Polynom-Ring}(\mathbb{Z}/n)) \neq 0$ .

Let  $n$  be a non trivial natural number. Observe that there exists a ring which is infinite and has characteristic  $n$ .

#### 5. UNITS AND IRREDUCIBLE POLYNOMIALS

Let us note that there exists an integral domain which is non almost left invertible.

Let  $R$  be a non almost left invertible integral domain. One can verify that there exists a non-unit of  $R$  which is non zero and  $\mathbb{Z}^R$  is non almost left invertible.

Let  $R$  be an integral domain. Observe that  $\text{Polynom-Ring}(R)$  is non almost left invertible.

Now we state the propositions:

- (34) Let us consider an integral domain  $R$ . Then  $R$  is a field if and only if for every non-unit  $a$  of  $R$ ,  $a = 0_R$ .
- (35) Let us consider an integral domain  $R$ , and an element  $a$  of  $R$ . Then  $a \mid R$  is a unit of  $\text{Polynom-Ring}(R)$  if and only if  $a$  is a unit of  $R$ . The theorem is a consequence of (1), (20), (18), and (19).
- (36) Let us consider an integral domain  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . If  $p$  is a unit of  $\text{Polynom-Ring}(F)$ , then  $\deg p = 0$ . The theorem is a consequence of (1).
- (37) Let us consider a field  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Then  $p$  is a unit of  $\text{Polynom-Ring}(F)$  if and only if  $\deg p = 0$ . The theorem is a consequence of (1), (20), and (18).
- (38) Let us consider an integral domain  $R$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(R)$ . Suppose  $p$  is a unit of  $\text{Polynom-Ring}(R)$ . Then  $p$  is non zero and constant. The theorem is a consequence of (36) and (15).
- (39) Let us consider a field  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Then  $p$  is a unit of  $\text{Polynom-Ring}(F)$  if and only if  $p$  is non zero and constant. The theorem is a consequence of (37) and (15).

Let  $R$  be an integral domain. One can check that every element of  $\text{Polynom-Ring}(R)$  which is non constant is also non zero and non unital.

Let  $F$  be an integral domain. Let us observe that every element of the carrier of  $\text{Polynom-Ring}(F)$  which is non constant is also non zero and non unital.

Let  $F$  be a field. Observe that every element of  $\text{Polynom-Ring}(F)$  which is non zero and constant is also unital and every element of  $\text{Polynom-Ring}(F)$  which is unital is also non zero and constant and every element of the carrier of  $\text{Polynom-Ring}(F)$  which is non zero and constant is also unital and every element of the carrier of  $\text{Polynom-Ring}(F)$  which is unital is also non zero and constant.

Now we state the propositions:

- (40) Let us consider an integral domain  $R$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(R)$ . Suppose there exists an element  $q$  of the carrier of  $\text{Polynom-Ring}(R)$  such that  $q \mid p$  and  $1 \leq \deg q < \deg p$ . Then  $p$  is reducible. The theorem is a consequence of (36).
- (41) Let us consider a field  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Then  $p$  is reducible if and only if  $p = \mathbf{0}_F$  or  $p$  is a unit of  $\text{Polynom-Ring}(F)$  or there exists an element  $q$  of the carrier of  $\text{Polynom-Ring}(F)$  such that  $q \mid p$  and  $1 \leq \deg q < \deg p$ . The theorem is a consequence of (1), (37), and (40).
- (42) Let us consider an integral domain  $R$ , and a monic element  $p$  of the carrier of  $\text{Polynom-Ring}(R)$ . If  $\deg p = 1$ , then  $p$  is irreducible. The theorem

is a consequence of (36), (20), (16), (10), (12), and (35).

(43) There exists a non monic element  $p$  of the carrier of  $\text{Polynom-Ring}(\mathbb{Z}^{\mathbb{R}})$  such that

(i)  $\deg p = 1$ , and

(ii)  $p$  is reducible.

The theorem is a consequence of (16), (10), (12), (15), (35), and (36).

(44) Let us consider a field  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . If  $\deg p = 1$ , then  $p$  is irreducible. The theorem is a consequence of (36), (20), (21), and (35).

(45) Let us consider an algebraic closed field  $F$ , and an element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Then  $p$  is irreducible if and only if  $\deg p = 1$ . The theorem is a consequence of (36) and (44).

(46) Let us consider a field  $F$ . Then  $F$  is algebraic closed if and only if for every monic element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ ,  $p$  is irreducible iff  $\deg p = 1$ . The theorem is a consequence of (37), (41), (28), and (45).

Let  $R$  be an integral domain. Note that there exists an element of  $\text{Polynom-Ring}(R)$  which is irreducible and there exists an element of the carrier of  $\text{Polynom-Ring}(R)$  which is irreducible.

Let  $R$  be a ring. Let us observe that there exists an element of  $\text{Polynom-Ring}(R)$  which is reducible and there exists an element of the carrier of  $\text{Polynom-Ring}(R)$  which is reducible. Let  $R$  be an integral domain.

Note that  $\text{IRR}(\text{Polynom-Ring}(R))$  is non empty.

Let  $F$  be a field. Observe that every element of  $\text{Polynom-Ring}(F)$  which is constant is also reducible and every element of the carrier of  $\text{Polynom-Ring}(F)$  which is constant is also reducible and every element of  $\text{Polynom-Ring}(F)$  which is irreducible is also non constant and every element of the carrier of  $\text{Polynom-Ring}(F)$  which is irreducible is also non constant.

## 6. THE FIELD $F[X]/\langle p \rangle$

Let  $F$  be a field and  $p$  be an element of the carrier of  $\text{Polynom-Ring}(F)$ . Let us note that  $\frac{\text{Polynom-Ring}(F)}{\{p\}\text{-ideal}}$  is Abelian, add-associative, right zeroed, right complementable, commutative, associative, well unital, and distributive.

Let  $p$  be an irreducible element of the carrier of  $\text{Polynom-Ring}(F)$ . Observe that  $\frac{\text{Polynom-Ring}(F)}{\{p\}\text{-ideal}}$  is non degenerated and almost left invertible.

Let  $p$  be a polynomial over  $F$ . The functor  $\text{PolyMultMod}(p)$  yielding a binary operation on  $\text{Polynom-Ring}(F)$  is defined by

(Def. 7) for every polynomials  $r, q$  over  $F$ ,  $it(r, q) = r * q \text{ mod } p$ .

Let  $p$  be a non constant element of the carrier of  $\text{Polynom-Ring}(F)$ . The functor  $\text{Polynom-Ring}(p)$  yielding a strict double loop structure is defined by

- (Def. 8) the carrier of  $it = \{q, \text{ where } q \text{ is a polynomial over } F : \deg q < \deg p\}$  and the addition of  $it = (\text{the addition of } \text{Polynom-Ring}(F)) \upharpoonright (\text{the carrier of } it)$  and the multiplication of  $it = \text{PolyMultMod}(p) \upharpoonright (\text{the carrier of } it)$  and the one of  $it = \mathbf{1}.F$  and the zero of  $it = \mathbf{0}.F$ .

Observe that  $\text{Polynom-Ring}(p)$  is non degenerated and  $\text{Polynom-Ring}(p)$  is Abelian, add-associative, right zeroed, and right complementable and  $\text{Polynom-Ring}(p)$  is associative, well unital, and distributive.

The functor  $\text{PolyMod}(p)$  yielding a function from  $\text{Polynom-Ring}(F)$  into  $\text{Polynom-Ring}(p)$  is defined by

- (Def. 9) for every polynomial  $q$  over  $F$ ,  $it(q) = q \text{ mod } p$ .

Observe that  $\text{PolyMod}(p)$  is additive, multiplicative, and unity-preserving and  $\text{Polynom-Ring}(p)$  is  $(\text{Polynom-Ring}(F))$ -homomorphic and  $\text{PolyMod}(p)$  is onto.

Let us consider a field  $F$  and a non constant element  $p$  of the carrier of  $\text{Polynom-Ring}(F)$ . Now we state the propositions:

- (47)  $\ker \text{PolyMod}(p) = \{p\}$ -ideal. The theorem is a consequence of (1) and (3).  
 (48)  $\frac{\text{Polynom-Ring}(F)}{\{p\}\text{-ideal}}$  and  $\text{Polynom-Ring}(p)$  are isomorphic. The theorem is a consequence of (47).

Let  $F$  be a field and  $p$  be a non constant element of the carrier of  $\text{Polynom-Ring}(F)$ . Observe that  $\text{Polynom-Ring}(p)$  is commutative.

Let  $p$  be an irreducible element of the carrier of  $\text{Polynom-Ring}(F)$ . Observe that  $\text{Polynom-Ring}(p)$  is almost left invertible.

### 7. POLYNOMIAL GCDs

Let  $L$  be a non empty multiplicative magma,  $x, y$  be elements of  $L$ , and  $z$  be an element of  $L$ . We say that  $z$  is  $x,y$ -GCD if and only if

- (Def. 10)  $z \mid x$  and  $z \mid y$  and for every element  $r$  of  $L$  such that  $r \mid x$  and  $r \mid y$  holds  $r \mid z$ .

Let  $L$  be a GCD domain. Note that there exists an element of  $L$  which is  $x,y$ -GCD.

A GCD of  $x$  and  $y$  is an  $x,y$ -GCD element of  $L$ . Now we state the proposition:

- (49) Let us consider a GCD domain  $L$ , elements  $x, y$  of  $L$ , and GCDs  $u, v$  of  $x$  and  $y$ . Then  $u$  is associated to  $v$ .

Let  $L$  be a GCD domain and  $x, y$  be elements of  $L$ . One can verify that every element of  $L$  which is  $x,y$ -GCD is also  $y,x$ -GCD.

Let  $F$  be a field and  $p, q$  be elements of the carrier of  $\text{Polynom-Ring}(F)$ . The functor  $\text{gcd}(p, q)$  yielding an element of the carrier of  $\text{Polynom-Ring}(F)$  is defined by

- (Def. 11) (i)  $it = \mathbf{0}.F$ , **if**  $p = \mathbf{0}.F$  and  $q = \mathbf{0}.F$ ,  
 (ii)  $it$  is GCD of  $p$  and  $q$  and monic, **otherwise**.

One can check that the functor  $\text{gcd}(p, q)$  is commutative.

Let  $p, q$  be elements of  $\text{Polynom-Ring}(F)$ . Let us note that the functor  $\text{gcd}(p, q)$  is commutative.

Let  $p, q$  be elements of the carrier of  $\text{Polynom-Ring}(F)$ . Let us observe that  $\text{gcd}(p, q)$  is  $p,q$ -GCD.

Let  $p, q$  be elements of  $\text{Polynom-Ring}(F)$ . Observe that  $\text{gcd}(p, q)$  is  $p,q$ -GCD.

Let  $p$  be an element of the carrier of  $\text{Polynom-Ring}(F)$  and  $q$  be a non zero element of the carrier of  $\text{Polynom-Ring}(F)$ . Note that  $\text{gcd}(p, q)$  is non zero and monic.

Let  $p$  be an element of  $\text{Polynom-Ring}(F)$  and  $q$  be a non zero element of  $\text{Polynom-Ring}(F)$ . Let us observe that  $\text{gcd}(p, q)$  is non zero and monic.

Let  $p, q$  be zero elements of the carrier of  $\text{Polynom-Ring}(F)$ . Let us note that  $\text{gcd}(p, q)$  is zero.

Let  $p, q$  be zero elements of  $\text{Polynom-Ring}(F)$ . One can verify that  $\text{gcd}(p, q)$  is zero.

Now we state the propositions:

- (50) Let us consider a field  $F$ , and elements  $p, q$  of the carrier of  $\text{Polynom-Ring}(F)$ . Then
- (i)  $\text{gcd}(p, q) \mid p$ , and
  - (ii)  $\text{gcd}(p, q) \mid q$ , and
  - (iii) for every element  $r$  of the carrier of  $\text{Polynom-Ring}(F)$  such that  $r \mid p$  and  $r \mid q$  holds  $r \mid \text{gcd}(p, q)$ .
- (51) Let us consider a field  $F$ , and elements  $p, q$  of  $\text{Polynom-Ring}(F)$ . Then
- (i)  $\text{gcd}(p, q) \mid p$ , and
  - (ii)  $\text{gcd}(p, q) \mid q$ , and
  - (iii) for every element  $r$  of  $\text{Polynom-Ring}(F)$  such that  $r \mid p$  and  $r \mid q$  holds  $r \mid \text{gcd}(p, q)$ .

Let  $F$  be a field and  $p, q$  be polynomials over  $F$ . The functor  $\text{gcd}(p, q)$  yielding a polynomial over  $F$  is defined by



(Def. 12) there exist elements  $a, b$  of Polynom-Ring( $F$ ) such that  $a = p$  and  $b = q$  and  $it = \gcd(a, b)$ .

Observe that the functor  $\gcd(p, q)$  is commutative.

Let  $p$  be a polynomial over  $F$  and  $q$  be a non zero polynomial over  $F$ . Let us note that  $\gcd(p, q)$  is non zero and monic.

Let  $p, q$  be zero polynomials over  $F$ . One can verify that  $\gcd(p, q)$  is zero.

Now we state the propositions:

(52) Let us consider a field  $F$ , and polynomials  $p, q$  over  $F$ . Then

(i)  $\gcd(p, q) \mid p$ , and

(ii)  $\gcd(p, q) \mid q$ , and

(iii) for every polynomial  $r$  over  $F$  such that  $r \mid p$  and  $r \mid q$  holds  $r \mid \gcd(p, q)$ .

The theorem is a consequence of (1).

(53) Let us consider a field  $F$ , a polynomial  $p$  over  $F$ , a non zero polynomial  $q$  over  $F$ , and a monic polynomial  $s$  over  $F$ . Then  $s = \gcd(p, q)$  if and only if  $s \mid p$  and  $s \mid q$  and for every polynomial  $r$  over  $F$  such that  $r \mid p$  and  $r \mid q$  holds  $r \mid s$ . The theorem is a consequence of (52).

#### REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] H. Heuser. *Lehrbuch der Analysis*. B.G. Teubner Stuttgart, 1990.
- [3] Steven H. Weintraub. *Galois Theory*. Springer-Verlag, 2 edition, 2009.

*Received June 30, 2016*

---



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.