

Contents

Formaliz. Math. 26 (4)

Pythagorean Tuning: Pentatonic and Heptatonic Scale By ROLAND COGHETTO	239
Fundamental Properties of Fuzzy Implications By ADAM GRABOWSKI	271
Zariski Topology By YASUSHIGE WATASE	277

Continued on inside back cover

Pythagorean Tuning: Pentatonic and Heptatonic Scale

Roland Coghetto 
Rue de la Brasserie 5
7100 La Louvière, Belgium

Summary. In this article, using the Mizar system [3], [4], we define a structure [1], [6] in order to build a Pythagorean pentatonic scale and a Pythagorean heptatonic scale¹ [5], [7].

MSC: 00A65 97M80 03B35

Keywords: music; Pythagorean tuning; pentatonic scale; heptatonic scale

MML identifier: MUSIC_S1, version: 8.1.08 5.53.1335

1. PRELIMINARIES

Now we state the proposition:

- (1) Let us consider an object r . Then $r \in \mathbb{R}_{+\cup\{0\}} \setminus \{0\}$ if and only if r is a positive real number.

Note that there exists a rational number which is positive.

The functor \mathbb{Q}_+ yielding a non empty subset of $\mathbb{R}_{+\cup\{0\}}$ is defined by the term

- (Def. 1) the set of all r where r is a positive rational number.

Now we state the propositions:

- (2) Let us consider an object r . Then r is an element of \mathbb{Q}_+ if and only if r is a positive rational number.
- (3) $\mathbb{Q}_{+\cup\{0\}} \subseteq \mathbb{Q}$.

¹https://en.wikipedia.org/wiki/Pythagorean_tuning

The functor \mathbb{R}_+ yielding a non empty subset of $\mathbb{R}_{+\cup\{0\}}$ is defined by the term

(Def. 2) $\mathbb{R}_{+\cup\{0\}} \setminus \{0\}$.

Now we state the propositions:

(4) $\mathbb{N}_+ \subseteq \mathbb{Q}_+$.

(5) $\mathbb{N}_+ \subseteq \mathbb{R}_+$. The theorem is a consequence of (1).

(6) $\mathbb{Q}_+ \subseteq \mathbb{R}_+$. The theorem is a consequence of (2) and (1).

2. REAL FREQUENCY

We consider structures of music which extend 1-sorted structures and are systems

\langle a carrier, an equidistance, a Ratio \rangle

where the carrier is a set, the equidistance is a relation between (the carrier) \times (the carrier) and (the carrier) \times (the carrier), the Ratio is a function from (the carrier) \times (the carrier) into the carrier.

Let S be a structure of music and a, b, c, d be elements of S . We say that $\overline{ab} \cong \overline{cd}$ if and only if

(Def. 3) $\langle\langle a, b \rangle, \langle c, d \rangle\rangle \in$ the equidistance of S .

Let x, y be elements of \mathbb{R}_+ . The functor \mathbb{R} -ratio(x, y) yielding an element of \mathbb{R}_+ is defined by

(Def. 4) there exist positive real numbers r, s such that $x = r$ and $s = y$ and $it = \frac{s}{r}$.

Now we state the proposition:

(7) Let us consider elements a, b, c, d of \mathbb{R}_+ . Then \mathbb{R} -ratio(a, b) = \mathbb{R} -ratio(c, d) if and only if \mathbb{R} -ratio(b, a) = \mathbb{R} -ratio(d, c).

The functor \mathbb{R} -ratio yielding a function from $\mathbb{R}_+ \times \mathbb{R}_+$ into \mathbb{R}_+ is defined by

(Def. 5) for every element x of $\mathbb{R}_+ \times \mathbb{R}_+$, there exist elements y, z of \mathbb{R}_+ such that $x = \langle y, z \rangle$ and $it(x) = \mathbb{R}$ -ratio(y, z).

The functor eq- \mathbb{R} -ratio yielding a relation between $\mathbb{R}_+ \times \mathbb{R}_+$ and $\mathbb{R}_+ \times \mathbb{R}_+$ is defined by

(Def. 6) for every elements x, y of $\mathbb{R}_+ \times \mathbb{R}_+$, $\langle x, y \rangle \in it$ iff there exist elements a, b, c, d of \mathbb{R}_+ such that $x = \langle a, b \rangle$ and $y = \langle c, d \rangle$ and \mathbb{R} -ratio(a, b) = \mathbb{R} -ratio(c, d).

The functor \mathbb{R} -music yielding a structure of music is defined by the term

(Def. 7) $\langle \mathbb{R}_+, \text{eq-}\mathbb{R}\text{-ratio}, \mathbb{R}\text{-ratio} \rangle$.

Now we state the propositions:

- (8) (i) \mathbb{R} -music is not empty, and
 - (ii) the carrier of \mathbb{R} -music $\subseteq \mathbb{R}_+$, and
 - (iii) for every elements f_1, f_2, f_3, f_4 of \mathbb{R} -music, $\overline{f_1 f_2} \cong \overline{f_3 f_4}$ iff (the Ratio of \mathbb{R} -music)(f_1, f_2) = (the Ratio of \mathbb{R} -music)(f_3, f_4).
- (9) Let us consider elements f_1, f_2, f_3 of \mathbb{R} -music. Suppose (the Ratio of \mathbb{R} -music)(f_1, f_2) = (the Ratio of \mathbb{R} -music)(f_1, f_3). Then $f_2 = f_3$.
- (10) $\mathbb{N}_+ \subseteq$ the carrier of \mathbb{R} -music.
- (11) Let us consider an element fr of \mathbb{R} -music, and a non zero natural number n . Then there exists an element h of \mathbb{R} -music such that $\langle fr, h \rangle \in [\langle 1, n \rangle]_\alpha$, where α is the equidistance of \mathbb{R} -music. The theorem is a consequence of (1) and (8).
- (12) Let us consider elements f_1, f_2, f_3 of \mathbb{R} -music. Suppose (the Ratio of \mathbb{R} -music)(f_1, f_1) = (the Ratio of \mathbb{R} -music)(f_2, f_3). Then $f_2 = f_3$.
- (13) Let us consider elements $f_1, f_2, f_8, f_6, f_9, f_7$ of \mathbb{R} -music, positive real numbers r_1, r_2 , and non zero natural numbers n, m . Suppose $f_8 = n \cdot r_1$ and $f_6 = m \cdot r_1$ and $f_9 = n \cdot r_2$ and $f_7 = m \cdot r_2$. Then $\overline{f_8 f_6} \cong \overline{f_9 f_7}$. The theorem is a consequence of (8).
- (14) Let us consider elements f_1, f_2, f_3, f_4 of \mathbb{R} -music. Then (the Ratio of \mathbb{R} -music)(f_1, f_2) = (the Ratio of \mathbb{R} -music)(f_3, f_4) if and only if (the Ratio of \mathbb{R} -music)(f_2, f_1) = (the Ratio of \mathbb{R} -music)(f_4, f_3). The theorem is a consequence of (7).

3. RATIONAL FREQUENCY

Let x, y be elements of \mathbb{Q}_+ . The functor \mathbb{Q} -ratio(x, y) yielding an element of \mathbb{Q}_+ is defined by

(Def. 8) there exist positive rational numbers r, s such that $x = r$ and $s = y$ and $it = \frac{s}{r}$.

Now we state the proposition:

- (15) Let us consider elements a, b, c, d of \mathbb{Q}_+ . Then \mathbb{Q} -ratio(a, b) = \mathbb{Q} -ratio(c, d) if and only if \mathbb{Q} -ratio(b, a) = \mathbb{Q} -ratio(d, c).

The functor \mathbb{Q} -ratio yielding a function from $\mathbb{Q}_+ \times \mathbb{Q}_+$ into \mathbb{Q}_+ is defined by

(Def. 9) for every element x of $\mathbb{Q}_+ \times \mathbb{Q}_+$, there exist elements y, z of \mathbb{Q}_+ such that $x = \langle y, z \rangle$ and $it(x) = \mathbb{Q}$ -ratio(y, z).

The functor eq- \mathbb{Q} -ratio yielding a relation between $\mathbb{Q}_+ \times \mathbb{Q}_+$ and $\mathbb{Q}_+ \times \mathbb{Q}_+$ is defined by

(Def. 10) for every elements x, y of $\mathbb{Q}_+ \times \mathbb{Q}_+$, $\langle x, y \rangle \in it$ iff there exist elements a, b, c, d of \mathbb{Q}_+ such that $x = \langle a, b \rangle$ and $y = \langle c, d \rangle$ and $\mathbb{Q}\text{-ratio}(a, b) = \mathbb{Q}\text{-ratio}(c, d)$.

The functor \mathbb{Q} -music yielding a structure of music is defined by the term

(Def. 11) $\langle \mathbb{Q}_+, \text{eq-}\mathbb{Q}\text{-ratio}, \mathbb{Q}\text{-ratio} \rangle$.

Now we state the propositions:

- (16) (i) \mathbb{Q} -music is not empty, and
 (ii) the carrier of \mathbb{Q} -music $\subseteq \mathbb{R}_+$, and
 (iii) for every elements f_1, f_2, f_3, f_4 of \mathbb{Q} -music, $\overline{f_1 f_2} \cong \overline{f_3 f_4}$ iff (the Ratio of \mathbb{Q} -music)(f_1, f_2) = (the Ratio of \mathbb{Q} -music)(f_3, f_4).

The theorem is a consequence of (6).

- (17) Let us consider elements f_1, f_2, f_3 of \mathbb{Q} -music. Suppose (the Ratio of \mathbb{Q} -music)(f_1, f_2) = (the Ratio of \mathbb{Q} -music)(f_1, f_3). Then $f_2 = f_3$.
- (18) $\mathbb{N}_+ \subseteq$ the carrier of \mathbb{Q} -music.
- (19) Let us consider an element fr of \mathbb{Q} -music, and a non zero natural number n . Then there exists an element h of \mathbb{Q} -music such that $\langle fr, h \rangle \in [\langle 1, n \rangle]_\alpha$, where α is the equidistance of \mathbb{Q} -music. The theorem is a consequence of (2) and (16).
- (20) Let us consider elements f_1, f_2, f_3 of \mathbb{Q} -music. Suppose (the Ratio of \mathbb{Q} -music)(f_1, f_1) = (the Ratio of \mathbb{Q} -music)(f_2, f_3). Then $f_2 = f_3$.
- (21) Let us consider an element fr of \mathbb{Q} -music. Then there exists a positive real number r such that
 (i) $fr = r$, and
 (ii) for every non zero natural number n , $n \cdot r$ is an element of \mathbb{Q} -music.

The theorem is a consequence of (2).

- (22) Let us consider elements $f_1, f_2, f_8, f_6, f_9, f_7$ of \mathbb{Q} -music, positive rational numbers r_1, r_2 , and non zero natural numbers n, m . Suppose $f_8 = n \cdot r_1$ and $f_6 = m \cdot r_1$ and $f_9 = n \cdot r_2$ and $f_7 = m \cdot r_2$. Then $\overline{f_8 f_6} \cong \overline{f_9 f_7}$. The theorem is a consequence of (16).
- (23) Let us consider elements f_1, f_2, f_3, f_4 of \mathbb{Q} -music. Then (the Ratio of \mathbb{Q} -music)(f_1, f_2) = (the Ratio of \mathbb{Q} -music)(f_3, f_4) if and only if (the Ratio of \mathbb{Q} -music)(f_2, f_1) = (the Ratio of \mathbb{Q} -music)(f_4, f_3). The theorem is a consequence of (15).

4. MUSICAL STRUCTURE AND SOME AXIOMS

Let S be a structure of music. We say that S is satisfying real if and only if
 (Def. 12) the carrier of $S \subseteq \mathbb{R}_+$.

We say that S is equidistant-ratio equivalent if and only if

(Def. 13) for every elements f_1, f_2, f_3, f_4 of S , $\overline{f_1 f_2} \cong \overline{f_3 f_4}$ iff (the Ratio of S)(f_1, f_2) = (the Ratio of S)(f_3, f_4).

We say that S is satisfying interval if and only if

(Def. 14) for every elements f_1, f_2, f_3 of S such that (the Ratio of S)(f_1, f_2) = (the Ratio of S)(f_1, f_3) holds $f_2 = f_3$.

We say that S is unison-ratio stable if and only if

(Def. 15) for every elements f_1, f_2, f_3 of S such that (the Ratio of S)(f_1, f_1) = (the Ratio of S)(f_2, f_3) holds $f_2 = f_3$.

We say that S is ratio symmetric if and only if

(Def. 16) for every elements f_1, f_2, f_3, f_4 of S , (the Ratio of S)(f_1, f_2) = (the Ratio of S)(f_3, f_4) iff (the Ratio of S)(f_2, f_1) = (the Ratio of S)(f_4, f_3).

We say that S is natural if and only if

(Def. 17) $\mathbb{N}_+ \subseteq$ the carrier of S .

We say that S is harmonic closed if and only if

(Def. 18) for every element fr of S and for every non zero natural number n , there exists an element h of S such that $\langle fr, h \rangle \in [\langle 1, n \rangle]_\alpha$, where α is the equidistance of S .

Note that there exists a structure of music which is harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, and non empty.

Let us note that the functor \mathbb{R} -music yields a harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Observe that the functor \mathbb{Q} -music yields a harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Now we state the propositions:

(24) Let us consider a natural structure of music S . Then every non zero natural number is an element of S .

(25) Let us consider an equidistant-ratio equivalent structure of music M , and elements a, b of M . Then $\overline{ab} \cong \overline{ab}$.

(26) Let us consider an equidistant-ratio equivalent structure of music M , and elements a, b, c, d of M . Then $\overline{ab} \cong \overline{cd}$ if and only if $\overline{cd} \cong \overline{ab}$.

- (27) Let us consider an equidistant-ratio equivalent structure of music M , and elements a, b, c, d, e, f of M . Suppose $\overline{ab} \cong \overline{cd}$ and $\overline{cd} \cong \overline{ef}$. Then $\overline{ab} \cong \overline{ef}$.
- (28) Let us consider a satisfying interval, equidistant-ratio equivalent structure of music S , and elements a, b, c of S . Then $\overline{ab} \cong \overline{ac}$ if and only if $b = c$. The theorem is a consequence of (25).

From now on M denotes an equidistant-ratio equivalent structure of music and a, b, c, d, e, f denote elements of M .

Now we state the propositions:

- (29) $\overline{aa} \cong \overline{aa}$.
- (30) The equidistance of M is reflexive in $(\text{the carrier of } M) \times (\text{the carrier of } M)$. The theorem is a consequence of (25).
- (31) Suppose M is not empty. Then
- (i) the equidistance of M is reflexive, and
 - (ii) $\text{field}(\text{the equidistance of } M) = (\text{the carrier of } M) \times (\text{the carrier of } M)$.

The theorem is a consequence of (30).

- (32) The equidistance of M is symmetric in $(\text{the carrier of } M) \times (\text{the carrier of } M)$. The theorem is a consequence of (26).
- (33) The equidistance of M is transitive in $(\text{the carrier of } M) \times (\text{the carrier of } M)$. The theorem is a consequence of (27).
- (34) The equidistance of M is an equivalence relation of $(\text{the carrier of } M) \times (\text{the carrier of } M)$. The theorem is a consequence of (30), (32), and (33).
- (35) Let us consider a ratio symmetric, equidistant-ratio equivalent structure of music M , and elements a, b, c, d of M . Then $\overline{ab} \cong \overline{cd}$ if and only if $\overline{ba} \cong \overline{dc}$.
- (36) Let us consider a unison-ratio stable, equidistant-ratio equivalent structure of music S , and elements a, b, c of S . If $\overline{aa} \cong \overline{bc}$, then $b = c$.

Let S be a natural, satisfying interval, harmonic closed, equidistant-ratio equivalent structure of music, fr be an element of S , and n be a non zero natural number. The n -harmonic of fr in S yielding an element of S is defined by

(Def. 19) $\langle fr, it \rangle \in [(1, n)]_\alpha$, where α is the equidistance of S .

We say that S is harmonic linear if and only if

(Def. 20) for every element fr of S and for every non zero natural number n , there exists a positive real number f such that $fr = f$ and the n -harmonic of fr in $S = n \cdot f$.

Now we state the propositions:

(37) \mathbb{R} -music is harmonic linear. The theorem is a consequence of (1) and (24).

(38) \mathbb{Q} -music is harmonic linear. The theorem is a consequence of (2) and (24).

One can check that there exists a harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is harmonic linear.

One can check that the functor \mathbb{R} -music yields a harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Let us note that the functor \mathbb{Q} -music yields a harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be a harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that M is harmonic stable if and only if (Def. 21) $\overline{\text{for every elements } f_1, f_2 \text{ of } M \text{ and for every non zero natural numbers } n, m, \text{ the } n\text{-harmonic of } f_1 \text{ in } M \text{ the } m\text{-harmonic of } f_1 \text{ in } M \cong \text{the } n\text{-harmonic of } f_2 \text{ in } M \text{ the } m\text{-harmonic of } f_2 \text{ in } M.}$

Now we state the propositions:

(39) \mathbb{R} -music is harmonic stable. The theorem is a consequence of (1) and (13).

(40) \mathbb{Q} -music is harmonic stable. The theorem is a consequence of (2) and (22).

Observe that there exists a harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is harmonic stable.

One can verify that the functor \mathbb{R} -music yields a harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Observe that the functor \mathbb{Q} -music yields a harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be a harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music and fr be an element of M . The functors: the set of unison of fr in M , the set of octave of fr in M , the set of fifth of fr in M , the set of fourth of fr in M , and the set of major sixth of fr in M yielding subsets of $(\text{the carrier of } M) \times (\text{the carrier of } M)$ are defined by terms

(Def. 22) [\langle the 1-harmonic of fr in M , the 1-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 23) [\langle the 1-harmonic of fr in M , the 2-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 24) [\langle the 2-harmonic of fr in M , the 3-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 25) [\langle the 3-harmonic of fr in M , the 4-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 26) [\langle the 3-harmonic of fr in M , the 5-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

respectively. The functors: the set of major third of fr in M , the set of minor third of fr in M , the set of minor sixth of fr in M , the set of major tone of fr in M , and the set of minor tone of fr in M yielding subsets of (the carrier of M) \times (the carrier of M) are defined by terms

(Def. 27) [\langle the 4-harmonic of fr in M , the 5-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 28) [\langle the 5-harmonic of fr in M , the 6-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 29) [\langle the 5-harmonic of fr in M , the 8-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 30) [\langle the 8-harmonic of fr in M , the 9-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

(Def. 31) [\langle the 9-harmonic of fr in M , the 10-harmonic of fr in M \rangle_α , where α is the equidistance of M ,

respectively. The functors: the set of unison of M , the set of octave of M , the set of fifth of M , the set of fourth of M , and the set of major sixth of M yielding subsets of (the carrier of M) \times (the carrier of M) are defined by terms

(Def. 32) [$\langle 1, 1 \rangle_\alpha$, where α is the equidistance of M ,

(Def. 33) [$\langle 1, 2 \rangle_\alpha$, where α is the equidistance of M ,

(Def. 34) [$\langle 2, 3 \rangle_\alpha$, where α is the equidistance of M ,

(Def. 35) [$\langle 3, 4 \rangle_\alpha$, where α is the equidistance of M ,

(Def. 36) [$\langle 3, 5 \rangle_\alpha$, where α is the equidistance of M ,

respectively. The functors: the set of major third of M , the set of minor third of M , the set of minor sixth of M , the set of major tone of M , and the set of minor tone of M yielding subsets of (the carrier of M) \times (the carrier of M) are defined by terms

(Def. 37) [$\langle 4, 5 \rangle_\alpha$, where α is the equidistance of M ,

(Def. 38) $[\{5, 6\}]_\alpha$, where α is the equidistance of M ,

(Def. 39) $[\{5, 8\}]_\alpha$, where α is the equidistance of M ,

(Def. 40) $[\{8, 9\}]_\alpha$, where α is the equidistance of M ,

(Def. 41) $[\{9, 10\}]_\alpha$, where α is the equidistance of M ,

respectively. Let S be a harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that S is fifth constructible if and only if

(Def. 42) for every element fr of S , there exists an element q of S such that $\langle fr, q \rangle \in$ the set of fifth of S .

Now we state the propositions:

(41) Let us consider an element fr of \mathbb{R} -music. Then there exist positive real numbers f, q_1 such that

(i) $f = fr$, and

(ii) $q_1 = \frac{(3 \text{ qua real number})}{2} \cdot f$, and

(iii) $\langle f, q_1 \rangle \in$ the set of fifth of \mathbb{R} -music.

The theorem is a consequence of (1) and (24).

(42) \mathbb{R} -music is fifth constructible. The theorem is a consequence of (41) and (1).

(43) Let us consider an element fr of \mathbb{Q} -music. Then there exist positive rational numbers f, q_1 such that

(i) $f = fr$, and

(ii) $q_1 = \frac{(3 \text{ qua rational number})}{2} \cdot f$, and

(iii) $\langle f, q_1 \rangle \in$ the set of fifth of \mathbb{Q} -music.

The theorem is a consequence of (2) and (24).

(44) \mathbb{Q} -music is fifth constructible. The theorem is a consequence of (43) and (2).

Let us observe that there exists a harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is fifth constructible.

Let us note that the functor \mathbb{R} -music yields a fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Let us note that the functor \mathbb{Q} -music yields a fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be a fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music and fr be an element of M . The fifth of fr in M yielding an element of M is defined by

(Def. 43) $\langle fr, it \rangle \in$ the set of fifth of M .

Now we state the propositions:

(45) Let us consider a fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music M , and an element fr of M . Then the set of fifth of fr in $M =$ the set of fifth of M . The theorem is a consequence of (24) and (27).

(46) Let us consider an element fr of \mathbb{R} -music. Then there exists a positive real number f such that

(i) $fr = f$, and

(ii) the fifth of fr in \mathbb{R} -music $= \frac{(3 \text{ qua real number})}{2} \cdot f$.

The theorem is a consequence of (1) and (41).

(47) Let us consider an element fr of \mathbb{Q} -music. Then there exists a positive rational number f such that

(i) $fr = f$, and

(ii) the fifth of fr in \mathbb{Q} -music $= \frac{(3 \text{ qua rational number})}{2} \cdot f$.

The theorem is a consequence of (2) and (43).

Let M be a fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that M is classical fifth if and only if

(Def. 44) for every element fr of M , there exists a positive real number f such that $fr = f$ and the fifth of fr in $M = \frac{(3 \text{ qua real number})}{2} \cdot f$.

One can verify that there exists a fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is classical fifth.

One can verify that the functor \mathbb{R} -music yields a classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

One can check that the functor \mathbb{Q} -music yields a classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

5. HARMONIC

Now we state the propositions:

- (48) Let us consider a harmonic closed, natural, unison-ratio stable, satisfying interval, equidistant-ratio equivalent structure of music M , and an element fr of M . Then the 1-harmonic of fr in $M = fr$. The theorem is a consequence of (36).
- (49) Let us consider a harmonic stable, harmonic closed, natural, unison-ratio stable, satisfying interval, equidistant-ratio equivalent structure of music M , and elements a, b of M . Then $\overline{aa} \cong \overline{bb}$. The theorem is a consequence of (48).
- (50) Let us consider a harmonic stable, harmonic linear, harmonic closed, natural, unison-ratio stable, satisfying interval, equidistant-ratio equivalent structure of music M , and an element fr of M . Then the set of octave of fr in $M =$ the set of octave of M . The theorem is a consequence of (48), (27), and (24).
- (51) Let us consider a fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent, non empty structure of music M , and an element fr of M . Then there exists a sequence s_{11} of M such that
 - (i) $s_{11}(0) = fr$, and
 - (ii) for every natural number n , $\langle s_{11}(n), s_{11}(n + 1) \rangle \in$ the set of fifth of M .

PROOF: Define $\mathcal{P}[\text{set}, \text{set}, \text{set}] \equiv$ there exist positive real numbers x, y such that $\langle \mathcal{S}_2, \mathcal{S}_3 \rangle \in$ the set of fifth of M . For every natural number n and for every element x of M , there exists an element y of M such that $\mathcal{P}[n, x, y]$. Consider s_{11} being a sequence of M such that $s_{11}(0) = fr$ and for every natural number n , $\mathcal{P}[n, s_{11}(n), s_{11}(n + 1)]$. \square

Let M be a structure of music and a, b, c be elements of M . We say that b is between a and c if and only if

- (Def. 45) there exist positive real numbers r_1, r_2, r_3 such that $a = r_1$ and $b = r_2$ and $c = r_3$ and $r_1 \leq r_2 < r_3$.

Let S be a harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that S is octave constructible if and only if

- (Def. 46) for every element fr of S , there exists an element o of S such that $\langle fr, o \rangle \in$ the set of octave of S .

Now we state the propositions:

(52) Let us consider an element fr of \mathbb{R} -music. Then there exist positive real numbers f, q_1 such that

(i) $f = fr$, and

(ii) $q_1 = 2 \cdot f$, and

(iii) $\langle f, q_1 \rangle \in$ the set of octave of \mathbb{R} -music.

The theorem is a consequence of (1) and (24).

(53) \mathbb{R} -music is octave constructible. The theorem is a consequence of (52) and (1).

(54) Let us consider an element fr of \mathbb{Q} -music. Then there exist positive rational numbers f, q_1 such that

(i) $f = fr$, and

(ii) $q_1 = 2 \cdot f$, and

(iii) $\langle f, q_1 \rangle \in$ the set of octave of \mathbb{Q} -music.

The theorem is a consequence of (2) and (24).

(55) \mathbb{Q} -music is octave constructible. The theorem is a consequence of (54) and (2).

Let us note that there exists a classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is octave constructible.

Let us observe that the functor \mathbb{R} -music yields an octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Let us note that the functor \mathbb{Q} -music yields an octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be an octave constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music and fr be an element of M . The octave of fr in M yielding an element of M is defined by

(Def. 47) $\langle fr, it \rangle \in$ the set of octave of M .

Let M be a satisfying real, non empty structure of music and r be an element of M . The functor \mathbb{Q}_r yielding a positive real number is defined by the term

(Def. 48) r .

Let M be an octave constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that M is classical octave if and only if

(Def. 49) for every element fr of M , there exists a positive real number f such that $fr = f$ and the octave of fr in $M = 2 \cdot f$.

Now we state the propositions:

(56) \mathbb{R} -music is classical octave. The theorem is a consequence of (52) and (1).

(57) \mathbb{Q} -music is classical octave. The theorem is a consequence of (54) and (2).

One can verify that there exists an octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is classical octave.

Observe that the functor \mathbb{R} -music yields a classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Observe that the functor \mathbb{Q} -music yields a classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be an octave constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music. We say that M is octave descending constructible if and only if

(Def. 50) for every element fr of M , there exists an element o of M such that $\langle o, fr \rangle \in$ the set of octave of M .

Now we state the propositions:

(58) Let us consider an element fr of \mathbb{R} -music. Then there exist positive real numbers f, q_1 such that

(i) $f = fr$, and

(ii) $q_1 = \frac{(1 \text{ qua real number})}{2} \cdot f$, and

(iii) $\langle q_1, f \rangle \in$ the set of octave of \mathbb{R} -music.

The theorem is a consequence of (1), (24), and (35).

(59) \mathbb{R} -music is octave descending constructible. The theorem is a consequence of (58) and (1).

(60) Let us consider an element fr of \mathbb{Q} -music. Then there exist positive rational numbers f , q_1 such that

(i) $f = fr$, and

(ii) $q_1 = \frac{(1 \text{ qua rational number})}{2} \cdot f$, and

(iii) $\langle q_1, f \rangle \in$ the set of octave of \mathbb{Q} -music.

The theorem is a consequence of (2), (24), and (35).

(61) \mathbb{Q} -music is octave descending constructible. The theorem is a consequence of (60) and (2).

One can verify that there exists a classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music which is octave descending constructible.

One can verify that the functor \mathbb{R} -music yields an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Note that the functor \mathbb{Q} -music yields an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music.

Let M be an octave descending constructible, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent structure of music and fr be an element of M . The octave descending of fr in M yielding an element of M is defined by

(Def. 51) $\langle it, fr \rangle \in$ the set of octave of M .

Now we state the propositions:

(62) Let us consider an octave descending constructible, classical octave, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music M , and an element fr of M . Then there exists a positive real number r such that

(i) $fr = r$, and

(ii) the octave descending of fr in $M = \frac{r}{2}$.

The theorem is a consequence of (1).

(63) Let us consider classical octave, octave constructible, classical fifth, fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structures of music M_1, M_2 , an element f_1 of M_1 , and an element f_2 of M_2 . Suppose $f_1 = f_2$. Then

- (i) the fifth of f_1 in $M_1 =$ the fifth of f_2 in M_2 , and
- (ii) the octave of f_1 in $M_1 =$ the octave of f_2 in M_2 .

(64) Let us consider octave descending constructible, classical octave, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structures of music M_1, M_2 , an element fr_1 of M_1 , and an element fr_2 of M_2 . Suppose $fr_1 = fr_2$. Then the octave descending of fr_1 in $M_1 =$ the octave descending of fr_2 in M_2 . The theorem is a consequence of (62).

Let M be an octave descending constructible, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent structure of music and f_{10}, fr be elements of M . The reduct fifth of the fr with fundamental frequency f_{10} in M yielding an element of M is defined by the term

$$(\text{Def. 52}) \quad \left\{ \begin{array}{l} \text{the fifth of } fr \text{ in } M, \text{ if the fifth of } fr \text{ in } M \text{ is between } f_{10} \text{ and the} \\ \text{octave of } f_{10} \text{ in } M, \\ \text{the octave descending of (the fifth of } fr \text{ in } M) \text{ in } M, \text{ otherwise.} \end{array} \right.$$

Now we state the propositions:

(65) Let us consider octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structures of music M_1, M_2 , elements fr_1, f_{11} of M_1 , and elements fr_2, f_{12} of M_2 . Suppose $fr_1 = fr_2$ and $f_{11} = f_{12}$. Then the reduct fifth of the fr_1 with fundamental frequency f_{11} in $M_1 =$ the reduct fifth of the fr_2 with fundamental frequency f_{12} in M_2 . The theorem is a consequence of (63) and (64).

(66) Let us consider a classical fifth, fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music M , and an element fr of M . Then there exist positive real numbers r, s such that

- (i) $r = fr$, and
- (ii) $s = \frac{(3 \text{ qua real number})}{2} \cdot r$, and
- (iii) the fifth of fr in $M = s$.

(67) Let us consider an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic closed, natural, satisfying interval, equidistant-ratio equivalent structure of music M , and elements f_{10} , fr of M . Suppose fr is between f_{10} and the octave of f_{10} in M . Then there exist positive real numbers r_1, r_2, r_3 such that

(i) $f_{10} = r_1$, and

(ii) $fr = r_2$, and

(iii) the octave of f_{10} in $M = 2 \cdot r_1$, and

(iv) $r_1 \leq r_2 \leq 2 \cdot r_1$.

(68) Let us consider an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music M , and elements f_{10} , fr of M . Suppose fr is between f_{10} and the octave of f_{10} in M . Then the reduct fifth of the fr with fundamental frequency f_{10} in M is between f_{10} and the octave of f_{10} in M . The theorem is a consequence of (67) and (62).

A space of music is an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic stable, harmonic linear, harmonic closed, natural, ratio symmetric, unison-ratio stable, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music. Now we state the propositions:

(69) \mathbb{R} -music is a space of music.

(70) \mathbb{Q} -music is a space of music.

6. SPIRAL OF FIFTHS

Now we state the proposition:

(71) Let us consider an octave descending constructible, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, non empty structure of music M , and elements f_{10} , fr of M . Then there exists a sequence s_{11} of M such that

(i) $s_{11}(0) = fr$, and

(ii) for every natural number n , $s_{11}(n + 1) =$ the reduct fifth of the $s_{11}(n)$ with fundamental frequency f_{10} in M .

PROOF: Define $\mathcal{P}[\text{set, set, set}] \equiv$ there exist elements x, y of M such that $x = \mathbb{S}_2$ and $y = \mathbb{S}_3$ and y is the reduct fifth of the x with fundamental frequency f_{10} in M . For every natural number n and for every element x of M , there exists an element y of M such that $\mathcal{P}[n, x, y]$. Consider s_{11} being a sequence of M such that $s_{11}(0) = fr$ and for every natural number n , $\mathcal{P}[n, s_{11}(n), s_{11}(n + 1)]$. \square

Let M be an octave descending constructible, octave constructible, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, non empty structure of music and f_{10}, fr be elements of M . The spiral of fifths of fr with fundamental frequency f_{10} in M yielding a sequence of M is defined by

(Def. 53) $it(0) = fr$ and for every natural number n , $it(n + 1) =$ the reduct fifth of the $it(n)$ with fundamental frequency f_{10} in M .

From now on M denotes an octave descending constructible, classical octave, octave constructible, classical fifth, fifth constructible, harmonic closed, natural, ratio symmetric, satisfying interval, equidistant-ratio equivalent, satisfying real, non empty structure of music and f_{10}, fr denote elements of M .

Now we state the propositions:

(72) Suppose fr is between f_{10} and the octave of f_{10} in M . Let us consider a natural number n . Then (the spiral of fifths of fr with fundamental frequency f_{10} in M)(n) is between f_{10} and the octave of f_{10} in M .

PROOF: Define $\mathcal{P}[\text{natural number}] \equiv$ (the spiral of fifths of fr with fundamental frequency f_{10} in M)(\mathbb{S}_1) is between f_{10} and the octave of f_{10} in M . For every natural number k such that $\mathcal{P}[k]$ holds $\mathcal{P}[k + 1]$. For every natural number k , $\mathcal{P}[k]$ from [2, Sch. 2]. \square

(73) (The spiral of fifths of f_{10} with fundamental frequency f_{10} in M)(1) = $\frac{(3 \text{ qua real number})}{2} \cdot (@f_{10})$. The theorem is a consequence of (66).

(74) (The spiral of fifths of f_{10} with fundamental frequency f_{10} in M)(2) = $\frac{(9 \text{ qua real number})}{8} \cdot (@f_{10})$. The theorem is a consequence of (73), (66), and (62).

(75) (The spiral of fifths of f_{10} with fundamental frequency f_{10} in M)(3) = $\frac{(27 \text{ qua real number})}{16} \cdot (@f_{10})$. The theorem is a consequence of (74) and (66).

(76) (The spiral of fifths of f_{10} with fundamental frequency f_{10} in M)(4) = $\frac{(81 \text{ qua real number})}{64} \cdot (@f_{10})$. The theorem is a consequence of (75), (66), and (62).

(77) (The spiral of fifths of f_{10} with fundamental frequency f_{10} in M)(5) = $\frac{(243 \text{ qua real number})}{128} \cdot (@f_{10})$. The theorem is a consequence of (76) and (66).

(78) $\frac{@(the \text{ spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(2)}{@fr} =$

- $\frac{3 \cdot 3 \text{ qua real number}}{2 \cdot 2 \cdot 2}$. The theorem is a consequence of (74).
- (79) $\frac{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(4)}}{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(2)}} = \frac{3 \cdot 3 \text{ qua real number}}{2 \cdot 2 \cdot 2}$. The theorem is a consequence of (74) and (76).
- (80) $\frac{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(1)}}{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(4)}} = \frac{32 \text{ qua real number}}{27}$. The theorem is a consequence of (73) and (76).
- (81) $\frac{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(3)}}{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(1)}} = \frac{9 \text{ qua real number}}{8}$. The theorem is a consequence of (73) and (75).
- (82) $\frac{\textcircled{\text{(the octave of } fr \text{ in } M)}}{\textcircled{\text{(the spiral of fifths of } fr \text{ with fundamental frequency } fr \text{ in } M)(3)}} = \frac{32 \text{ qua real number}}{27}$. The theorem is a consequence of (75).

Let M be a space of music and s_{10} be an element of $(\text{the carrier of } M)^2$. We say that s_{10} is monotonic if and only if

- (Def. 54) there exists an element fr of M and there exist positive real numbers r_1, r_2 such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $r_1 < r_2$ and $s_{10}(2) = \text{the octave of } fr \text{ in } M$.

Let s_{10} be an element of $(\text{the carrier of } M)^3$. We say that s_{10} is ditonic if and only if

- (Def. 55) there exists an element fr of M and there exist positive real numbers r_1, r_2, r_3 such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $r_1 < r_2 < r_3$ and $s_{10}(3) = \text{the octave of } fr \text{ in } M$.

Let s_{10} be an element of $(\text{the carrier of } M)^4$. We say that s_{10} is tritonic if and only if

- (Def. 56) there exists an element fr of M and there exist positive real numbers r_1, r_2, r_3, r_4 such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $r_1 < r_2 < r_3$ and $r_3 < r_4$ and $s_{10}(4) = \text{the octave of } fr \text{ in } M$.

Let s_{10} be an element of $(\text{the carrier of } M)^5$. We say that s_{10} is tetratonic if and only if

- (Def. 57) there exists an element fr of M and there exist positive real numbers r_1, r_2, r_3, r_4, r_5 such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $s_{10}(5) = r_5$ and $r_1 < r_2 < r_3$ and $r_3 < r_4 < r_5$ and $s_{10}(5) = \text{the octave of } fr \text{ in } M$.

Let n be a natural number and s_{10} be an element of $(\text{the carrier of } M)^n$. We say that s_{10} is pentatonic if and only if

- (Def. 58) $n = 6$ and there exists an element fr of M and there exist positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6$ such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $s_{10}(5) = r_5$ and $s_{10}(6) =$

r_6 and $r_1 < r_2 < r_3$ and $r_3 < r_4 < r_5$ and $r_5 < r_6$ and $s_{10}(6) =$ the octave of fr in M .

Let s_{10} be an element of (the carrier of M)⁷. We say that s_{10} is hexatonic if and only if

(Def. 59) there exists an element fr of M and there exist positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6, r_7$ such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $s_{10}(5) = r_5$ and $s_{10}(6) = r_6$ and $s_{10}(7) = r_7$ and $r_1 < r_2 < r_3$ and $r_3 < r_4 < r_5$ and $r_5 < r_6 < r_7$ and $s_{10}(7) =$ the octave of fr in M .

Let n be a natural number and s_{10} be an element of (the carrier of M)ⁿ. We say that s_{10} is heptatonic if and only if

(Def. 60) $n = 8$ and there exists an element fr of M and there exist positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6, r_7, r_8$ such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $s_{10}(5) = r_5$ and $s_{10}(6) = r_6$ and $s_{10}(7) = r_7$ and $s_{10}(8) = r_8$ and $r_1 < r_2 < r_3$ and $r_3 < r_4 < r_5$ and $r_5 < r_6 < r_7$ and $r_7 < r_8$ and $s_{10}(8) =$ the octave of fr in M .

Let s_{10} be an element of (the carrier of M)⁹. We say that s_{10} is octatonic if and only if

(Def. 61) there exists an element fr of M and there exist positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6, r_7, r_8, r_9$ such that $s_{10}(1) = fr$ and $s_{10}(1) = r_1$ and $s_{10}(2) = r_2$ and $s_{10}(3) = r_3$ and $s_{10}(4) = r_4$ and $s_{10}(5) = r_5$ and $s_{10}(6) = r_6$ and $s_{10}(7) = r_7$ and $s_{10}(8) = r_8$ and $s_{10}(9) = r_9$ and $r_1 < r_2 < r_3$ and $r_3 < r_4 < r_5$ and $r_5 < r_6 < r_7$ and $r_7 < r_8 < r_9$ and $s_{10}(9) =$ the octave of fr in M .

7. PENTATONIC PYTHAGOREAN SCALE

Let M be a space of music and fr be an element of M . The pentatonic pythagorean scale of fr in M yielding an element of (the carrier of M)⁶ is defined by

(Def. 62) $it(1) = fr$ and $it(2) =$ (the spiral of fifths of fr with fundamental frequency fr in M)(2) and $it(3) =$ (the spiral of fifths of fr with fundamental frequency fr in M)(4) and $it(4) =$ (the spiral of fifths of fr with fundamental frequency fr in M)(1) and $it(5) =$ (the spiral of fifths of fr with fundamental frequency fr in M)(3) and $it(6) =$ the octave of fr in M .

From now on M denotes a space of music and f_{10}, fr, f_1, f_2 denote elements of M .

Now we state the proposition:

(83) The pentatonic pythagorean scale of fr in M is pentatonic. The theorem is a consequence of (74), (76), (73), and (75).

Let M be a space of music and f_1, f_2 be elements of M . The interval between f_1 and f_2 yielding a positive real number is defined by

(Def. 63) there exist positive real numbers r_1, r_2 such that $r_1 = f_1$ and $r_2 = f_2$ and $it = \frac{r_2}{r_1}$.

The pythagorean tone yielding a positive real number is defined by the term

(Def. 64) $\frac{(9 \text{ qua real number})}{8}$.

The pythagorean semiditone yielding a positive real number is defined by the term

(Def. 65) $\frac{(32 \text{ qua real number})}{27}$.

The pythagorean major third yielding a positive real number is defined by the term

(Def. 66) (the pythagorean tone) · (the pythagorean tone).

The pythagorean pure major third yielding a positive real number is defined by the term

(Def. 67) $\frac{(5 \text{ qua real number})}{4}$.

The syntonic comma yielding a positive real number is defined by the term

(Def. 68) $\frac{\text{the pythagorean major third}}{\text{the pythagorean pure major third}}$.

Now we state the propositions:

(84) The syntonic comma = $\frac{(81 \text{ qua real number})}{80}$.

(85) The pythagorean tone < the pythagorean semiditone.

(86) (The pythagorean tone) · (the pythagorean tone) · (the pythagorean semiditone) · (the pythagorean tone) · (the pythagorean semiditone) = 2.

Let M be a space of music and fr be an element of M . The functors: the first degree of pentatonic scale of fr in M , the second degree of pentatonic scale of fr in M , the third degree of pentatonic scale of fr in M , the fourth degree of pentatonic scale of fr in M , and the fifth degree of pentatonic scale of fr in M yielding elements of M are defined by terms

(Def. 69) (the pentatonic pythagorean scale of fr in M)(1),

(Def. 70) (the pentatonic pythagorean scale of fr in M)(2),

(Def. 71) (the pentatonic pythagorean scale of fr in M)(3),

(Def. 72) (the pentatonic pythagorean scale of fr in M)(4),

(Def. 73) (the pentatonic pythagorean scale of fr in M)(5),

respectively. The octave of pentatonic scale of fr in M yielding an element of M is defined by the term

(Def. 74) the octave of fr in M .

Now we state the propositions:

- (87) There exist elements r_1, r_2 of \mathbb{R}_+ such that the interval between f_1 and $f_2 = \mathbb{R}\text{-ratio}(r_1, r_2)$.
- (88) Let us consider positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6$. Suppose (the pentatonic pythagorean scale of fr in M)(1) = r_1 and (the pentatonic pythagorean scale of fr in M)(2) = r_2 and (the pentatonic pythagorean scale of fr in M)(3) = r_3 and (the pentatonic pythagorean scale of fr in M)(4) = r_4 and (the pentatonic pythagorean scale of fr in M)(5) = r_5 and (the pentatonic pythagorean scale of fr in M)(6) = r_6 . Then

- (i) $\frac{r_2}{r_1} = \frac{(9 \text{ qua real number})}{8}$, and
- (ii) $\frac{r_3}{r_2} = \frac{(9 \text{ qua real number})}{8}$, and
- (iii) $\frac{r_4}{r_3} = \frac{(32 \text{ qua real number})}{27}$, and
- (iv) $\frac{r_5}{r_4} = \frac{(9 \text{ qua real number})}{8}$, and
- (v) $\frac{r_6}{r_5} = \frac{(32 \text{ qua real number})}{27}$.

The theorem is a consequence of (83), (78), (79), (80), (81), and (82).

- (89) There exist positive real numbers $r_1, r_2, r_3, r_4, r_5, r_6$ such that

- (i) (the pentatonic pythagorean scale of fr in M)(1) = r_1 , and
- (ii) (the pentatonic pythagorean scale of fr in M)(2) = r_2 , and
- (iii) (the pentatonic pythagorean scale of fr in M)(3) = r_3 , and
- (iv) (the pentatonic pythagorean scale of fr in M)(4) = r_4 , and
- (v) (the pentatonic pythagorean scale of fr in M)(5) = r_5 , and
- (vi) (the pentatonic pythagorean scale of fr in M)(6) = r_6 , and
- (vii) $\frac{r_2}{r_1} = \frac{(9 \text{ qua real number})}{8}$, and
- (viii) $\frac{r_3}{r_2} = \frac{(9 \text{ qua real number})}{8}$, and
- (ix) $\frac{r_4}{r_3} = \frac{(32 \text{ qua real number})}{27}$, and
- (x) $\frac{r_5}{r_4} = \frac{(9 \text{ qua real number})}{8}$, and
- (xi) $\frac{r_6}{r_5} = \frac{(32 \text{ qua real number})}{27}$.

The theorem is a consequence of (1) and (88).

- (90) $\frac{(9 \text{ qua real number})}{8} = \frac{(9 \text{ qua rational number})}{8}$.

- (91) (i) the interval between the first degree of pentatonic scale of fr in M and (the second degree of pentatonic scale of fr in M) = the pythagorean tone, and

- (ii) the interval between the second degree of pentatonic scale of fr in M and (the third degree of pentatonic scale of fr in M) = the pythagorean tone, and
- (iii) the interval between the third degree of pentatonic scale of fr in M and (the fourth degree of pentatonic scale of fr in M) = the pythagorean semiditone, and
- (iv) the interval between the fourth degree of pentatonic scale of fr in M and (the fifth degree of pentatonic scale of fr in M) = the pythagorean tone, and
- (v) the interval between the fifth degree of pentatonic scale of fr in M and (the octave of pentatonic scale of fr in M) = the pythagorean semiditone.

The theorem is a consequence of (89).

- (92) the fifth of fr in M is between fr and the octave of fr in M .

Let us consider positive real numbers r_1, r_2 . Now we state the propositions:

- (93) Suppose $f_1 = r_1$ and $f_2 = r_2$ and $r_2 = \frac{(4 \text{ qua real number})}{3} \cdot r_1$. Then

- (i) the fifth of f_2 in $M = 2 \cdot r_1$, and
- (ii) the fifth of f_2 in M is not between f_1 and the octave of f_1 in M .

- (94) Suppose $f_1 = r_1$ and $f_2 = r_2$ and $r_2 = \frac{(4 \text{ qua real number})}{3} \cdot r_1$. Then

- (i) if the fifth of f_2 in M is between f_{10} and the octave of f_{10} in M , then the octave descending of (the reduct fifth of the f_2 with fundamental frequency f_{10} in M) in $M = f_1$, and
- (ii) if the fifth of f_2 in M is not between f_{10} and the octave of f_{10} in M , then the reduct fifth of the f_2 with fundamental frequency f_{10} in $M = f_1$.

The theorem is a consequence of (62).

- (95) Suppose $f_1 = r_1$ and $f_2 = r_2$ and $r_2 = \frac{(4 \text{ qua real number})}{3} \cdot r_1$. Then the reduct fifth of the f_2 with fundamental frequency f_1 in $M = f_1$. The theorem is a consequence of (94) and (93).

8. HEPTATONIC PYTHAGOREAN SCALE

Let S be a space of music. We say that S is fourth constructible if and only if

- (Def. 75) for every element fr of S , there exists an element q of S such that $\langle fr, q \rangle \in$ the set of fourth of S .

Now we state the propositions:

(96) Let us consider a space of music M . Suppose $M = \mathbb{R}$ -music. Let us consider an element fr of M . Then there exist positive real numbers f, q_1 such that

- (i) $f = fr$, and
- (ii) $q_1 = \frac{(4 \text{ qua real number})}{3} \cdot f$, and
- (iii) $\langle f, q_1 \rangle \in$ the set of fourth of M .

The theorem is a consequence of (1) and (24).

(97) \mathbb{R} -music is fourth constructible. The theorem is a consequence of (96) and (1).

One can verify that there exists a space of music which is fourth constructible.

Let M be a fourth constructible space of music and fr be an element of M . The fourth of fr in M yielding an element of M is defined by

(Def. 76) $\langle fr, it \rangle \in$ the set of fourth of M .

We say that M is classical fourth if and only if

(Def. 77) for every element fr of M , there exists a positive real number f such that $fr = f$ and the fourth of fr in $M = \frac{(4 \text{ qua real number})}{3} \cdot f$.

Now we state the proposition:

(98) Let us consider a fourth constructible space of music M . Suppose $M = \mathbb{R}$ -music. Let us consider an element fr of M . Then there exists a positive real number f such that

- (i) $fr = f$, and
- (ii) the fourth of fr in $M = \frac{(4 \text{ qua real number})}{3} \cdot f$.

The theorem is a consequence of (1) and (96).

Let us note that there exists a fourth constructible space of music which is classical fourth.

Let M be a satisfying real, non empty structure of music. We say that M is euclidean if and only if

(Def. 78) for every elements f_1, f_2 of M , (the Ratio of M)(f_1, f_2) = $\frac{f_2}{f_1}$.

One can verify that there exists a satisfying real, non empty structure of music which is euclidean and every satisfying real, non empty structure of music which is euclidean is also satisfying interval and every satisfying real, non empty structure of music which is euclidean is also unison-ratio stable and every satisfying real, non empty structure of music which is euclidean is also ratio symmetric and there exists a classical fourth, fourth constructible space of music which is euclidean.

A heptatonic pythagorean score is a classical fourth, fourth constructible space of music. From now on H denotes a heptatonic pythagorean score and fr denotes an element of H .

Let H be a heptatonic pythagorean score and fr be an element of H . The heptatonic pythagorean scale of fr in H yielding an element of (the carrier of H)⁸ is defined by

(Def. 79) $it(1)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(1) and $it(2)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(3) and $it(3)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(5) and $it(4)$ = the fourth of fr in H and $it(5)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(2) and $it(6)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(4) and $it(7)$ = (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(6) and $it(8)$ = the octave of (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(1) in H .

Now we state the propositions:

- (99) the fourth of fr in H is between fr and the octave of fr in H .
- (100) Let us consider a natural number n . Then (the spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(n) is between fr and the octave of fr in H .
- (101) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(1) = fr . The theorem is a consequence of (66) and (62).
- (102) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(2) = $\frac{(3 \text{ qua real number})}{2} \cdot (@fr)$. The theorem is a consequence of (101) and (66).
- (103) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(3) = $\frac{(9 \text{ qua real number})}{8} \cdot (@fr)$. The theorem is a consequence of (102), (66), and (62).
- (104) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(4) = $\frac{(27 \text{ qua real number})}{16} \cdot (@fr)$. The theorem is a consequence of (103) and (66).
- (105) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(5) = $\frac{(81 \text{ qua real number})}{64} \cdot (@fr)$. The theorem is a consequence of (104), (66), and (62).
- (106) (The spiral of fifths of (the fourth of fr in H) with fundamental frequency fr in H)(6) = $\frac{(243 \text{ qua real number})}{128} \cdot (@fr)$. The theorem is a consequence of

(105) and (66).

- (107) (i) (the heptatonic pythagorean scale of fr in H)(1) = $1 \cdot (^@fr)$, and
 (ii) (the heptatonic pythagorean scale of fr in H)(2) = $\frac{(9 \text{ qua real number})}{8} \cdot (^@fr)$, and
 (iii) (the heptatonic pythagorean scale of fr in H)(3) = $\frac{(81 \text{ qua real number})}{64} \cdot (^@fr)$, and
 (iv) (the heptatonic pythagorean scale of fr in H)(4) = $\frac{(4 \text{ qua real number})}{3} \cdot (^@fr)$, and
 (v) (the heptatonic pythagorean scale of fr in H)(5) = $\frac{(3 \text{ qua real number})}{2} \cdot (^@fr)$, and
 (vi) (the heptatonic pythagorean scale of fr in H)(6) = $\frac{(27 \text{ qua real number})}{16} \cdot (^@fr)$, and
 (vii) (the heptatonic pythagorean scale of fr in H)(7) = $\frac{(243 \text{ qua real number})}{128} \cdot (^@fr)$, and

(viii) (the heptatonic pythagorean scale of fr in H)(8) = $2 \cdot (^@fr)$.

The theorem is a consequence of (101), (103), (105), (102), (104), and (106).

- (108) The heptatonic pythagorean scale of fr in H is heptatonic. The theorem is a consequence of (107).

The pythagorean semitone yielding a positive real number is defined by the term

(Def. 80) $\frac{(256 \text{ qua real number})}{243}$.

Now we state the propositions:

(109) $\frac{\text{the pythagorean tone}}{2} < \text{the pythagorean semitone}$.

(110) $(\text{The pythagorean tone}) \cdot (\text{the pythagorean tone}) \cdot (\text{the pythagorean semitone}) \cdot (\text{the pythagorean tone}) \cdot (\text{the pythagorean tone}) \cdot (\text{the pythagorean tone}) \cdot (\text{the pythagorean semitone}) = 2$.

Let H be a heptatonic pythagorean score and fr be an element of H . The functors: the first degree of heptatonic scale of fr in H , the second degree of heptatonic scale of fr in H , the third degree of heptatonic scale of fr in H , the fourth degree of heptatonic scale of fr in H , and the fifth degree of heptatonic scale of fr in H yielding elements of H are defined by terms

(Def. 81) (the heptatonic pythagorean scale of fr in H)(1),

(Def. 82) (the heptatonic pythagorean scale of fr in H)(2),

(Def. 83) (the heptatonic pythagorean scale of fr in H)(3),

(Def. 84) (the heptatonic pythagorean scale of fr in H)(4),

(Def. 85) (the heptatonic pythagorean scale of fr in H)(5),

respectively. The functors: the sixth degree of heptatonic scale of fr in H , the seventh degree of heptatonic scale of fr in H , and the eight degree of heptatonic scale of fr in H yielding elements of H are defined by terms

(Def. 86) (the heptatonic pythagorean scale of fr in H)(6),

(Def. 87) (the heptatonic pythagorean scale of fr in H)(7),

(Def. 88) the octave of fr in H ,

respectively. Now we state the proposition:

(111) (i) the interval between the first degree of heptatonic scale of fr in H and (the second degree of heptatonic scale of fr in H) = the pythagorean tone, and

(ii) the interval between the second degree of heptatonic scale of fr in H and (the third degree of heptatonic scale of fr in H) = the pythagorean tone, and

(iii) the interval between the third degree of heptatonic scale of fr in H and (the fourth degree of heptatonic scale of fr in H) = the pythagorean semitone, and

(iv) the interval between the fourth degree of heptatonic scale of fr in H and (the fifth degree of heptatonic scale of fr in H) = the pythagorean tone, and

(v) the interval between the fifth degree of heptatonic scale of fr in H and (the sixth degree of heptatonic scale of fr in H) = the pythagorean tone, and

(vi) the interval between the sixth degree of heptatonic scale of fr in H and (the seventh degree of heptatonic scale of fr in H) = the pythagorean tone, and

(vii) the interval between the seventh degree of heptatonic scale of fr in H and (the eight degree of heptatonic scale of fr in H) = the pythagorean semitone.

The theorem is a consequence of (107).

From now on H denotes a heptatonic pythagorean score and fr denotes an element of H .

Let M be a space of music, n be a natural number, and s_{10} be an element of (the carrier of M) ^{n} . Assume s_{10} is heptatonic. We say that s_{10} is perfect fifth if and only if

(Def. 89) $\langle s_{10}(1), s_{10}(5) \rangle, \langle s_{10}(2), s_{10}(6) \rangle, \langle s_{10}(3), s_{10}(7) \rangle, \langle s_{10}(4), s_{10}(8) \rangle \in$ the set of fifth of M .

Now we state the proposition:

(112) Let us consider an euclidean heptatonic pythagorean score H , and an element fr of H . Then the heptatonic pythagorean scale of fr in H is perfect fifth. The theorem is a consequence of (108), (107), and (24).

Let H be a heptatonic pythagorean score and fr be an element of H . The heptatonic pythagorean scale ascending of fr in H yielding an element of (the carrier of H)⁸ is defined by the term

(Def. 90) the heptatonic pythagorean scale of (the octave of fr in H) in H .

Now we state the propositions:

(113) (i) (the heptatonic pythagorean scale ascending of fr in H)(1) = $2 \cdot (\textcircled{fr})$, and

(ii) (the heptatonic pythagorean scale ascending of fr in H)(2) = $\frac{9 \text{ qua real number}}{4} \cdot (\textcircled{fr})$, and

(iii) (the heptatonic pythagorean scale ascending of fr in H)(3) = $\frac{81 \text{ qua real number}}{32} \cdot (\textcircled{fr})$, and

(iv) (the heptatonic pythagorean scale ascending of fr in H)(4) = $\frac{8 \text{ qua real number}}{3} \cdot (\textcircled{fr})$, and

(v) (the heptatonic pythagorean scale ascending of fr in H)(5) = $(3 \text{ qua real number}) \cdot (\textcircled{fr})$, and

(vi) (the heptatonic pythagorean scale ascending of fr in H)(6) = $\frac{27 \text{ qua real number}}{8} \cdot (\textcircled{fr})$, and

(vii) (the heptatonic pythagorean scale ascending of fr in H)(7) = $\frac{243 \text{ qua real number}}{64} \cdot (\textcircled{fr})$, and

(viii) (the heptatonic pythagorean scale ascending of fr in H)(8) = $4 \cdot (\textcircled{fr})$.
The theorem is a consequence of (107).

(114) (The heptatonic pythagorean scale of fr in H)(8) = (the heptatonic pythagorean scale ascending of fr in H)(1). The theorem is a consequence of (107) and (113).

(115) (i) the interval between the fifth degree of heptatonic scale of fr in H and (the second degree of heptatonic scale of (the octave of fr in H) in H) = $\frac{(3 \text{ qua real number})}{2}$, and

(ii) the interval between the sixth degree of heptatonic scale of fr in H and (the third degree of heptatonic scale of (the octave of fr in H) in H) = $\frac{(3 \text{ qua real number})}{2}$, and

(iii) the interval between the seventh degree of heptatonic scale of fr in H and (the fourth degree of heptatonic scale of (the octave of fr in H) in H) $\neq \frac{(3 \text{ qua real number})}{2}$, and

- (iv) the interval between the eight degree of heptatonic scale of fr in H and (the fifth degree of heptatonic scale of (the octave of fr in H) in H) = $\frac{(3 \text{ qua real number})}{2}$.

The theorem is a consequence of (107) and (113).

- (116) Let us consider an euclidean heptatonic pythagorean score H , and elements f_1, f_2 of H . Then the interval between f_1 and $f_2 =$ (the Ratio of H)(f_1, f_2).

- (117) Let us consider an euclidean heptatonic pythagorean score H , and an element fr of H . Then

- (i) \langle (the heptatonic pythagorean scale of fr in H)(5), (the heptatonic pythagorean scale ascending of fr in H)(2) \rangle, \langle (the heptatonic pythagorean scale of fr in H)(6), (the heptatonic pythagorean scale ascending of fr in H)(3) $\rangle \in$ the set of fifth of H , and
- (ii) \langle (the heptatonic pythagorean scale of fr in H)(7), (the heptatonic pythagorean scale ascending of fr in H)(4) $\rangle \notin$ the set of fifth of H .

The theorem is a consequence of (115), (24), and (116).

Let H be a space of music, n be a non zero, natural number, s_{10} be an element of (the carrier of H) n , and i be a natural number. The functor $\#_i^{s_{10}}$ yielding an element of H is defined by the term

(Def. 91)
$$\begin{cases} s_{10}(i), & \text{if } i \in \text{Seg } n, \\ \text{the element of } H, & \text{otherwise.} \end{cases}$$

Assume s_{10} is heptatonic. We say that s_{10} is dorian if and only if

- (Def. 92) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_2$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_2$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is hypodorian if and only if

- (Def. 93) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_2$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_2$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is phrygian if and only if

- (Def. 94) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval

between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_2$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_2$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is hypophrygian if and only if

- (Def. 95) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_2$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_2$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is lydian if and only if

- (Def. 96) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_2$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_2$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is hypolydian if and only if

- (Def. 97) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_2$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_2$.

Assume s_{10} is heptatonic. We say that s_{10} is mixolydian if and only if

- (Def. 98) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_2$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_2$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is hypomixolydian if and only if

- (Def. 99) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_2$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_2$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is eolian if and only if

- (Def. 100) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_2$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_2$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is hypoeolian if and only if

- (Def. 101) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_2$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_1$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_2$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Assume s_{10} is heptatonic. We say that s_{10} is ionan if and only if

- (Def. 102) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_2$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_1$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_2$.

Assume s_{10} is heptatonic. We say that s_{10} is hypoionan if and only if

- (Def. 103) there exist positive real numbers t_1, t_2 such that $t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_1 \cdot t_2 \cdot t_2 = 2$ and the interval between $\#_1^{s_{10}}$ and $\#_2^{s_{10}} = t_1$ and the interval between $\#_2^{s_{10}}$ and $\#_3^{s_{10}} = t_1$ and the interval between $\#_3^{s_{10}}$ and $\#_4^{s_{10}} = t_2$ and the interval between $\#_4^{s_{10}}$ and $\#_5^{s_{10}} = t_1$ and the interval between $\#_5^{s_{10}}$ and $\#_6^{s_{10}} = t_1$ and the interval between $\#_6^{s_{10}}$ and $\#_7^{s_{10}} = t_2$ and the interval between $\#_7^{s_{10}}$ and $\#_8^{s_{10}} = t_1$.

Now we state the proposition:

- (118) The heptatonic pythagorean scale of fr in H is ionan. The theorem is a consequence of (108), (107), and (111).

ACKNOWLEDGEMENT: I would like to thank H el ene Cambier (Professor of music history at the Music Academy La Louvi ere) and Marie Barbier (musical instrument: Recorder) for valuable suggestions.

REFERENCES

- [1] Grzegorz Bancerek. On the structure of Mizar types. In Herman Geuvers and Fairouz Kamareddine, editors, *Electronic Notes in Theoretical Computer Science*, volume 85, pages 69–85. Elsevier, 2003.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8_17.
- [4] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.
- [5] François Baskevitch. *Les représentations de la propagation du son, d’Aristote à l’Encyclopédie*. PhD thesis, Université de Nantes, France, 2008.
- [6] Adam Grabowski, Artur Korniłowicz, and Christoph Schwarzweller. On algebraic hierarchies in mathematical repository of Mizar. In M. Ganzha, L. Maciaszek, and M. Paprzycki, editors, *Proceedings of the 2016 Federated Conference on Computer Science and Information Systems (FedCSIS)*, volume 8 of *Annals of Computer Science and Information Systems*, pages 363–371, 2016. doi:10.15439/2016F520.
- [7] Bernard Parzysz and Yves Hellegouarch. *Musique et mathématique: (suivi de) Gammes naturelles*. Number 53. Association des Professeurs de Mathématiques de l’Enseignement Public (APMEP), Paris, 1984.

Accepted September 29, 2018

Fundamental Properties of Fuzzy Implications

Adam Grabowski 
Institute of Informatics
University of Białystok
Poland

Summary. In the article we continue in the Mizar system [8], [2] the formalization of fuzzy implications according to the monograph of Baczyński and Jayaram “*Fuzzy Implications*” [1]. We develop a framework of Mizar attributes allowing us for a smooth proving of basic properties of these fuzzy connectives [9]. We also give a set of theorems about the ordering of nine fundamental implications: Łukasiewicz (I_{LK}), Gödel (I_{GD}), Reichenbach (I_{RC}), Kleene-Dienes (I_{KD}), Goguen (I_{GG}), Rescher (I_{RS}), Yager (I_{YG}), Weber (I_{WB}), and Fodor (I_{FD}).

This work is a continuation of the development of fuzzy sets in Mizar [6]; it could be used to give a variety of more general operations on fuzzy sets [13]. The formalization follows [10], [5], and [4].

MSC: 03B52 68T37 03B35

Keywords: fuzzy implication; fuzzy set; fuzzy logic

MML identifier: FUZIMPL2, version: 8.1.08 5.53.1335

0. INTRODUCTION

There are two fundamental aims of this Mizar article: first of all, I wanted to introduce in the Mizar Mathematical Library how nine basic fuzzy implications formally defined in [4] are ordered – and this result is given in Section 2 as a formal counterpart of Example 1.1.6, p. 3 of [1].

On the other hand, in the final section I prove the formal characterization of fundamental fuzzy implications in terms of four elementary properties [12] expressed in Table 1.4 of [1], p. 10 (note the absence of the continuity of the operators in our version of this presentation). Here

- (NP) – the left neutrality property,
- (EP) – the exchange principle,
- (IP) – the identity principle,
- (OP) – the ordering property.

Actually, this is the part of Example 1.3.2, p. 9 from [1]:

Fuzzy implication	(NP)	(EP)	(IP)	(OP)
I_{LK}	+	+	+	+
I_{GD}	+	+	+	+
I_{RC}	+	+	–	–
I_{KD}	+	+	–	–
I_{GG}	+	+	+	+
I_{RS}	–	–	+	+
I_{YG}	+	+	–	–
I_{WB}	+	+	+	–
I_{FD}	+	+	+	+

Additionally, Section 4 contains registrations of clusters of adjectives allowing for further work in more automated framework within fuzzy sets [3] – this is the Mizar version of Lemma 1.3.3 and 1.3.4 from [1]. Such automatization can be especially useful in the hybridization of fuzzy and rough approaches [7].

1. PRELIMINARIES

We introduce the notation I_{LK} as a synonym of the Łukasiewicz implication and I_{GD} as a synonym of the Gödel implication. We introduce I_{RC} as a synonym of the Reichenbach implication and I_{KD} as a synonym of the Kleene-Dienes implication.

We introduce I_{GG} as a synonym of the Goguen implication and I_{RS} as a synonym of the Rescher implication. We introduce I_{YG} as a synonym of the Yager implication and I_{WB} as a synonym of the Weber implication and I_{FD} as a synonym of the Fodor implication.

From now on x, y denote elements of $[0, 1]$. Now we state the propositions:

- (1) $\square^1 = (\text{AffineMap}(1, 0)) \upharpoonright]0, +\infty[$.

PROOF: Set $f = \square^1$. Set $g = (\text{AffineMap}(1, 0)) \upharpoonright]0, +\infty[$. For every object x such that $x \in \text{dom } f$ holds $f(x) = g(x)$. \square

- (2) Let us consider real numbers a, b . Then

- (i) $\text{AffineMap}(a, b)$ is differentiable on \mathbb{R} , and
 - (ii) for every real number x , $(\text{AffineMap}(a, b))'(x) = a$.
- (3) If $0 < x < 1$ and $0 < y < 1$, then $(\square^x + (\text{AffineMap}(-x, x - 1))) \upharpoonright]0, 1[$ is increasing.
- PROOF: Set $f_1 = \square^x$. Set $f_2 = \text{AffineMap}(-x, x - 1)$. Reconsider $Y =]0, 1[$ as an open subset of \mathbb{R} . Set $f = f_1 + f_2$. Set $A =]0, +\infty[$. f_2 is differentiable on A . $f_1 \upharpoonright A$ is differentiable on A . f_2 is differentiable on Y . For every real number y such that $y \in Y$ holds $0 < f'(y)$ by [11, (21)], (2). \square
- (4) Let us consider a real number u . Suppose $u \in]0, 1[$. Then $(\square^x + (\text{AffineMap}(-x, x - 1)))(u) = u^x - 1 + x - x \cdot u$.

2. THE ORDERING OF FUZZY IMPLICATIONS

Now we state the propositions:

- (5) (i) if $x \leq y$, then $(I_{LK})(x, y) = 1$, and
- (ii) if $x > y$, then $(I_{LK})(x, y) = 1 - x + y$.
- (6) (i) if $x = 0$, then $(I_{GG})(x, y) = 1$, and
- (ii) if $x > 0$, then $(I_{GG})(x, y) = \min(1, \frac{y}{x})$.
- (7) $I_{KD} \leq I_{RC} \leq I_{LK} \leq I_{WB}$.
- (8) $I_{RS} \leq I_{GD} \leq I_{GG} \leq I_{LK} \leq I_{WB}$.
- (9) $I_{RC} \leq I_{LK} \leq I_{WB}$.
- (10) $I_{KD} \leq I_{FD} \leq I_{LK} \leq I_{WB}$.
- (11) $I_{RS} \leq I_{GD} \leq I_{FD} \leq I_{LK} \leq I_{WB}$.

3. ADDITIONAL PROPERTIES OF FUZZY IMPLICATIONS

Let I be a binary operation on $[0, 1]$. We say that I satisfies (NP) if and only if

- (Def. 1) for every element y of $[0, 1]$, $I(1, y) = y$.

We say that I satisfies (EP) if and only if

- (Def. 2) for every elements x, y, z of $[0, 1]$, $I(x, I(y, z)) = I(y, I(x, z))$.

We say that I satisfies (IP) if and only if

- (Def. 3) for every element x of $[0, 1]$, $I(x, x) = 1$.

We say that I satisfies (OP) if and only if

- (Def. 4) for every elements x, y of $[0, 1]$, $I(x, y) = 1$ iff $x \leq y$.

In the sequel I denotes a binary operation on $[0, 1]$.

Let I be a binary operation on $[0, 1]$. We introduce the notation I satisfies (NC) as a synonym of I is 01-dominant and I satisfies (I1) as a synonym of I is antitone w.r.t. 1st coordinate.

We introduce I satisfies (I2) as a synonym of I is isotone w.r.t. 2nd coordinate and I satisfies (I3) as a synonym of I is 00-dominant and I satisfies (I4) as a synonym of I is 11-dominant and I satisfies (I5) as a synonym of I is 10-weak.

4. DEPENDENCIES BETWEEN CHOSEN PROPERTIES

Now we state the proposition:

(12) If I satisfies (LB), then I satisfies (I3) and (NC).

One can verify that every binary operation on $[0, 1]$ which satisfies (LB) satisfies also (I3) and (NC).

Now we state the proposition:

(13) If I satisfies (RB), then I satisfies (I4) and (NC).

One can check that every binary operation on $[0, 1]$ which satisfies (RB) satisfies also (I4) and (NC).

Now we state the proposition:

(14) If I satisfies (NP), then I satisfies (I4) and (I5).

Note that every binary operation on $[0, 1]$ which satisfies (NP) satisfies also (I4) and (I5).

Now we state the proposition:

(15) If I satisfies (IP), then I satisfies (I3) and (I4).

Let us note that every binary operation on $[0, 1]$ which satisfies (IP) satisfies also (I3) and (I4).

Now we state the proposition:

(16) If I satisfies (OP), then I satisfies (I3), (I4), (NC), (LB), (RB), and (IP).

One can verify that every binary operation on $[0, 1]$ which satisfies (OP) satisfies also (I3), (I4), (NC), (LB), (RB), and (IP).

Now we state the proposition:

(17) If I satisfies (EP) and (OP), then I satisfies (I1), (I3), (I4), (I5), (LB), (RB), (NC), (NP), and (IP).

One can verify that every binary operation on $[0, 1]$ which satisfies (EP) and (OP) satisfies also (I1), (I5), and (NP).

5. PROPERTIES OF NINE CLASSICAL FUZZY IMPLICATIONS

Let us note that I_{LK} satisfies (NP), (EP), (IP), and (OP).

I_{GD} satisfies (NP), (EP), (IP), and (OP).

I_{RC} satisfies (NP) and (EP) but does not satisfy (IP) and (OP).

I_{KD} satisfies (NP) and (EP) but does not satisfy (IP) and (OP).

I_{GG} satisfies (NP), (EP), (IP), and (OP).

Let us note that I_{RS} satisfies (IP) and (OP) but does not satisfy (NP) and (EP).

I_{YG} satisfies (NP) and (EP) but does not satisfy (IP) and (OP).

I_{WB} satisfies (NP), (EP), and (IP) but does not satisfy (OP).

I_{FD} satisfies (NP), (EP), (IP), and (OP).

I_0 satisfies (EP) but does not satisfy (NP), (IP), and (OP).

I_1 satisfies (EP) and (IP) but does not satisfy (NP) and (OP).

REFERENCES

- [1] Michał Baczyński and Balasubramanian Jayaram. *Fuzzy Implications*. Springer Publishing Company, Incorporated, 2008. doi:10.1007/978-3-540-69082-5.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Małuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.
- [3] Didier Dubois and Henri Prade. *Fuzzy Sets and Systems: Theory and Applications*. Academic Press, New York, 1980.
- [4] Adam Grabowski. Formal introduction to fuzzy implications. *Formalized Mathematics*, 25(3):241–248, 2017. doi:10.1515/forma-2017-0023.
- [5] Adam Grabowski. Basic formal properties of triangular norms and conorms. *Formalized Mathematics*, 25(2):93–100, 2017. doi:10.1515/forma-2017-0009.
- [6] Adam Grabowski. On the computer certification of fuzzy numbers. In M. Ganzha, L. Maciaszek, and M. Paprzycki, editors, *2013 Federated Conference on Computer Science and Information Systems (FedCSIS)*, Federated Conference on Computer Science and Information Systems, pages 51–54, 2013.
- [7] Adam Grabowski and Takashi Mitsuishi. Initial comparison of formal approaches to fuzzy and rough sets. In Leszek Rutkowski, Marcin Korytkowski, Rafal Scherer, Ryszard Tadeusiewicz, Lotfi A. Zadeh, and Jacek M. Zurada, editors, *Artificial Intelligence and Soft Computing - 14th International Conference, ICAISC 2015, Zakopane, Poland, June 14-18, 2015, Proceedings, Part I*, volume 9119 of *Lecture Notes in Computer Science*, pages 160–171. Springer, 2015. doi:10.1007/978-3-319-19324-3_15.
- [8] Adam Grabowski, Artur Korniłowicz, and Adam Naumowicz. Four decades of Mizar. *Journal of Automated Reasoning*, 55(3):191–198, 2015. doi:10.1007/s10817-015-9345-1.
- [9] Petr Hájek. *Metamathematics of Fuzzy Logic*. Dordrecht: Kluwer, 1998.
- [10] Takashi Mitsuishi, Noboru Endou, and Yasunari Shidama. The concept of fuzzy set and membership function and basic properties of fuzzy set operation. *Formalized Mathematics*, 9(2):351–356, 2001.
- [11] Yasunari Shidama. The Taylor expansions. *Formalized Mathematics*, 12(2):195–200, 2004.
- [12] Philippe Smets and Paul Magrez. Implication in fuzzy logic. *International Journal of Approximate Reasoning*, 1(4):327–347, 1987. doi:10.1016/0888-613X(87)90023-5.

- [13] Lotfi Zadeh. Fuzzy sets. *Information and Control*, 8(3):338–353, 1965. doi:10.1016/S0019-9958(65)90241-X.

Accepted September 29, 2018

Zariski Topology

Yasushige Watase
Suginami-ku Matsunoki
3-21-6 Tokyo, Japan

Summary. We formalize in the Mizar system [3], [4] basic definitions of commutative ring theory such as prime spectrum, nilradical, Jacobson radical, local ring, and semi-local ring [5], [6], then formalize proofs of some related theorems along with the first chapter of [1].

The article introduces the so-called Zariski topology. The set of all prime ideals of a commutative ring A is called the prime spectrum of A denoted by Spectrum A . A new functor Spec generates Zariski topology to make Spectrum A a topological space. A different role is given to Spec as a map from a ring morphism of commutative rings to that of topological spaces by the following manner: for a ring homomorphism $h : A \rightarrow B$, we defined $(\text{Spec } h) : \text{Spec } B \rightarrow \text{Spec } A$ by $(\text{Spec } h)(\mathfrak{p}) = h^{-1}(\mathfrak{p})$ where $\mathfrak{p} \in \text{Spec } B$.

MSC: 14A05 16D25 68T99 03B35

Keywords: prime spectrum; local ring; semi-local ring; nilradical; Jacobson radical; Zariski topology

MML identifier: TOPZARI1, version: 8.1.08 5.53.1335

1. PRELIMINARIES: SOME PROPERTIES OF IDEALS

From now on R denotes a commutative ring, A, B denote non degenerated, commutative rings, h denotes a function from A into B , I, I_1, I_2 denote ideals of A , J, J_1, J_2 denote proper ideals of A , p denotes a prime ideal of A .

S denotes non empty subset of A , E, E_1, E_2 denote subsets of A , a, b, f denote elements of A , n denotes a natural number, and x denotes object.

Let us consider A and S . The functor $\text{Ideals}(A, S)$ yielding a subset of Ideals A is defined by the term

(Def. 1) $\{I, \text{ where } I \text{ is an ideal of } A : S \subseteq I\}$.

Let us observe that $\text{Ideals}(A, S)$ is non empty.

Now we state the proposition:

- (1) $\text{Ideals}(A, S) = \text{Ideals}(A, S\text{-ideal})$.

PROOF: $\text{Ideals}(A, S) \subseteq \text{Ideals}(A, S\text{-ideal})$. Consider y being an ideal of A such that $x = y$ and $S\text{-ideal} \subseteq y$. \square

Let A be a unital, non empty multiplicative loop with zero structure and a be an element of A . We say that a is nilpotent if and only if

- (Def. 2) there exists a non zero natural number k such that $a^k = 0_A$.

Let us note that 0_A is nilpotent and there exists an element of A which is nilpotent.

Let us consider A . Observe that 1_A is non nilpotent.

Let us consider f . The functor $\text{MultClSet}(f)$ yielding a subset of A is defined by the term

- (Def. 3) the set of all f^i where i is a natural number.

Let us observe that $\text{MultClSet}(f)$ is multiplicatively closed.

Now we state the propositions:

- (2) Let us consider a natural number n . Then $(1_A)^n = 1_A$.

PROOF: Define $\mathcal{P}[\text{natural number}] \equiv (1_A)^{\mathfrak{S}_1} = 1_A$. For every natural number n , $\mathcal{P}[n]$. \square

- (3) $1_A \notin \sqrt{J}$. The theorem is a consequence of (2).

- (4) $\text{MultClSet}(1_A) = \{1_A\}$. The theorem is a consequence of (2).

Let us consider A, J , and f . The functor $\text{Ideals}(A, J, f)$ yielding a subset of $\text{Ideals } A$ is defined by the term

- (Def. 4) $\{I, \text{ where } I \text{ is a subset of } A : I \text{ is a proper ideal of } A \text{ and } J \subseteq I \text{ and } I \cap \text{MultClSet}(f) = \emptyset\}$.

Let us consider A, J , and f . Now we state the propositions:

- (5) If $f \notin \sqrt{J}$, then $J \in \text{Ideals}(A, J, f)$.

- (6) If $f \notin \sqrt{J}$, then $\text{Ideals}(A, J, f)$ has the upper Zorn property w.r.t. $\subseteq_{\text{Ideals}(A, J, f)}$.

PROOF: Set $S = \text{Ideals}(A, J, f)$. Set $P = \subseteq_S$. For every set Y such that $Y \subseteq S$ and $P \upharpoonright^2 Y$ is a linear order there exists a set x such that $x \in S$ and for every set y such that $y \in Y$ holds $\langle y, x \rangle \in P$. \square

- (7) If $f \notin \sqrt{J}$, then there exists a prime ideal m of A such that $f \notin m$ and $J \subseteq m$.

PROOF: Set $S = \text{Ideals}(A, J, f)$. Set $P = \subseteq_S$. Consider I being a set such that I is maximal in P . Consider p being a subset of A such that $p = I$ and p is a proper ideal of A and $J \subseteq p$ and $p \cap \text{MultClSet}(f) = \emptyset$. p is a quasi-prime ideal of A . \square

(8) There exists a maximal ideal m of A such that $J \subseteq m$.

PROOF: $1_A \notin \sqrt{J}$. Set $S = \text{Ideals}(A, J, 1_A)$. Set $P = \subseteq_S$. Consider I being a set such that I is maximal in P . Consider p being a subset of A such that $p = I$ and p is a proper ideal of A and $J \subseteq p$ and $p \cap \text{MultClSet}(1_A) = \emptyset$. For every ideal q of A such that $p \subseteq q$ holds $q = p$ or q is not proper. \square

(9) There exists a prime ideal m of A such that $J \subseteq m$. The theorem is a consequence of (8).

(10) If a is a non-unit of A , then there exists a maximal ideal m of A such that $a \in m$. The theorem is a consequence of (8).

2. SPECTRUM OF PRIME IDEALS (SPECTRUM) AND MAXIMAL IDEALS (M-SPECTRUM)

Let R be a commutative ring. The spectrum of R yielding a family of subsets of R is defined by the term

(Def. 5) $\left\{ \begin{array}{l} \{I, \text{ where } I \text{ is an ideal of } R : I \text{ is quasi-prime and } I \neq \Omega_R\}, \\ \quad \text{if } R \text{ is not degenerated,} \\ \emptyset, \text{ otherwise.} \end{array} \right.$

Let us consider A . Observe that the spectrum of A yields a family of subsets of A and is defined by the term

(Def. 6) the set of all I where I is a prime ideal of A .

Observe that the spectrum of A is non empty.

Let us consider R . The functor $\text{m-Spectrum}(R)$ yielding a family of subsets of R is defined by the term

(Def. 7) $\left\{ \begin{array}{l} \{I, \text{ where } I \text{ is an ideal of } R : I \text{ is quasi-maximal and } I \neq \Omega_R\}, \\ \quad \text{if } R \text{ is not degenerated,} \\ \emptyset, \text{ otherwise.} \end{array} \right.$

Let us consider A . Observe that the functor $\text{m-Spectrum}(A)$ yields a family of subsets of the carrier of A and is defined by the term

(Def. 8) the set of all I where I is a maximal ideal of A .

Observe that $\text{m-Spectrum}(A)$ is non empty.

3. LOCAL AND SEMI-LOCAL RING

Let us consider A . We say that A is local if and only if

(Def. 9) there exists an ideal m of A such that $\text{m-Spectrum}(A) = \{m\}$.

We say that A is semi-local if and only if

(Def. 10) $\mathfrak{m}\text{-Spectrum}(A)$ is finite.

Now we state the propositions:

(11) If $x \in I$ and I is a proper ideal of A , then x is a non-unit of A .

(12) If for every objects m_1, m_2 such that $m_1, m_2 \in \mathfrak{m}\text{-Spectrum}(A)$ holds $m_1 = m_2$, then A is local.

(13) If for every x such that $x \in \Omega_A \setminus J$ holds x is a unit of A , then A is local. The theorem is a consequence of (8), (11), and (12).

In the sequel m denotes a maximal ideal of A . Now we state the propositions:

(14) If $a \in \Omega_A \setminus m$, then $\{a\}$ -ideal $+ m = \Omega_A$.

(15) If for every a such that $a \in m$ holds $1_A + a$ is a unit of A , then A is local.

PROOF: For every x such that $x \in \Omega_A \setminus m$ holds x is a unit of A . \square

Let us consider R . Let E be a subset of R . The functor $\text{PrimeIdeals}(R, E)$ yielding a subset of the spectrum of R is defined by the term

(Def. 11)
$$\begin{cases} \{p, \text{ where } p \text{ is an ideal of } R : p \text{ is quasi-prime and } p \neq \Omega_R \text{ and } E \subseteq p\}, \\ \quad \text{if } R \text{ is not degenerated,} \\ \emptyset, \text{ otherwise.} \end{cases}$$

Let us consider A . Let E be a subset of A . Let us note that the functor $\text{PrimeIdeals}(A, E)$ yields a subset of the spectrum of A and is defined by the term

(Def. 12) $\{p, \text{ where } p \text{ is a prime ideal of } A : E \subseteq p\}$.

Let us consider J . Observe that $\text{PrimeIdeals}(A, J)$ is non empty.

From now on p denotes a prime ideal of A and k denotes a non zero natural number. Now we state the proposition:

(16) If $a \notin p$, then $a^k \notin p$.

4. NILRADICAL AND JACOBSON RADICAL

Let us consider A . The functor $\text{nilrad}(A)$ yielding a subset of A is defined by the term

(Def. 13) the set of all a where a is a nilpotent element of A .

Now we state the proposition:

(17) $\text{nilrad}(A) = \sqrt{\{0_A\}}$.

Let us consider A . One can verify that $\text{nilrad}(A)$ is non empty and $\text{nilrad}(A)$ is closed under addition as a subset of A and $\text{nilrad}(A)$ is left and right ideal as a subset of A .

Now we state the propositions:

- (18) $\sqrt{J} = \bigcap \text{PrimeIdeals}(A, J)$. The theorem is a consequence of (16), (7), and (9).
- (19) $\text{nilrad}(A) = \bigcap (\text{the spectrum of } A)$. The theorem is a consequence of (17) and (18).
- (20) $I \subseteq \sqrt{I}$.
- (21) If $I \subseteq J$, then $\sqrt{I} \subseteq \sqrt{J}$.

PROOF: Consider s_1 being an element of A such that $s_1 = s$ and there exists an element n of \mathbb{N} such that $s_1^n \in I$. Consider n_1 being an element of \mathbb{N} such that $s_1^{n_1} \in I$. $n_1 \neq 0$ by [7, (8)], [2, (19)]. \square

Let us consider A . The functor $J\text{-Rad}(A)$ yielding a subset of A is defined by the term

(Def. 14) $\bigcap \text{m-Spectrum}(A)$.

5. CONSTRUCTION OF ZARISKI TOPOLOGY OF THE PRIME SPECTRUM OF A

Now we state the propositions:

- (22) $\text{PrimeIdeals}(A, S) \subseteq \text{Ideals}(A, S)$.
 - (23) $\text{PrimeIdeals}(A, S) = \text{Ideals}(A, S) \cap (\text{the spectrum of } A)$. The theorem is a consequence of (22).
 - (24) $\text{PrimeIdeals}(A, S) = \text{PrimeIdeals}(A, S\text{-ideal})$. The theorem is a consequence of (23) and (1).
 - (25) If $I \subseteq p$, then $\sqrt{I} \subseteq p$.
- PROOF: Consider s_1 being an element of A such that $s_1 = s$ and there exists an element n of \mathbb{N} such that $s_1^n \in I$. Consider n_1 being an element of \mathbb{N} such that $s_1^{n_1} \in I$. $n_1 \neq 0$. \square
- (26) If $\sqrt{I} \subseteq p$, then $I \subseteq p$. The theorem is a consequence of (20).
 - (27) $\text{PrimeIdeals}(A, \sqrt{S\text{-ideal}}) = \text{PrimeIdeals}(A, S\text{-ideal})$. The theorem is a consequence of (26) and (25).
 - (28) If $E_2 \subseteq E_1$, then $\text{PrimeIdeals}(A, E_1) \subseteq \text{PrimeIdeals}(A, E_2)$.
 - (29) $\text{PrimeIdeals}(A, J_1) = \text{PrimeIdeals}(A, J_2)$ if and only if $\sqrt{J_1} = \sqrt{J_2}$. The theorem is a consequence of (18) and (27).
 - (30) If $I_1 * I_2 \subseteq p$, then $I_1 \subseteq p$ or $I_2 \subseteq p$.

PROOF: If it is not true that $I_1 \subseteq p$ or $I_2 \subseteq p$, then $I_1 * I_2 \not\subseteq p$. \square

- (31) $\text{PrimeIdeals}(A, \{1_A\}) = \emptyset$.
- (32) The spectrum of $A = \text{PrimeIdeals}(A, \{0_A\})$.
- (33) Let us consider non empty subsets E_1, E_2 of A . Then there exists a non empty subset E_3 of A such that $\text{PrimeIdeals}(A, E_1) \cup \text{PrimeIdeals}(A, E_2) = \text{PrimeIdeals}(A, E_3)$.

PROOF: Set $I_1 = E_1$ -ideal. Set $I_2 = E_2$ -ideal. Reconsider $I_3 = I_1 * I_2$ as an ideal of A . $\text{PrimeIdeals}(A, E_1) = \text{PrimeIdeals}(A, I_1)$. $\text{PrimeIdeals}(A, I_3) \subseteq \text{PrimeIdeals}(A, I_1) \cup \text{PrimeIdeals}(A, I_2)$. $\text{PrimeIdeals}(A, I_1) \cup \text{PrimeIdeals}(A, I_2) \subseteq \text{PrimeIdeals}(A, I_3)$. $\text{PrimeIdeals}(A, I_3) = \text{PrimeIdeals}(A, E_1) \cup \text{PrimeIdeals}(A, E_2)$. \square

- (34) Let us consider a family G of subsets of the spectrum of A . Suppose for every set S such that $S \in G$ there exists a non empty subset E of A such that $S = \text{PrimeIdeals}(A, E)$. Then there exists a non empty subset F of A such that $\text{Intersect}(G) = \text{PrimeIdeals}(A, F)$. The theorem is a consequence of (28).

Let us consider A . The functor $\text{Spec}(A)$ yielding a strict topological space is defined by

- (Def. 15) the carrier of $it =$ the spectrum of A and for every subset F of it , F is closed iff there exists a non empty subset E of A such that $F = \text{PrimeIdeals}(A, E)$.

Note that $\text{Spec}(A)$ is non empty. Now we state the proposition:

- (35) Let us consider points P, Q of $\text{Spec}(A)$. Suppose $P \neq Q$. Then there exists a subset V of $\text{Spec}(A)$ such that
- (i) V is open, and
 - (ii) $P \in V$ and $Q \notin V$ or $Q \in V$ and $P \notin V$.

Note that there exists a commutative ring which is degenerated. Let R be a degenerated, commutative ring. Let us observe that ADTS (the spectrum of R) is T_0 . Let us consider A . Observe that $\text{Spec}(A)$ is T_0 .

6. CONTINUOUS MAP OF ZARISKI TOPOLOGY ASSOCIATED WITH A RING HOMOMORPHISM

From now on M_0 denotes an ideal of B . Now we state the proposition:

- (36) If h inherits ring homomorphism, then $h^{-1}(M_0)$ is an ideal of A .

In the sequel M_0 denotes a prime ideal of B .

- (37) If h inherits ring homomorphism, then $h^{-1}(M_0)$ is a prime ideal of A .

PROOF: For every elements x, y of A such that $x \cdot y \in h^{-1}(M_0)$ holds $x \in h^{-1}(M_0)$ or $y \in h^{-1}(M_0)$. $h^{-1}(M_0) \neq$ the carrier of A . \square

Let us consider A, B , and h . Assume h inherits ring homomorphism. The functor $\text{Spec}(h)$ yielding a function from $\text{Spec}(B)$ into $\text{Spec}(A)$ is defined by

- (Def. 16) for every point x of $\text{Spec}(B)$, $it(x) = h^{-1}(x)$.

Now we state the propositions:

(38) If h inherits ring homomorphism, then $\text{Spec}(h)^{-1} \text{PrimeIdeals}(A, E) = \text{PrimeIdeals}(B, h^\circ E)$.

PROOF: $\text{Spec}(h)^{-1} \text{PrimeIdeals}(A, E) \subseteq \text{PrimeIdeals}(B, h^\circ E)$. Consider q being a prime ideal of B such that $x = q$ and $h^\circ E \subseteq q$. $h^{-1}(q)$ is a prime ideal of A . \square

(39) If h inherits ring homomorphism, then $\text{Spec}(h)$ is continuous. The theorem is a consequence of (38).

REFERENCES

- [1] Michael Francis Atiyah and Ian Grant Macdonald. *Introduction to Commutative Algebra*, volume 2. Addison-Wesley Reading, 1969.
- [2] Jonathan Backer, Piotr Rudnicki, and Christoph Schwarzweiler. Ring ideals. *Formalized Mathematics*, 9(3):565–582, 2001.
- [3] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8_17.
- [4] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, and Karol Pąk. The role of the Mizar Mathematical Library for interactive proof development in Mizar. *Journal of Automated Reasoning*, 61(1):9–32, 2018. doi:10.1007/s10817-017-9440-6.
- [5] Shigeru Iitaka. *Algebraic Geometry: An Introduction to Birational Geometry of Algebraic Varieties*. Springer-Verlag New York, Inc., 1982.
- [6] Shigeru Iitaka. *Ring Theory (in Japanese)*. Kyoritsu Shuppan Co., Ltd., 2013.
- [7] Christoph Schwarzweiler. The binomial theorem for algebraic structures. *Formalized Mathematics*, 9(3):559–564, 2001.

Accepted October 16, 2018
